

Facet Joint Intervention	
Order Details	
Order Date *	MM/DD/YYYY
Procedure ordered *	☐ 64490 - Inj paravert f jnt c/t 1 lev ☐ 64493 - Inj paravert f jnt l/s 1 lev ☐ 64633 - Destroy cerv/thor facet jnt ☐ 64635 - Destroy lumb/sac facet jnt
Order Id *	Type a value
Ordering Physician Name *	Type a value
Ordering Physician NPI *	Type a value
Requested Intervention Details	
What is the primary HCPC code for the requested procedure? *	☐ 64490 - Inj paravert f jnt c/t 1 lev ☐ 64493 - Inj paravert f jnt l/s 1 lev ☐ 64633 - Destroy cerv/thor facet jnt ☐ 64635 - Destroy lumb/sac facet jnt
How many additional levels are being requested? *	☐ None, Primary only ☐ One additional level: 64491 - Inj paravert f jnt c/t 2 lev. (Primary level, and second level.) Reminder: one to two levels, either unilateral or bilateral are allowed per session per spine region. ☐ Two additional levels: 64492 - Inj paravert f jnt c/t 3 lev. (Primary level, second level and a third level.) Provider education: Three or four levels is not medically necessary and not covered. Reminder: one to two levels, either unilateral or bilateral are allowed per session per spine region. ☐ Three additional levels. (Primary level, second level, third level and a fourth level.) Provider education: Three or four levels is not medically necessary and not covered. Reminder: one to two levels, either unilateral or bilateral are allowed per session per spine region.
Specify the additional level being requested: *	Type a value
How many additional levels are being requested? *	☐ None, Primary only ☐ One additional level: 64494 - Inj paravert f jnt l/s 2 lev. (Primary level, and second level.) Reminder: one to two levels, unilateral or bilateral are allowed per session per spine region. ☐ Two additional levels: 64495 - Inj paravert f jnt l/s 3 lev. (Primary level, second level and a third level.) Provider education: Three or four levels is not medically necessary and not covered. Reminder: one to two levels, either unilateral or bilateral are allowed per session per spine region. ☐ Three additional levels. (Primary level, second level, third leve and a fourth level.) Provider education: Three or four levels is not medically necessary and not covered. Reminder: one to two levels, either unilateral or bilateral are allowed per session per spine region

be reimbursed per rolling 12 months. *

If RFA is being performed what were the qualifying MBB first injection and MBB second injection dates?

Medial branch block (MBB) first injection date:

MM/DD/YYYY

Medial branch block (MBB) second injection date:

MM/DD/YYYY

Provider Education: For each covered spinal region no more than two (2) radiofrequency sessions will be reimbursed per rolling 12 months. *

Yes

No

Not Answered

Provider Education: Cyst aspiration/rupture may be repeated once per individual cyst and only if there is 50% or more consistent improvement in pain for at least three (3) months. *

Yes

No

Not Answered

Are there treatments such as anticoagulation interruption of therapy that would support consideration for additional levels?

Yes

No

Not Answered

Is general anesthesia, moderate sedation, or monitored anesthesia care going to be used?

Yes

No

Not Answered

What is the specific condition or symptoms that make sedation medically necessary? Select all that are well documented.

Patient has a medical condition that would interfere with the procedure.

Patient does not have the cognitive capacity to cooperate.

Patient does not have the functional capacity to cooperate.

Patient does not have the capacity to remain motionless during the procedure.

Patient is not able to tolerate remaining in a painful position for a prolonged period of time.

Other

Specify other

Type a value

If the requested intervention will not meet the medical necessity criteria for anesthesia/sedation, was an advance beneficiary notice of non-coverage (ABN) given to the patient?

Yes

No

Not Answered

Is this request an expedited request?

Yes

No

Not Answered

What is the specific condition that demonstrates any delays can jeopardize the patient? Specify:

Type a value

What is the date for the requested expedited procedure to be performed?

MM/DD/YYYY

Will fluoroscopy or CT Guidance be used during this intervention?

Yes

No

Not Answered

What FDA approved agents will be injected? Select all that apply.

Contrast agent

Corticosteroids

Anti-inflammatory

Anesthetic

Anesthetic injected:

Procaine

Lidocaine

Tetracaine

Prilocaine

Etidocaine

Bupivacaine

Ropivacaine

Does the patient have an implanted electrical device that requires consideration?

Yes

No

Not Answered

Provider has/will follow manufacturer instructions to ensure patient's safety relevant to the planned procedure?

Yes

No

Not Answered

Diagnosis(es) and Hx of Relevant Procedures

The patient's primary diagnosis that supports approval for a facet joint intervention procedure.

Cervical/Thoracic

Spondylosis without myelopathy or radiculopathy, cervical region

Spondylosis without myelopathy or radiculopathy, cervicothoracic region

Spondylosis without myelopathy or radiculopathy, thoracic region

Spondylosis without myelopathy or radiculopathy, thoracolumbar region

Other spondylosis, cervical region

Other spondylosis, cervicothoracic region

Other spondylosis, thoracic region

Other spondylosis, thoracolumbar region

Ankylosing hyperostosis [Forestier], cervical region

Ankylosing hyperostosis [Forestier], cervicothoracic region

Ankylosing hyperostosis [Forestier], thoracic region

Ankylosing hyperostosis [Forestier], thoracolumbar region

Other

Specify other *

Type a value

Lumbar/Sacral

Spondylosis without myelopathy or radiculopathy, lumbar region

Spondylosis without myelopathy or radiculopathy, lumbosacral region

Other spondylosis, lumbar region

Other spondylosis, lumbosacral region

Ankylosing hyperostosis [Forestier], lumbar region

Ankylosing hyperostosis [Forestier], lumbosacral region

Other

Specify other *

Type a value

The patient's primary diagnosis that supports approval and allowed for facet cyst rupture procedures only.

Cervical/Thoracic *

Other bursal cyst, unspecified site

Other bursal cyst, other site

Other specified dorsopathies

Other specified dorsopathies, cervical

Other specified dorsopathies, cervicothoracic

Other specified dorsopathies, thoracic

Other specified dorsopathies, thoracolumbar

Other

Specify other *

Type a value

Lumbar/Sacral *

Other bursal cyst, unspecified site

Other bursal cyst, other site

Other specified dorsopathies

Other specified dorsopathies, lumbar

Other specified dorsopathies, lumbosacral

Other specified dorsopathies, sacral

Other

Specify other *

Type a value

Other relevant diagnosis or supported findings:

Cervical/Thoracic

Spondylosis without myelopathy or radiculopathy, cervical region

Spondylosis without myelopathy or radiculopathy, cervicothoracic region

Spondylosis without myelopathy or radiculopathy, thoracic region

	☐ Spondylosis without myelopathy or radiculopathy, thoracolumbar region ☐ Other spondylosis, cervical region ☐ Other spondylosis, cervicothoracic region ☐ Other spondylosis, thoracic region ☐ Other spondylosis, thoracolumbar region ☐ Ankylosing hyperostosis [Forestier], cervical region ☐ Ankylosing hyperostosis [Forestier], cervicothoracic region ☐ Ankylosing hyperostosis [Forestier], thoracic region ☐ Ankylosing hyperostosis [Forestier], thoracolumbar region ☐ Other
Specify other *	Type a value
Lumbar/Sacral	☐ Spondylosis without myelopathy or radiculopathy, lumbar region ☐ Spondylosis without myelopathy or radiculopathy, lumbosacral region ☐ Other spondylosis, lumbar region ☐ Other spondylosis, lumbosacral region ☐ Ankylosing hyperostosis [Forestier], lumbar region ☐ Ankylosing hyperostosis [Forestier], lumbosacral region ☐ Other
Specify other *	Type a value
For the requested therapeutic intra-articular [IA] injection(s), select the Diagnosis(es) that explain anatomic restrictions and why an MBB cannot be performed. *	☐ Posterior Fusion ☐ Scoliosis ☐ Pseudoarthrosis ☐ Presence of a sixth lumbar vertebra ☐ Sacralization of the fifth lumbar vertebra ☐ Severe Osteophytes (bone spurs) ☐ Obesity or excessive adipose tissue ☐ Severe Kyphosis ☐ Prior surgical scarring ☐ Other
Specify other	Type a value
Does the patient record reflect accurate/active ICD10 codes, eg., old diagnosis(es) have been coded to 'history of' if no longer active/being treated?	☐ Yes ☐ No ☒ Not Answered
Non-facet pathology per clinical assessment or diagnostic studies has ruled out other sources of the patient's pain, including but not limited to fractures, tumors, infections or significant deformity.	☐ Negative for non-facet pathology sources: Medical record indicates all of the following are ABSENT: active spine fractures, spine tumors, active infections, sciatica, hip osteoarthritis, sacroiliac impingement, untreated radiculopathy, neurogenic claudication, myofascial pain, compression fractures, disc herniation, osteophytes, rheumatoid arthritis, fibromyalgia. ☐ Positive for non-facet pathology sources: Medical record indicates one or more of the following are PRESENT: active spine fractures, spine tumors, active infections, sciatica, hip osteoarthritis, sacroiliac impingement, untreated radiculopathy, neurogenic claudication, myofascial pain, compression fractures, disc herniation, osteophytes, rheumatoid arthritis, fibromyalgia.
Hx of relevant spine surgery or procedures:	Type a value
Supporting Clinical Findings	
The supportive findings for the requested procedure, select one.	☐ Neck pain and its progression. ☐ Axial low back pain and its progression.
Is there an initial pain onset encounter date PRIOR to the identified chronic pain baseline encounter, supporting the progression of chronic pain being identified, evaluated, and documented.	MM/DD/YYYY
Which pain scale was used at the initial encounter? Provider education: Pain assessment must be performed and documented at baseline, after each procedure and at each follow-up using the same pain scale for each assessment.	☐ Numerical Pain Rating Scale (NRS) (0-10) ☐ Visual Analog Scale (VAS) Score ☐ Wong-Baker FACES ☐ Verbal Rating Scale for pain intensity ☐ Pain Assessment in Advanced Dementia Scale (PAINAD) ☐ Other ☐ No tool used
Pain duration	Type a number
Pain Intensity or Severity (Pain Score)	Type a number
Which functional assessment (disability) scale was used at the initial encounter if applicable? Provider education: The SAME scale must be used at baseline, after each procedure and at each follow-up as applicable.)	☐ Pain Disability Index ☐ Oswestry Disability Index (ODI) ☐ Oswestry Low Back Pain Disability Questionnaire (OSW) ☐ Quebec Low Back Pain Disability Score (QUE) ☐ Roland Morris Pain Scale ☐ Back Pain Functional Scale (BPFS) ☐ PROMIS profile domains which have been found to correlate well with the ODI scale ☐ No functional assessment tool used
Which functional assessment (disability) scale was used at the initial encounter if applicable? Provider education: The SAME scale must be used at baseline, after each procedure and at each follow-up as applicable.)	☐ Pain Disability Index ☐ Oswestry Disability Index (ODI) ☐ Roland Morris Pain Scale ☐ Back Pain Functional Scale (BPFS) ☐ PROMIS profile domains which have been found to correlate well with the ODI scale ☐ No functional assessment tool used
Score/Results:	Type a number
Pain Intensity Score:	Type a number
Pain Intensity Score:	☐ No pain or can tolerate pain without pain killers ☐ Able to manage pain without pain killers ☐ Complete relief of pain with pain killers ☐ Moderate relief of pain with pain killers ☐ Very little relief of pain with pain killers ☐ No relief/no effect, and does not use due to not helpful
Pain Intensity Score:	☐ No pain ☐ Mild ☐ Moderate ☐ Severe ☐ Very Severe
Disability Index Score/Results:	☐ 0 - 4 No disability ☐ 5 - 14 Mild disability ☐ 15 - 24 Moderate disability ☐ 25 - 34 Severe disability ☐ 35 - 50 Completely disabled
Pain interference with ADLs Score/Results:	Type a number
The baseline encounter date. The date the provider performed the required baseline evaluation for pain that has now been present for three months or greater and not responding to conservative pain management measures.	MM/DD/YYYY
Baseline encounter supportive findings, select one.	☐ Chronic neck pain greater than three months duration. ☐ Chronic low axial back pain greater than three months duration.
Chronic neck pain region	☐ Pain in cervical spine. ☐ Pain in thoracic spine.
Chronic low back pain region	☐ Pain in lumber spine. ☐ Pain in sacral spine.

Which pain scale was used at the baseline visit encounter? Provider education: Pain assessment must be performed and documented at baseline, after each procedure and at each follow-up using the same pain scale for each assessment.	☐ Numerical Pain Rating Scale (NRS) (0-10) ☐ Visual Analog Scale (VAS) Score ☐ Wong-Baker FACES ☐ Verbal Rating Scale for pain intensity ☐ Pain Assessment in Advanced Dementia Scale (PAINAD) ☐ Other ☐ No tool used
Pain duration	Type a number
Pain Intensity or Severity (Pain Score)	Type a number
Which functional assessment (disability) scale was used at the baseline visit? Provider education: The SAME scale must be used at baseline, after each procedure and at each follow-up as applicable.)	☐ Pain Disability Index ☐ Oswestry Disability Index (ODI) ☐ Oswestry Low Back Pain Disability Questionnaire (OSW) ☐ Quebec Low Back Pain Disability Scare (QUE) ☐ Roland Morris Pain Scale ☐ Back Pain Functional Scale (BPFS) ☐ PROMIS profile domains which have been found to correlate well with the ODI scale ☐ No functional assessment tool used
Which functional assessment (disability) scale was used at the baseline visit? Provider education: The SAME scale must be used at baseline, after each procedure and at each follow-up as applicable.)	☐ Pain Disability Index ☐ Oswestry Disability Index (ODI) ☐ Roland Morris Pain Scale ☐ Back Pain Functional Scale (BPFS) ☐ PROMIS profile domains which have been found to correlate well with the ODI scale ☐ No functional assessment tool used
Score/Results:	Type a number
Pain Intensity Score:	Type a number
Pain Intensity Score:	☐ No pain or can tolerate pain without pain killers ☐ Able to manage pain without pain killers ☐ Complete relief of pain with pain killers ☐ Moderate relief of pain with pain killers ☐ Very little relief of pain with pain killers ☐ No relief/no effect, and does not use due to not helpful
Pain Intensity Score:	☐ No pain ☐ Mild ☐ Moderate ☐ Severe ☐ Very Severe
Disability Index Score/Results:	☐ 0 - 4 No disability ☐ 5 - 14 Mild disability ☐ 15 - 24 Moderate disability ☐ 25 - 34 Severe disability ☐ 35 - 50 Completely disabled
Pain interference with ADLs Score/Results:	Type a number
Past pain medications attempted:	☐ Non-opioid analgesic(s) ☐ Opioid analgesic(s) ☐ Nonsteroidal anti-inflammatory drugs (NSAIDs) ☐ Muscle relaxants ☐ Anti-seizure medications ☐ Antidepressants ☐ Other
Specify other:	Type a value
Patient's therapeutic response:	☐ Poor/minimal ☐ Fair response ☐ Good response ☐ Excellent ☐ Complete
Past non-pharmacologic treatments attempted and failed to manage pain. Select all that apply.	☐ Cold Therapy or Heat Therapy ☐ Exercise(s) for pain management ☐ Physical Therapy ☐ Acupuncture ☐ Spinal manipulation ☐ Application of biopsychosocial treatment techniques ☐ Diet ☐ Other
Past non-pharmacologic treatments attempted and failed to manage pain. Select all that apply.	☐ Cold Therapy or Heat Therapy ☐ Exercise(s) for pain management ☐ Acupuncture ☐ Application of biopsychosocial treatment techniques ☐ Diet ☐ Other
Specify other:	Type a value
Patient's therapeutic response:	☐ Poor/minimal ☐ Fair response ☐ Good response ☐ Excellent ☐ Complete
Post Procedure Pain	
Which of the following procedures have been performed and their documented follow up results?	☐ N/A initial request ☐ 1st diagnostic injection follow up ☐ 2nd diagnostic injection follow up ☐ Initial RFA intervention follow up ☐ Second/subsequent RFA intervention follow up ☐ Therapeutic intervention follow up ☐ Facet cyst Aspiration/Rupture pain follow up
What was the date of the 1st diagnostic procedure performed?	MM/DD/YYYY
What was the date of the 1st diagnostic injection pain follow up?	MM/DD/YYYY
What was the date of the 2nd diagnostic procedure performed?	MM/DD/YYYY
What was the date of the 2nd diagnostic injection pain follow up?	MM/DD/YYYY
What was the date of the initial RFA procedure performed?	MM/DD/YYYY
What was the date of the initial RFA intervention pain follow up?	MM/DD/YYYY
What was the date of the Second/subsequent RFA procedure performed?	MM/DD/YYYY
What was the date of the Second/subsequent RFA intervention follow up?	MM/DD/YYYY
What was the date of the therapeutic procedure performed?	MM/DD/YYYY
What was the date of the therapeutic intervention pain follow up?	MM/DD/YYYY
What was the date of the IA procedure for cyst rupture performed?	MM/DD/YYYY
What was the date of the IA injection performed with synovial cyst aspiration pain follow up?	MM/DD/YYYY
Did the patient experienced pain relief results after the first diagnostic procedure that indicates there has been a consistent positive pain relief response of at least 80% ?	☐ Yes ☐ No ☒ Not Answered
Did the patient experienced pain relief results after the 2nd diagnostic procedure that indicates there has been a consistent positive pain relief response of at least 80% ?	☐ Yes ☐ No ☒ Not Answered
Did the patient experience pain relief results after the RFA procedure that indicates there has been a consistent positive pain relief response of at least 80% ?	☐ Yes ☐ No ☒ Not Answered
Did the patient experience pain relief results after the second RFA procedure that indicates there has been a	☐ Yes ☐ No ☒ Not Answered

consistent positive pain relief response of at least 50% improvement for at least six months or at least 50% improvement in ADL performance?	
Did the patient experience pain relief results after the therapeutic procedure that indicates there has been at least consistent 50% pain relief for at least three (3) months from the prior therapeutic procedure or at least 50% consistent improvement in the ability to perform previously painful movements and ADLs as compared to baseline measurement using the same scale? Yes No Not Answered	
The duration of pain relief is consistent with the agent used. Specify the duration?	Type a number
Which pain scale was used at this follow up that reflects the reported pain relief? Provider education: Pain assessment must be performed and documented at baseline, after each procedure and at each follow-up using the same pain scale for each assessment.	Numerical Pain Rating Scale (NRS) (0-10) Visual Analog Scale (VAS) Score Wong-Baker FACES Verbal Rating Scale for pain intensity Pain Assessment in Advanced Dementia Scale (PAINAD) Other No tool used
Pain duration	Type a number
Pain Intensity or Severity (Pain Score)	Type a number
Which functional assessment scale was used at this follow up? Provider education: The SAME scale must be used at baseline, after each procedure and at each follow-up as applicable.)	Pain Disability Index Oswestry Disability Index (ODI) Oswestry Low Back Pain Disability Questionnaire (OSW) Quebec Low Back Pain Disability Scare (QUE) Roland Morris Pain Scale Back Pain Functional Scale (BPFS) PROMIS profile domains which have been found to correlate well with the ODI scale No functional assessment tool used
Score/Results:	Type a number
Pain Intensity Score:	Type a number
Pain Intensity Score:	No pain or can tolerate pain without pain killers Able to manage pain without pain killers Complete relief of pain with pain killers Moderate relief of pain with pain killers Very little relief of pain with pain killers No relief/no effect, and does not use due to not helpful
Pain Intensity Score:	No pain Mild Moderate Severe Very Severe
Disability Index Score/Results:	0 - 4 No disability 5 - 14 Mild disability 15 - 24 Moderate disability 25 - 34 Severe disability 35 - 50 Completely disabled
Pain interference with ADLs Score/Results:	Type a number
History and Physical	
The History and Physical note that supports the procedure:	Type a value
Diagnostic Test Results	
Diagnostic Imaging Report(s) Results available:	Type a value
+ Diagnostic Imaging Report(s) Results available:	