Response Tensors up to 4th rank in C_s

— polar tensors —

$$C^{(0,Q)} = (C^{(0,Q)})$$

$$C^{(0,Q)} = Q_0$$

$$C^{(1,Q)} = \begin{pmatrix} C_x^{(1,Q)} & 0 & C_z^{(1,Q)} \end{pmatrix}$$

$$C_x^{(1,Q)} = Q_x$$
$$C_z^{(1,Q)} = Q_z$$

$$S^{(2,Q)} = \begin{pmatrix} S_{xx}^{(2,Q)} & 0 & S_{xz}^{(2,Q)} \\ 0 & S_{yy}^{(2,Q)} & 0 \\ S_{xz}^{(2,Q)} & 0 & S_{zz}^{(2,Q)} \end{pmatrix}$$

$$S_{xx}^{(2,Q)} = Q_0 - Q_u + Q_v$$

$$S_{xz}^{(2,Q)} = Q_{zx}$$

$$S_{yy}^{(2,Q)} = Q_0 - Q_u - Q_v$$

$$S_{zz}^{(2,Q)} = Q_0 + 2Q_u$$

$$A^{(2,Q)} = \begin{pmatrix} 0 & 0 & A_{xz}^{(2,Q)} \\ 0 & 0 & 0 \\ -A_{xz}^{(2,Q)} & 0 & 0 \end{pmatrix}$$

$$A_{xz}^{(2,Q)} = -G_y$$

$$S^{(3,Q)} = \begin{pmatrix} S_{1x}^{(3,Q)} & 0 & S_{1z}^{(3,Q)} \\ S_{2x}^{(3,Q)} & 0 & S_{2z}^{(3,Q)} \\ S_{3x}^{(3,Q)} & 0 & S_{3z}^{(3,Q)} \\ 0 & S_{4y}^{(3,Q)} & 0 \\ S_{5x}^{(3,Q)} & 0 & S_{5z}^{(3,Q)} \\ 0 & S_{6y}^{(3,Q)} & 0 \end{pmatrix}$$

$$\begin{split} S_{1x}^{(3,Q)} &= 2Q_x[1] + Q_x[2] + 2Q_x^{\alpha} \\ S_{1z}^{(3,Q)} &= -2G_{xy}[1] + Q_z[2] - Q_z^{\alpha} + Q_z^{\beta} \\ S_{2x}^{(3,Q)} &= -2G_{yz}[1] + Q_x[2] - Q_x^{\alpha} + Q_x^{\beta} \\ S_{2z}^{(3,Q)} &= 2G_{xy}[1] + Q_z[2] - Q_z^{\alpha} - Q_z^{\beta} \\ S_{3x}^{(3,Q)} &= 2G_{yz}[1] + Q_x[2] - Q_x^{\alpha} - Q_x^{\beta} \\ S_{3z}^{(3,Q)} &= 2Q_z[1] + Q_z[2] + 2Q_z^{\alpha} \\ S_{4y}^{(3,Q)} &= -G_{xy}[1] + Q_z[1] - Q_z^{\alpha} - Q_z^{\beta} \\ S_{5x}^{(3,Q)} &= G_{xy}[1] + Q_z[1] - Q_x^{\alpha} + Q_z^{\beta} \\ S_{5z}^{(3,Q)} &= -G_{yz}[1] + Q_x[1] - Q_x^{\alpha} - Q_x^{\beta} \\ S_{6y}^{(3,Q)} &= G_{yz}[1] + Q_x[1] - Q_x^{\alpha} + Q_x^{\beta} \end{split}$$

$$A^{(3,Q)} = \begin{pmatrix} 0 & A_{4y}^{(3,Q)} & 0 \\ A_{5x}^{(3,Q)} & 0 & A_{5z}^{(3,Q)} \\ 0 & A_{6y}^{(3,Q)} & 0 \end{pmatrix}$$

$$A_{4y}^{(3,Q)} = G_{xy}[2] + Q_z[3]$$

$$A_{5x}^{(3,Q)} = G_{xy}[2] - Q_z[3]$$

$$A_{5z}^{(3,Q)} = G_{yz}[2] + Q_x[3]$$

$$A_{6y}^{(3,Q)} = G_{yz}[2] - Q_x[3]$$

$$S^{(4,Q)} = \begin{pmatrix} S_{11}^{(4,Q)} & S_{12}^{(4,Q)} & S_{13}^{(4,Q)} & 0 & S_{15}^{(4,Q)} & 0 \\ S_{12}^{(4,Q)} & S_{22}^{(4,Q)} & S_{23}^{(4,Q)} & 0 & S_{25}^{(4,Q)} & 0 \\ S_{13}^{(4,Q)} & S_{23}^{(4,Q)} & S_{33}^{(4,Q)} & 0 & S_{35}^{(4,Q)} & 0 \\ 0 & 0 & 0 & S_{44}^{(4,Q)} & 0 & S_{46}^{(4,Q)} \\ S_{15}^{(4,Q)} & S_{25}^{(4,Q)} & S_{35}^{(4,Q)} & 0 & S_{55}^{(4,Q)} \\ 0 & 0 & 0 & S_{46}^{(4,Q)} & 0 & S_{66}^{(4,Q)} \end{pmatrix}$$

$$\begin{split} S_{11}^{(4,Q)} &= Q_0[1] + 2Q_0[2] - Q_{4u} + Q_{4v} + 2Q_4 - 2Q_u[1] - 4Q_u[2] + 2Q_v[1] + 4Q_v[2] \\ S_{12}^{(4,Q)} &= Q_0[1] + 2Q_{4u} - Q_4 - 2Q_u[1] \\ S_{13}^{(4,Q)} &= Q_0[1] - Q_{4u} - Q_{4v} - Q_4 + Q_u[1] + Q_v[1] \\ S_{15}^{(4,Q)} &= -Q_{4y}^{\alpha} - Q_{4y}^{\beta} + Q_{zx}[1] + 2Q_{zx}[2] \\ S_{22}^{(4,Q)} &= Q_0[1] + 2Q_0[2] - Q_{4u} - Q_{4v} + 2Q_4 - 2Q_u[1] - 4Q_u[2] - 2Q_v[1] - 4Q_v[2] \\ S_{23}^{(4,Q)} &= Q_0[1] - Q_{4u} + Q_{4v} - Q_4 + Q_u[1] - Q_v[1] \\ S_{25}^{(4,Q)} &= 2Q_{4y}^{\beta} + Q_{zx}[1] \\ S_{33}^{(4,Q)} &= Q_0[1] + 2Q_0[2] + 2Q_{4u} + 2Q_4 + 4Q_u[1] + 8Q_u[2] \\ S_{44}^{(4,Q)} &= Q_0[2] - Q_{4u} + Q_{4v} - Q_4 + Q_u[2] - Q_v[2] \\ S_{46}^{(4,Q)} &= 2Q_{4y}^{\beta} + Q_{zx}[2] \\ S_{55}^{(4,Q)} &= Q_0[2] - Q_{4u} - Q_{4v} - Q_4 + Q_u[2] + Q_v[2] \\ S_{6c}^{(4,Q)} &= Q_0[2] + 2Q_{4v} - Q_4 - 2Q_u[2] \end{split}$$

$$\bar{S}^{(4,Q)} = \begin{pmatrix} 0 & \bar{S}^{(4,Q)}_{12} & \bar{S}^{(4,Q)}_{13} & 0 & \bar{S}^{(4,Q)}_{15} & 0 \\ -\bar{S}^{(4,Q)}_{12} & 0 & \bar{S}^{(4,Q)}_{23} & 0 & \bar{S}^{(4,Q)}_{25} & 0 \\ -\bar{S}^{(4,Q)}_{13} & -\bar{S}^{(4,Q)}_{23} & 0 & 0 & \bar{S}^{(4,Q)}_{35} & 0 \\ 0 & 0 & 0 & 0 & \bar{S}^{(4,Q)}_{35} & 0 \\ -\bar{S}^{(4,Q)}_{15} & -\bar{S}^{(4,Q)}_{25} & -\bar{S}^{(4,Q)}_{35} & 0 & 0 & 0 \\ 0 & 0 & 0 & -\bar{S}^{(4,Q)}_{46} & 0 & 0 \end{pmatrix}$$

$$\begin{split} \bar{S}_{12}^{(4,Q)} &= 4G_{xyz}[1] - 2Q_v[3] \\ \bar{S}_{13}^{(4,Q)} &= -4G_{xyz}[1] + 3Q_u[3] - Q_v[3] \\ \bar{S}_{15}^{(4,Q)} &= -2G_y[1] + 2G_y^{\alpha}[1] + 2G_y^{\beta}[1] + Q_{zx}[3] \\ \bar{S}_{23}^{(4,Q)} &= 4G_{xyz}[1] + 3Q_u[3] + Q_v[3] \\ \bar{S}_{25}^{(4,Q)} &= -4G_y^{\beta}[1] + Q_{zx}[3] \\ \bar{S}_{35}^{(4,Q)} &= 2G_y[1] - 2G_y^{\alpha}[1] + 2G_y^{\beta}[1] + Q_{zx}[3] \\ \bar{S}_{46}^{(4,Q)} &= G_y[1] + 4G_y^{\alpha}[1] \end{split}$$

$$A^{(4,Q)} = \begin{pmatrix} A_{xx}^{(4,Q)} & 0 & A_{xz}^{(4,Q)} \\ 0 & A_{yy}^{(4,Q)} & 0 \\ A_{xz}^{(4,Q)} & 0 & A_{zz}^{(4,Q)} \end{pmatrix}$$

$$A_{xx}^{(4,Q)} = Q_0[3] - 2Q_u[6] + 2Q_v[6]$$

$$A_{xz}^{(4,Q)} = 2Q_{zx}[6]$$

$$A_{yy}^{(4,Q)} = Q_0[3] - 2Q_u[6] - 2Q_v[6]$$

$$A_{zz}^{(4,Q)} = Q_0[3] + 4Q_u[6]$$

$$\bar{A}^{(4,Q)} = \begin{pmatrix} 0 & 0 & \bar{A}_{xz}^{(4,Q)} \\ 0 & 0 & 0 \\ -\bar{A}_{xz}^{(4,Q)} & 0 & 0 \end{pmatrix}$$

$$\bar{A}_{xz}^{(4,Q)} = -G_y[6]$$

$$M^{(4,Q)} = \begin{pmatrix} 0 & M_{1y}^{(4,Q)} & 0 \\ 0 & M_{2y}^{(4,Q)} & 0 \\ 0 & M_{3y}^{(4,Q)} & 0 \\ M_{4x}^{(4,Q)} & 0 & M_{4z}^{(4,Q)} \\ 0 & M_{5y}^{(4,Q)} & 0 \\ M_{6x}^{(4,Q)} & 0 & M_{6z}^{(4,Q)} \end{pmatrix}$$

$$\begin{split} M_{1y}^{(4,Q)} &= G_y[3] - G_y^{\alpha}[2] - G_y^{\beta}[2] + 2Q_{zx}[4] \\ M_{2y}^{(4,Q)} &= 2G_y[2] + G_y[3] + 2G_y^{\alpha}[2] \\ M_{3y}^{(4,Q)} &= G_y[3] - G_y^{\alpha}[2] + G_y^{\beta}[2] - 2Q_{zx}[4] \\ M_{4x}^{(4,Q)} &= G_{xyz}[2] - 3Q_u[4] - Q_v[4] \\ M_{4z}^{(4,Q)} &= G_y[2] - G_y^{\alpha}[2] + G_y^{\beta}[2] + Q_{zx}[4] \\ M_{5y}^{(4,Q)} &= G_{xyz}[2] + 3Q_u[4] - Q_v[4] \\ M_{6x}^{(4,Q)} &= G_y[2] - G_y^{\alpha}[2] - G_y^{\beta}[2] - Q_{zx}[4] \\ M_{6x}^{(4,Q)} &= G_{xyz}[2] + 2Q_v[4] \end{split}$$

$$\bar{M}^{(4,Q)} = \begin{pmatrix} 0 & 0 & 0 & \bar{M}_{x4}^{(4,Q)} & 0 & \bar{M}_{x6}^{(4,Q)} \\ \bar{M}_{y1}^{(4,Q)} & \bar{M}_{y2}^{(4,Q)} & \bar{M}_{y3}^{(4,Q)} & 0 & \bar{M}_{y5}^{(4,Q)} & 0 \\ 0 & 0 & 0 & \bar{M}_{z4}^{(4,Q)} & 0 & \bar{M}_{z6}^{(4,Q)} \end{pmatrix}$$

$$\begin{split} \bar{M}_{x4}^{(4,Q)} &= G_{xyz}[3] - 3Q_u[5] - Q_v[5] \\ \bar{M}_{x6}^{(4,Q)} &= G_y[4] - G_y^{\alpha}[3] - G_y^{\beta}[3] - Q_{zx}[5] \\ \bar{M}_{y1}^{(4,Q)} &= G_y[5] - G_y^{\alpha}[3] - G_y^{\beta}[3] + 2Q_{zx}[5] \\ \bar{M}_{y2}^{(4,Q)} &= 2G_y[4] + G_y[5] + 2G_y^{\alpha}[3] \\ \bar{M}_{y3}^{(4,Q)} &= G_y[5] - G_y^{\alpha}[3] + G_y^{\beta}[3] - 2Q_{zx}[5] \\ \bar{M}_{y5}^{(4,Q)} &= G_{xyz}[3] + 3Q_u[5] - Q_v[5] \\ \bar{M}_{z4}^{(4,Q)} &= G_y[4] - G_y^{\alpha}[3] + G_y^{\beta}[3] + Q_{zx}[5] \\ \bar{M}_{z6}^{(4,Q)} &= G_{xyz}[3] + 2Q_v[5] \end{split}$$

$$C^{(1,G)} = \begin{pmatrix} 0 & C_y^{(1,G)} & 0 \end{pmatrix}$$

$$C_y^{(1,G)} = G_y$$

$$S^{(2,G)} = \begin{pmatrix} 0 & S_{xy}^{(2,G)} & 0 \\ S_{xy}^{(2,G)} & 0 & S_{yz}^{(2,G)} \\ 0 & S_{yz}^{(2,G)} & 0 \end{pmatrix}$$

$$S_{xy}^{(2,G)} = G_{xy}$$
$$S_{yz}^{(2,G)} = G_{yz}$$

$$S_{yz}^{(2,G)} = G_y$$

$$A^{(2,G)} = \begin{pmatrix} 0 & A_{xy}^{(2,G)} & 0\\ -A_{xy}^{(2,G)} & 0 & A_{yz}^{(2,G)}\\ 0 & -A_{yz}^{(2,G)} & 0 \end{pmatrix}$$

$$A_{xy}^{(2,G)} = Q_x$$

$$A_{xy}^{(2,G)} = Q_z$$
$$A_{yz}^{(2,G)} = Q_x$$

$$S^{(3,G)} = \begin{pmatrix} 0 & S_{1y}^{(3,G)} & 0 \\ 0 & S_{2y}^{(3,G)} & 0 \\ 0 & S_{3y}^{(3,G)} & 0 \\ S_{4x}^{(3,G)} & 0 & S_{4z}^{(3,G)} \\ 0 & S_{5y}^{(3,G)} & 0 \\ S_{6x}^{(3,G)} & 0 & S_{6z}^{(3,G)} \end{pmatrix}$$

$$\begin{split} S_{1y}^{(3,G)} &= G_y[2] - G_y^\alpha - G_y^\beta + 2Q_{zx}[1] \\ S_{2y}^{(3,G)} &= 2G_y[1] + G_y[2] + 2G_y^\alpha \\ S_{3y}^{(3,G)} &= G_y[2] - G_y^\alpha + G_y^\beta - 2Q_{zx}[1] \\ S_{4x}^{(3,G)} &= G_{xyz} - 3Q_u[1] - Q_v[1] \\ S_{4z}^{(3,G)} &= G_y[1] - G_y^\alpha + G_y^\beta + Q_{zx}[1] \\ S_{5y}^{(3,G)} &= G_{xyz} + 3Q_u[1] - Q_v[1] \\ S_{6x}^{(3,G)} &= G_y[1] - G_y^\alpha - G_y^\beta - Q_{zx}[1] \end{split}$$

$$A^{(3,G)} = \begin{pmatrix} A_{4x}^{(3,G)} & 0 & A_{4z}^{(3,G)} \\ 0 & A_{5y}^{(3,G)} & 0 \\ A_{6x}^{(3,G)} & 0 & A_{6z}^{(3,G)} \end{pmatrix}$$

 $S_{6z}^{(3,G)} = G_{xyz} + 2Q_v[1]$

$$A_{4x}^{(3,G)} = Q_0 - Q_u[2] + Q_v[2]$$

$$A_{4z}^{(3,G)} = -G_y[3] + Q_{zx}[2]$$

$$A_{5y}^{(3,G)} = Q_0 - Q_u[2] - Q_v[2]$$

$$A_{6x}^{(3,G)} = G_y[3] + Q_{zx}[2]$$

$$A_{6z}^{(3,G)} = Q_0 + 2Q_u[2]$$

$$S^{(4,G)} = \begin{pmatrix} 0 & 0 & 0 & S_{14}^{(4,G)} & 0 & S_{16}^{(4,G)} \\ 0 & 0 & 0 & S_{24}^{(4,G)} & 0 & S_{26}^{(4,G)} \\ 0 & 0 & 0 & S_{34}^{(4,G)} & 0 & S_{36}^{(4,G)} \\ S_{14}^{(4,G)} & S_{24}^{(4,G)} & S_{34}^{(4,G)} & 0 & S_{45}^{(4,G)} & 0 \\ 0 & 0 & 0 & S_{45}^{(4,G)} & 0 & S_{56}^{(4,G)} \\ S_{16}^{(4,G)} & S_{26}^{(4,G)} & S_{36}^{(4,G)} & 0 & S_{56}^{(4,G)} & 0 \end{pmatrix}$$

$$\begin{split} S_{14}^{(4,G)} &= 2G_{4x}^{\beta} + G_{yz}[1] \\ S_{16}^{(4,G)} &= G_{4z}^{\alpha} - G_{4z}^{\beta} + G_{xy}[1] + 2G_{xy}[2] \\ S_{24}^{(4,G)} &= G_{4x}^{\alpha} - G_{4x}^{\beta} + G_{yz}[1] + 2G_{yz}[2] \\ S_{26}^{(4,G)} &= -G_{4z}^{\alpha} - G_{4z}^{\beta} + G_{xy}[1] + 2G_{xy}[2] \\ S_{34}^{(4,G)} &= -G_{4x}^{\alpha} - G_{4x}^{\beta} + G_{yz}[1] + 2G_{yz}[2] \\ S_{36}^{(4,G)} &= 2G_{4z}^{\beta} + G_{xy}[1] \\ S_{45}^{(4,G)} &= 2G_{4z}^{\beta} + G_{xy}[2] \\ S_{56}^{(4,G)} &= 2G_{4x}^{\beta} + G_{yz}[2] \end{split}$$

$$\bar{S}^{(4,G)} = \begin{pmatrix} 0 & 0 & 0 & \bar{S}^{(4,G)}_{14} & 0 & \bar{S}^{(4,G)}_{16} \\ 0 & 0 & 0 & \bar{S}^{(4,G)}_{24} & 0 & \bar{S}^{(4,G)}_{26} \\ 0 & 0 & 0 & \bar{S}^{(4,G)}_{24} & 0 & \bar{S}^{(4,G)}_{36} \\ -\bar{S}^{(4,G)}_{14} & -\bar{S}^{(4,G)}_{24} & -\bar{S}^{(4,G)}_{34} & 0 & \bar{S}^{(4,G)}_{45} & 0 \\ 0 & 0 & 0 & -\bar{S}^{(4,G)}_{45} & 0 & \bar{S}^{(4,G)}_{56} \\ -\bar{S}^{(4,G)}_{16} & -\bar{S}^{(4,G)}_{26} & -\bar{S}^{(4,G)}_{36} & 0 & -\bar{S}^{(4,G)}_{56} & 0 \end{pmatrix}$$

$$\begin{split} \bar{S}_{14}^{(4,G)} &= G_{yz}[3] - 4Q_x^{\beta}[1] \\ \bar{S}_{16}^{(4,G)} &= G_{xy}[3] + 2Q_z[1] - 2Q_z^{\alpha}[1] + 2Q_z^{\beta}[1] \\ \bar{S}_{24}^{(4,G)} &= G_{yz}[3] + 2Q_x[1] - 2Q_x^{\alpha}[1] + 2Q_x^{\beta}[1] \\ \bar{S}_{26}^{(4,G)} &= G_{xy}[3] - 2Q_z[1] + 2Q_z^{\alpha}[1] + 2Q_z^{\beta}[1] \\ \bar{S}_{34}^{(4,G)} &= G_{yz}[3] - 2Q_x[1] + 2Q_x^{\alpha}[1] + 2Q_x^{\beta}[1] \\ \bar{S}_{36}^{(4,G)} &= G_{xy}[3] - 4Q_z^{\beta}[1] \\ \bar{S}_{45}^{(4,G)} &= -Q_z[1] - 4Q_z^{\alpha}[1] \\ \bar{S}_{56}^{(4,G)} &= -Q_x[1] - 4Q_x^{\alpha}[1] \end{split}$$

$$A^{(4,G)} = \begin{pmatrix} 0 & A_{xy}^{(4,G)} & 0 \\ A_{xy}^{(4,G)} & 0 & A_{yz}^{(4,G)} \\ 0 & A_{yz}^{(4,G)} & 0 \end{pmatrix}$$

$$A_{xy}^{(4,G)} = 2G_{xy}[6]$$
$$A_{yz}^{(4,G)} = 2G_{yz}[6]$$

$$\bar{A}^{(4,G)} = \begin{pmatrix} 0 & \bar{A}_{xy}^{(4,G)} & 0 \\ -\bar{A}_{xy}^{(4,G)} & 0 & \bar{A}_{yz}^{(4,G)} \\ 0 & -\bar{A}_{yz}^{(4,G)} & 0 \end{pmatrix}$$

$$\bar{A}_{xy}^{(4,G)} = Q_z[6]$$

 $\bar{A}_{yz}^{(4,G)} = Q_x[6]$

$$M^{(4,G)} = \begin{pmatrix} M_{1x}^{(4,G)} & 0 & M_{1z}^{(4,G)} \\ M_{2x}^{(4,G)} & 0 & M_{2z}^{(4,G)} \\ M_{3x}^{(4,G)} & 0 & M_{3z}^{(4,G)} \\ 0 & M_{4y}^{(4,G)} & 0 \\ M_{5x}^{(4,G)} & 0 & M_{5z}^{(4,G)} \\ 0 & M_{6y}^{(4,G)} & 0 \end{pmatrix}$$

$$\begin{split} M_{1x}^{(4,G)} &= 2Q_x[2] + Q_x[3] + 2Q_x^{\alpha}[2] \\ M_{1z}^{(4,G)} &= -2G_{xy}[4] + Q_z[3] - Q_z^{\alpha}[2] + Q_z^{\beta}[2] \\ M_{2x}^{(4,G)} &= -2G_{yz}[4] + Q_x[3] - Q_x^{\alpha}[2] + Q_x^{\beta}[2] \\ M_{2z}^{(4,G)} &= 2G_{xy}[4] + Q_z[3] - Q_z^{\alpha}[2] - Q_z^{\beta}[2] \\ M_{3x}^{(4,G)} &= 2G_{yz}[4] + Q_x[3] - Q_x^{\alpha}[2] - Q_x^{\beta}[2] \\ M_{3z}^{(4,G)} &= 2Q_z[2] + Q_z[3] + 2Q_z^{\alpha}[2] \\ M_{4y}^{(4,G)} &= -G_{xy}[4] + Q_z[2] - Q_z^{\alpha}[2] - Q_z^{\beta}[2] \\ M_{5x}^{(4,G)} &= G_{xy}[4] + Q_z[2] - Q_x^{\alpha}[2] + Q_z^{\beta}[2] \\ M_{5z}^{(4,G)} &= -G_{yz}[4] + Q_x[2] - Q_x^{\alpha}[2] - Q_x^{\beta}[2] \\ M_{6y}^{(4,G)} &= G_{yz}[4] + Q_x[2] - Q_x^{\alpha}[2] + Q_x^{\beta}[2] \\ M_{6y}^{(4,G)} &= G_{yz}[4] + Q_x[2] - Q_x^{\alpha}[2] + Q_x^{\beta}[2] \end{split}$$

$$\bar{M}^{(4,G)} = \begin{pmatrix} \bar{M}_{x1}^{(4,G)} & \bar{M}_{x2}^{(4,G)} & \bar{M}_{x3}^{(4,G)} & 0 & \bar{M}_{x5}^{(4,G)} & 0\\ 0 & 0 & 0 & \bar{M}_{y4}^{(4,G)} & 0 & \bar{M}_{y6}^{(4,G)}\\ \bar{M}_{z1}^{(4,G)} & \bar{M}_{z2}^{(4,G)} & \bar{M}_{z3}^{(4,G)} & 0 & \bar{M}_{z5}^{(4,G)} & 0 \end{pmatrix}$$

$$\begin{split} \bar{M}_{x1}^{(4,G)} &= 2Q_x[4] + Q_x[5] + 2Q_x^{\alpha}[3] \\ \bar{M}_{x2}^{(4,G)} &= -2G_{yz}[5] + Q_x[5] - Q_x^{\alpha}[3] + Q_x^{\beta}[3] \\ \bar{M}_{x3}^{(4,G)} &= 2G_{yz}[5] + Q_x[5] - Q_x^{\alpha}[3] - Q_x^{\beta}[3] \\ \bar{M}_{x5}^{(4,G)} &= G_{xy}[5] + Q_z[4] - Q_z^{\alpha}[3] + Q_z^{\beta}[3] \\ \bar{M}_{y4}^{(4,G)} &= -G_{xy}[5] + Q_z[4] - Q_z^{\alpha}[3] - Q_z^{\beta}[3] \\ \bar{M}_{y4}^{(4,G)} &= G_{yz}[5] + Q_x[4] - Q_x^{\alpha}[3] + Q_x^{\beta}[3] \\ \bar{M}_{y6}^{(4,G)} &= G_{yz}[5] + Q_x[4] - Q_x^{\alpha}[3] + Q_x^{\beta}[3] \\ \bar{M}_{z1}^{(4,G)} &= -2G_{xy}[5] + Q_z[5] - Q_z^{\alpha}[3] - Q_z^{\beta}[3] \\ \bar{M}_{z2}^{(4,G)} &= 2G_{xy}[5] + Q_z[5] + 2Q_z^{\alpha}[3] \\ \bar{M}_{z3}^{(4,G)} &= 2Q_z[4] + Q_z[5] + 2Q_z^{\alpha}[3] \\ \bar{M}_{z5}^{(4,G)} &= -G_{yz}[5] + Q_x[4] - Q_x^{\alpha}[3] - Q_x^{\beta}[3] \end{split}$$