

MPG No. 37.5.155 $\bar{6}m'2'$ (-62'm' setting) [Type III, hexagonal] [M tensor]

* Rank 0 tensor. * Rank 1 tensor.

$$\begin{bmatrix} 0 & 0 & M_z \end{bmatrix}$$

$$M_z = M_{pz}^{(1)}$$

* Rank 2 tensor (s). * Rank 2 tensor (a). * Rank 3 tensor (s).

$$\begin{bmatrix} 0 & 0 & M_{xxz} \\ 0 & 0 & M_{xxz} \\ 0 & 0 & M_{zzz} \\ 0 & M_{yzy} & 0 \\ M_{yzy} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$M_{xxz} = -M_{faz}^{(1)} + M_{pz}^{(1)}$$

$$M_{zzz} = 2M_{faz}^{(1)} + M_{pz}^{(1)} + 2M_{pz}^{(2)}$$

$$M_{yzy} = -M_{faz}^{(1)} + M_{pz}^{(2)}$$

* Rank 3 tensor (a).

$$\begin{bmatrix} 0 & M_{yzy} & 0 \\ -M_{yzy} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$M_{yzy} = M_{pz}^{(3)}$$

* Rank 4 tensor (sss).

$$\begin{bmatrix} 0 & 0 & 0 & 0 & M_{xxxx} & 0 \\ 0 & 0 & 0 & 0 & -M_{xxxx} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -M_{xxxx} \\ M_{xxxx} & -M_{xxxx} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -M_{xxxx} & 0 & 0 \end{bmatrix}$$

$$M_{xxxx} = M_{ga}^{(1)}$$

* Rank 4 tensor (ssa).

$$\begin{bmatrix} 0 & 0 & 0 & 0 & M_{xxxx} & 0 \\ 0 & 0 & 0 & 0 & -M_{xxxx} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & M_{xxxx} \\ -M_{xxxx} & M_{xxxx} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -M_{xxxx} & 0 & 0 \end{bmatrix}$$

$$M_{xxxx} = -2T_{f1}^{(1)}$$

* Rank 4 tensor (aas). * Rank 4 tensor (aaa). * Rank 4 tensor (sa).

$$\begin{bmatrix} 0 & M_{xxxx} & 0 \\ 0 & -M_{xxxx} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ M_{xxxx} & 0 & 0 \end{bmatrix}$$

$$M_{xxxx} = T_{f1}^{(2)}$$

* Rank 4 tensor (as).

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & M_{yzxy} \\ M_{yzxy} & -M_{yzxy} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$M_{yzxy} = T_{f1}^{(3)}$$

* Rank 4 tensor (s).

$$\begin{bmatrix} 0 & 0 & 0 & 0 & M_{xxxx} & 0 & 0 & M_{xxxz} & 0 \\ 0 & 0 & 0 & 0 & -M_{xxxx} & 0 & 0 & -M_{xxxz} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & M_{yzxy} & 0 & 0 & M_{yzxy} \\ -M_{yzxy} & M_{yzxy} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -M_{xxxz} & 0 & 0 & -M_{xxxz} & 0 & 0 \end{bmatrix}$$

$$M_{xxxx} = M_{ga}^{(1)} - 2T_{f1}^{(1)} + T_{f1}^{(2)}$$

$$M_{xxxz} = M_{ga}^{(1)} - 2T_{f1}^{(1)} - T_{f1}^{(2)}$$

$$M_{yzxy} = -M_{ga}^{(1)} - 2T_{f1}^{(1)}$$

* Rank 4 tensor (a).

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & M_{yzxy} & 0 & 0 & M_{yzxy} \\ M_{yzxy} & -M_{yzxy} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$M_{yzxy} = T_{f1}^{(3)}$$

* Rank 4 tensor (t).

$$\begin{bmatrix} 0 & 0 & M_{xxxz} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ -M_{xxxz} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ M_{xxxz} & 0 & 0 \\ 0 & 0 & -M_{xxxz} \\ 0 & -M_{xxxz} & 0 \end{bmatrix}$$

$$M_{xxxz} = M_{ga}^{(1)}$$