

SG No. 11 C_{2h}^2 $P2_1/m$ (b-axis setting) [monoclinic]

* plus set: $+ [0, 0, 0]$

Table 1: Wyckoff site: **2a**, site symmetry: -1

No.	position	mapping
1	$[0, 0, 0]$	$[1, 3]$
2	$[0, \frac{1}{2}, 0]$	$[2, 4]$

Table 2: Wyckoff site: **2b**, site symmetry: -1

No.	position	mapping
1	$[\frac{1}{2}, 0, 0]$	$[1, 3]$
2	$[\frac{1}{2}, \frac{1}{2}, 0]$	$[2, 4]$

Table 3: Wyckoff site: **2c**, site symmetry: -1

No.	position	mapping
1	$[0, 0, \frac{1}{2}]$	$[1, 3]$
2	$[0, \frac{1}{2}, \frac{1}{2}]$	$[2, 4]$

Table 4: Wyckoff site: **2d**, site symmetry: -1

No.	position	mapping
1	$[\frac{1}{2}, 0, \frac{1}{2}]$	$[1, 3]$
2	$[\frac{1}{2}, \frac{1}{2}, \frac{1}{2}]$	$[2, 4]$

Table 5: Wyckoff site: **2e**, site symmetry: m

No.	position	mapping
1	$[x, \frac{1}{4}, z]$	$[1, 4]$
2	$[-x, \frac{3}{4}, -z]$	$[2, 3]$

Table 6: Wyckoff site: 4f, site symmetry: 1

No.	position	mapping
1	$[x, y, z]$	[1]
2	$[-x, y + \frac{1}{2}, -z]$	[2]
3	$[-x, -y, -z]$	[3]
4	$[x, \frac{1}{2} - y, z]$	[4]