

Model for “graphene”

Generated on 2026-07-19 19:34:38 by MultiPie 2.2.4

General Condition

- Basis type: 1g
- SAMB selection:
 - Type: [Q, G, M, T]
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_{1g} , A_{1u} , A_{2g} , A_{2u} , B_{1g} , B_{1u} , B_{2g} , B_{2u} , E_{1g} , E_{1u} , E_{2g} , E_{2u}]
 - Spin (s): [0, 1]
- Atomic selection:
 - Type: [Q, G, M, T]
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_{1g} , A_{1u} , A_{2g} , A_{2u} , B_{1g} , B_{1u} , B_{2g} , B_{2u} , E_{1g} , E_{1u} , E_{2g} , E_{2u}]
 - Spin (s): [0, 1]
- Site-cluster selection:
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_{1g} , A_{1u} , A_{2g} , A_{2u} , B_{1g} , B_{1u} , B_{2g} , B_{2u} , E_{1g} , E_{1u} , E_{2g} , E_{2u}]
- Bond-cluster selection:
 - Type: [Q, G, M, T]
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_{1g} , A_{1u} , A_{2g} , A_{2u} , B_{1g} , B_{1u} , B_{2g} , B_{2u} , E_{1g} , E_{1u} , E_{2g} , E_{2u}]
- Max. neighbor: 10
- Search cell range: (-2, 3), (-2, 3), (-2, 3)
- Toroidal priority: **false**

Group and Unit Cell

- Group: SG No. 191 D_{6h}^1 $P6/mmm$ [hexagonal]
- Associated point group: PG No. 191 D_{6h} $6/mmm$ [hexagonal]
- Unit cell:
 - $a = 1.00000$, $b = 1.00000$, $c = 4.00000$, $\alpha = 90.0$, $\beta = 90.0$, $\gamma = 120.0$
- Lattice vectors (conventional cell):
 - $\mathbf{a}_1 = [1.00000, 0.00000, 0.00000]$
 - $\mathbf{a}_2 = [-0.50000, 0.86603, 0.00000]$
 - $\mathbf{a}_3 = [0.00000, 0.00000, 4.00000]$

Symmetry Operation

Table 1: Symmetry operation

#	SO	#	SO	#	SO	#	SO	#	SO
1	$\{1 0\}$	2	$\{3_{001}^+ 0\}$	3	$\{3_{001}^- 0\}$	4	$\{2_{001} 0\}$	5	$\{6_{001}^- 0\}$
6	$\{6_{001}^+ 0\}$	7	$\{2_{110} 0\}$	8	$\{2_{100} 0\}$	9	$\{2_{010} 0\}$	10	$\{2_{1-10} 0\}$
11	$\{2_{120} 0\}$	12	$\{2_{210} 0\}$	13	$\{-1 0\}$	14	$\{-3_{001}^+ 0\}$	15	$\{-3_{001}^- 0\}$
16	$\{m_{001} 0\}$	17	$\{-6_{001}^- 0\}$	18	$\{-6_{001}^+ 0\}$	19	$\{m_{110} 0\}$	20	$\{m_{100} 0\}$
21	$\{m_{010} 0\}$	22	$\{m_{1-10} 0\}$	23	$\{m_{120} 0\}$	24	$\{m_{210} 0\}$		

Harmonics

Table 2: Harmonics

#	symbol	irrep.	rank	X	multiplicity	component	symmetry
1	$\mathbb{Q}_0(A_{1g})$	A_{1g}	0	Q, T	-	-	1
2	$\mathbb{G}_1(A_{2g})$	A_{2g}	1	G, M	-	-	z
3	$\mathbb{Q}_3(B_{1u})$	B_{1u}	3	Q, T	-	-	$\frac{\sqrt{10}y(3x^2-y^2)}{4}$
4	$\mathbb{Q}_3(B_{2u})$	B_{2u}	3	Q, T	-	-	$\frac{\sqrt{10}x(x^2-3y^2)}{4}$
5	$\mathbb{Q}_{1,1}(E_{1u})$	E_{1u}	1	Q, T	-	1	x
6	$\mathbb{Q}_{1,2}(E_{1u})$					2	y

continued ...

Table 2

#	symbol	irrep.	rank	X	multiplicity	component	symmetry
7	$\mathbb{Q}_{2,1}(E_{2g})$	E_{2g}	2	Q, T	-	1	$\frac{\sqrt{3}(x-y)(x+y)}{2}$
8	$\mathbb{Q}_{2,2}(E_{2g})$					2	$-\sqrt{3}xy$

Basis in full matrix

Table 3: dimension = 2

#	orbital@atom(SL)	#	orbital@atom(SL)
0	$ p_z\rangle @C(1)$	1	$ p_z\rangle @C(2)$

Table 4: Atomic basis (orbital part only)

orbital	definition
$ p_x\rangle$	x
$ p_y\rangle$	y
$ p_z\rangle$	z

SAMB: 20 (all 20)

- **C : 'C' site-cluster**

- * bra: $\langle p_z |$

- * ket: $|p_z\rangle$

- * wyckoff: **2c**

$$\boxed{\text{z1}} \quad \mathbb{Q}_0^{(c)}(A_{1g}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_0^{(s)}(A_{1g})$$

$$\boxed{\text{z5}} \quad \mathbb{Q}_3^{(c)}(B_{1u}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_3^{(s)}(B_{1u})$$

- **C;C_001_1 : 'C'-'C' bond-cluster**

- * bra: $\langle p_z |$

- * ket: $|p_z\rangle$

- * wyckoff: **3a03f**

$$\boxed{\text{z2}} \quad \mathbb{Q}_0^{(c)}(A_{1g}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_0^{(b)}(A_{1g})$$

$$\boxed{\text{z6}} \quad \mathbb{T}_3^{(c)}(B_{1u}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_3^{(b)}(B_{1u})$$

$$\boxed{\text{z9}} \quad \mathbb{T}_{1,1}^{(c)}(E_{1u}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_{1,1}^{(b)}(E_{1u})}{2}$$

$$\boxed{\text{z10}} \quad \mathbb{T}_{1,2}^{(c)}(E_{1u}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_{1,2}^{(b)}(E_{1u})}{2}$$

$$\boxed{\text{z15}} \quad \mathbb{Q}_{2,1}^{(c)}(E_{2g}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_{2,1}^{(b)}(E_{2g})}{2}$$

$$\boxed{\text{z16}} \quad \mathbb{Q}_{2,2}^{(c)}(E_{2g}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_{2,2}^{(b)}(E_{2g})}{2}$$

- **C;C_002_1 : 'C'-'C' bond-cluster**

- * bra: $\langle p_z |$

- * ket: $|p_z\rangle$

- * wyckoff: **6b061**

$$\boxed{\text{z3}} \quad \mathbb{Q}_0^{(c)}(A_{1g}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_0^{(b)}(A_{1g})$$

$$\boxed{\text{z4}} \quad \mathbb{M}_1^{(c)}(A_{2g}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{M}_1^{(b)}(A_{2g})$$

$$\boxed{z7} \quad \mathbb{Q}_3^{(c)}(B_{1u}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_3^{(b)}(B_{1u})$$

$$\boxed{z8} \quad \mathbb{T}_3^{(c)}(B_{2u}) = \mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_3^{(b)}(B_{2u})$$

$$\boxed{z11} \quad \mathbb{Q}_{1,1}^{(c)}(E_{1u}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_{1,1}^{(b)}(E_{1u})}{2}$$

$$\boxed{z12} \quad \mathbb{Q}_{1,2}^{(c)}(E_{1u}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_{1,2}^{(b)}(E_{1u})}{2}$$

$$\boxed{z13} \quad \mathbb{T}_{1,1}^{(c)}(E_{1u}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_{1,1}^{(b)}(E_{1u})}{2}$$

$$\boxed{z14} \quad \mathbb{T}_{1,2}^{(c)}(E_{1u}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_{1,2}^{(b)}(E_{1u})}{2}$$

$$\boxed{z17} \quad \mathbb{Q}_{2,1}^{(c)}(E_{2g}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_{2,1}^{(b)}(E_{2g})}{2}$$

$$\boxed{z18} \quad \mathbb{Q}_{2,2}^{(c)}(E_{2g}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{Q}_{2,2}^{(b)}(E_{2g})}{2}$$

$$\boxed{z19} \quad \mathbb{T}_{2,1}^{(c)}(E_{2g}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_{2,1}^{(b)}(E_{2g})}{2}$$

$$\boxed{z20} \quad \mathbb{T}_{2,2}^{(c)}(E_{2g}) = \frac{\sqrt{2}\mathbb{Q}_0^{(a)}(A_{1g})\mathbb{T}_{2,2}^{(b)}(E_{2g})}{2}$$

Atomic SAMB

- bra: $\langle p_z |$
- ket: $|p_z\rangle$

$$\boxed{x1} \quad \mathbb{Q}_0^{(a)}(A_{1g}) = [1]$$

Cluster SAMB

- Site cluster

** Wyckoff: 2c

$$\boxed{\text{y1}} \quad \mathbb{Q}_0^{(s)}(A_{1g}) = \left[\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right]$$

$$\boxed{\text{y2}} \quad \mathbb{Q}_3^{(s)}(B_{1u}) = \left[\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right]$$

- Bond cluster

** Wyckoff: 3a@3f

$$\boxed{\text{y3}} \quad \mathbb{Q}_0^{(b)}(A_{1g}) = \left[\frac{\sqrt{3}}{3}, \frac{\sqrt{3}}{3}, \frac{\sqrt{3}}{3} \right]$$

$$\boxed{\text{y4}} \quad \mathbb{T}_3^{(b)}(B_{1u}) = \left[\frac{\sqrt{3}i}{3}, \frac{\sqrt{3}i}{3}, \frac{\sqrt{3}i}{3} \right]$$

$$\boxed{\text{y5}} \quad \mathbb{T}_{1,1}^{(b)}(E_{1u}) = \left[0, -\frac{\sqrt{2}i}{2}, \frac{\sqrt{2}i}{2} \right]$$

$$\boxed{\text{y6}} \quad \mathbb{T}_{1,2}^{(b)}(E_{1u}) = \left[\frac{\sqrt{6}i}{3}, -\frac{\sqrt{6}i}{6}, -\frac{\sqrt{6}i}{6} \right]$$

$$\boxed{\text{y7}} \quad \mathbb{Q}_{2,1}^{(b)}(E_{2g}) = \left[\frac{\sqrt{6}}{3}, -\frac{\sqrt{6}}{6}, -\frac{\sqrt{6}}{6} \right]$$

$$\boxed{\text{y8}} \quad \mathbb{Q}_{2,2}^{(b)}(E_{2g}) = \left[0, \frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right]$$

** Wyckoff: 6b@6l

$$\boxed{\text{y9}} \quad \mathbb{Q}_0^{(b)}(A_{1g}) = \left[\frac{\sqrt{6}}{6}, \frac{\sqrt{6}}{6}, \frac{\sqrt{6}}{6}, \frac{\sqrt{6}}{6}, \frac{\sqrt{6}}{6}, \frac{\sqrt{6}}{6} \right]$$

$$\boxed{\text{y10}} \quad \mathbb{M}_1^{(b)}(A_{2g}) = \left[\frac{\sqrt{6}i}{6}, \frac{\sqrt{6}i}{6}, \frac{\sqrt{6}i}{6}, \frac{\sqrt{6}i}{6}, \frac{\sqrt{6}i}{6}, \frac{\sqrt{6}i}{6} \right]$$

$$\boxed{\text{y11}} \quad \mathbb{Q}_3^{(b)}(B_{1u}) = \left[\frac{\sqrt{6}}{6}, \frac{\sqrt{6}}{6}, \frac{\sqrt{6}}{6}, -\frac{\sqrt{6}}{6}, -\frac{\sqrt{6}}{6}, -\frac{\sqrt{6}}{6} \right]$$

$$\boxed{\text{y12}} \quad \mathbb{T}_3^{(b)}(B_{2u}) = \left[\frac{\sqrt{6}i}{6}, \frac{\sqrt{6}i}{6}, \frac{\sqrt{6}i}{6}, -\frac{\sqrt{6}i}{6}, -\frac{\sqrt{6}i}{6}, -\frac{\sqrt{6}i}{6} \right]$$

$$\boxed{\text{y13}} \quad \mathbb{Q}_{1,1}^{(b)}(E_{1u}) = \left[0, -\frac{1}{2}, \frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right]$$

$$\boxed{\text{y14}} \quad \mathbb{Q}_{1,2}^{(b)}(E_{1u}) = \left[\frac{\sqrt{3}}{3}, -\frac{\sqrt{3}}{6}, -\frac{\sqrt{3}}{6}, -\frac{\sqrt{3}}{3}, \frac{\sqrt{3}}{6}, \frac{\sqrt{3}}{6} \right]$$

$$\boxed{\text{y15}} \quad \mathbb{T}_{1,1}^{(b)}(E_{1u}) = \left[\frac{\sqrt{3}i}{3}, -\frac{\sqrt{3}i}{6}, -\frac{\sqrt{3}i}{6}, -\frac{\sqrt{3}i}{3}, \frac{\sqrt{3}i}{6}, \frac{\sqrt{3}i}{6} \right]$$

$$\boxed{\text{y16}} \quad \mathbb{T}_{1,2}^{(b)}(E_{1u}) = \left[0, \frac{i}{2}, -\frac{i}{2}, 0, -\frac{i}{2}, \frac{i}{2} \right]$$

$$\boxed{\text{y17}} \quad \mathbb{Q}_{2,1}^{(b)}(E_{2g}) = \left[\frac{\sqrt{3}}{3}, -\frac{\sqrt{3}}{6}, -\frac{\sqrt{3}}{6}, \frac{\sqrt{3}}{3}, -\frac{\sqrt{3}}{6}, -\frac{\sqrt{3}}{6} \right]$$

$$\boxed{\text{y18}} \quad \mathbb{Q}_{2,2}^{(b)}(E_{2g}) = \left[0, \frac{1}{2}, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{1}{2} \right]$$

$$\boxed{\text{y19}} \quad \mathbb{T}_{2,1}^{(b)}(E_{2g}) = \left[0, \frac{i}{2}, -\frac{i}{2}, 0, \frac{i}{2}, -\frac{i}{2} \right]$$

$$\boxed{\text{y20}} \quad \mathbb{T}_{2,2}^{(b)}(E_{2g}) = \left[-\frac{\sqrt{3}i}{3}, \frac{\sqrt{3}i}{6}, \frac{\sqrt{3}i}{6}, -\frac{\sqrt{3}i}{3}, \frac{\sqrt{3}i}{6}, \frac{\sqrt{3}i}{6} \right]$$

— Site and Bond

Table 5: Orbital of each site

#	site	orbital
1	c	$ p_z\rangle$

Table 6: Neighbor and bra-ket of each bond

#	head	tail	neighbor	head (bra)	tail (ket)
1	c	c	[1,2]	[p]	[p]

— Site in Unit Cell —

Sites in (conventional) cell (no plus set), SL = sublattice

Table 7: 'C' (#1) site cluster (2c), -6m2

SL	position (s)	mapping
1	[0.33333, 0.66667, 0.00000]	[1,2,3,10,11,12,16,17,18,19,20,21]
2	[0.66667, 0.33333, 0.00000]	[4,5,6,7,8,9,13,14,15,22,23,24]

— Bond in Unit Cell —

Bonds in (conventional) cell (no plus set): tail, head = (SL, plus set), (N)D = (non)directional (listed up to 5th neighbor at most)

Table 8: 1-th 'C'-'C' [1] (#1) bond cluster (3a@3f), ND, $|v|=0.57735$ (cartesian)

SL	vector (v)	center (c)	mapping	head	tail	R (primitive)
1	[0.33333, 0.66667, 0.00000]	[0.50000, 0.00000, 0.00000]	[1,-4,-8,11,-13,16,20,-23]	(2,1)	(1,1)	[0,-1,0]
2	[-0.66667,-0.33333, 0.00000]	[0.00000, 0.50000, 0.00000]	[2,-5,-7,10,-14,17,19,-22]	(2,1)	(1,1)	[1,0,0]
3	[0.33333,-0.33333, 0.00000]	[0.50000, 0.50000, 0.00000]	[3,-6,-9,12,-15,18,21,-24]	(2,1)	(1,1)	[0,0,0]

Table 9: 2-th 'C'-'C' [1] (#2) bond cluster (6b@6l), ND, $|v|=1.0$ (cartesian)

SL	vector (v)	center (c)	mapping	head	tail	R (primitive)
1	[1.00000, 0.00000, 0.00000]	[0.83333, 0.66667, 0.00000]	[1,-11,16,-20]	(1,1)	(1,1)	[-1,0,0]
2	[0.00000, 1.00000, 0.00000]	[0.33333, 0.16667, 0.00000]	[2,-10,17,-19]	(1,1)	(1,1)	[0,-1,0]
3	[-1.00000,-1.00000, 0.00000]	[0.83333, 0.16667, 0.00000]	[3,-12,18,-21]	(1,1)	(1,1)	[1,1,0]
4	[-1.00000, 0.00000, 0.00000]	[0.16667, 0.33333, 0.00000]	[4,-8,13,-23]	(2,1)	(2,1)	[1,0,0]
5	[0.00000,-1.00000, 0.00000]	[0.66667, 0.83333, 0.00000]	[5,-7,14,-22]	(2,1)	(2,1)	[0,1,0]
6	[1.00000, 1.00000, 0.00000]	[0.16667, 0.83333, 0.00000]	[6,-9,15,-24]	(2,1)	(2,1)	[-1,-1,0]