

Model for “C2h1”

Generated on 2026-04-04 02:12:24 by MultiPie 2.1.3

General Condition

- Basis type: **lgs**
- SAMB selection:
 - Type: [Q, G]
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_g , A_u , B_g , B_u]
 - Spin (s): [0, 1]
- Atomic selection:
 - Type: [Q, G, M, T]
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_g , A_u , B_g , B_u]
 - Spin (s): [0, 1]
- Site-cluster selection:
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_g , A_u , B_g , B_u]
- Bond-cluster selection:
 - Type: [Q, G, M, T]
 - Rank: [0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
 - Irrep.: [A_g , A_u , B_g , B_u]
- Max. neighbor: 10
- Search cell range: (-2, 3), (-2, 3), (-2, 3)
- Toroidal priority: **false**

Group and Unit Cell

- Group: SG No. 10 C_{2h}^1 $P2/m$ (b-axis setting) [monoclinic]
- Associated point group: PG No. 10 C_{2h} $2/m$ (b-axis setting) [monoclinic]
- Unit cell:
 - $a = 1.00000$, $b = 1.20000$, $c = 1.00000$, $\alpha = 90.0$, $\beta = 90.0$, $\gamma = 90.0$
- Lattice vectors (conventional cell):
 - $\mathbf{a}_1 = [1.00000, 0.00000, 0.00000]$
 - $\mathbf{a}_2 = [0.00000, 1.20000, 0.00000]$
 - $\mathbf{a}_3 = [0.00000, 0.00000, 1.00000]$

Symmetry Operation

Table 1: Symmetry operation

#	SO	#	SO	#	SO	#	SO	#	SO
1	$\{1 0\}$	2	$\{2_{010} 0\}$	3	$\{-1 0\}$	4	$\{m_{010} 0\}$		

Harmonics

Table 2: Harmonics

#	symbol	irrep.	rank	X	multiplicity	component	symmetry
1	$\mathbb{Q}_0(A_g)$	A_g	0	Q, T	-	-	1
2	$\mathbb{G}_1(A_g)$	A_g	1	G, M	-	-	y
3	$\mathbb{Q}_2(A_g, 3)$	A_g	2	Q, T	3	-	$\sqrt{3}xz$
4	$\mathbb{G}_0(A_u)$	A_u	0	G, M	-	-	1
5	$\mathbb{Q}_1(A_u)$	A_u	1	Q, T	-	-	y
6	$\mathbb{G}_2(A_u, 2)$	A_u	2	G, M	2	-	$\frac{\sqrt{3}(x-y)(x+y)}{2}$
7	$\mathbb{G}_1(B_g, 1)$	B_g	1	G, M	1	-	x
8	$\mathbb{G}_1(B_g, 2)$	B_g	1	G, M	2	-	z
9	$\mathbb{Q}_2(B_g, 1)$	B_g	2	Q, T	1	-	$\sqrt{3}yz$
10	$\mathbb{Q}_2(B_g, 2)$	B_g	2	Q, T	2	-	$\sqrt{3}xy$

continued ...

Table 2

#	symbol	irrep.	rank	X	multiplicity	component	symmetry
11	$\mathbb{Q}_1(B_u, 1)$	B_u	1	Q, T	1	-	x
12	$\mathbb{Q}_1(B_u, 2)$	B_u	1	Q, T	2	-	z
13	$\mathbb{G}_2(B_u, 2)$	B_u	2	G, M	2	-	$\sqrt{3}xy$

Basis in full matrix

Table 3: dimension = 4

#	orbital@atom(SL)	#	orbital@atom(SL)	#	orbital@atom(SL)	#	orbital@atom(SL)
0	$ s, \uparrow\rangle @A(1)$	1	$ s, \downarrow\rangle @A(1)$	2	$ s, \uparrow\rangle @B(1)$	3	$ s, \downarrow\rangle @B(1)$

Table 4: Atomic basis (orbital part only)

orbital	definition
$ s\rangle$	1

SAMB: 34 (all 34)

- A : 'A' site-cluster

- * bra: $\langle s, \uparrow |, \langle s, \downarrow |$
- * ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$
- * wyckoff: **1a**

$$\boxed{\text{z1}} \quad \mathbb{Q}_0^{(c)}(A_g) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_0^{(s)}(A_g)$$

• **B : 'B' site-cluster**

- * bra: $\langle s, \uparrow |, \langle s, \downarrow |$
- * ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$
- * wyckoff: **1e**

$$\boxed{\text{z2}} \quad \mathbb{Q}_0^{(c)}(A_g) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_0^{(s)}(A_g)$$

• **A;A_001_1 : 'A'-'A' bond-cluster**

- * bra: $\langle s, \uparrow |, \langle s, \downarrow |$
- * ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$
- * wyckoff: **1a@1d**

$$\boxed{\text{z3}} \quad \mathbb{Q}_0^{(c)}(A_g) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_0^{(b)}(A_g)$$

$$\boxed{\text{z11}} \quad \mathbb{Q}_1^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z12}} \quad \mathbb{G}_0^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z27}} \quad \mathbb{Q}_1^{(1,-1;c)}(B_u, 2) = -\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(B_u, 1)$$

• **A;A_001_2 : 'A'-'A' bond-cluster**

- * bra: $\langle s, \uparrow |, \langle s, \downarrow |$
- * ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$
- * wyckoff: **1a@1c**

$$\boxed{\text{z4}} \quad \mathbb{Q}_0^{(c)}(A_g) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_0^{(b)}(A_g)$$

$$\boxed{\text{z13}} \quad \mathbb{Q}_1^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z14}} \quad \mathbb{G}_0^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z28}} \quad \mathbb{Q}_1^{(1,-1;c)}(B_u, 2) = -\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(B_u, 1)$$

• **B;A_001_1** : 'A'-'B' bond-cluster

* bra: $\langle s, \uparrow |, \langle s, \downarrow |$

* ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$

* wyckoff: **4a@4o**

$$\boxed{\text{z5}} \quad \mathbb{Q}_0^{(c)}(A_g) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_0^{(b)}(A_g)$$

$$\boxed{\text{z6}} \quad \mathbb{Q}_0^{(1,-1;c)}(A_g) = \mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{M}_1^{(b)}(B_g, 1)$$

$$\boxed{\text{z7}} \quad \mathbb{Q}_2^{(1,-1;c)}(A_g, 3) = \mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{M}_1^{(b)}(B_g, 1)$$

$$\boxed{\text{z8}} \quad \mathbb{G}_1^{(1,-1;c)}(A_g) = \mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_0^{(b)}(A_g)$$

$$\boxed{\text{z15}} \quad \mathbb{Q}_1^{(c)}(A_u) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_1^{(b)}(A_u)$$

$$\boxed{\text{z16}} \quad \mathbb{Q}_1^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z17}} \quad \mathbb{G}_0^{(1,-1;c)}(A_u) = \frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(A_u)}{2} + \frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(B_u, 1)}{2}$$

$$\boxed{\text{z18}} \quad \mathbb{G}_2^{(1,-1;c)}(A_u, 2) = -\frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(A_u)}{2} + \frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(B_u, 1)}{2}$$

$$\boxed{\text{z23}} \quad \mathbb{Q}_2^{(c)}(B_g, 1) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_2^{(b)}(B_g, 1)$$

$$\boxed{\text{z24}} \quad \mathbb{Q}_2^{(1,-1;c)}(B_g, 2) = \mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{M}_1^{(b)}(B_g, 1)$$

$$\boxed{\text{z25}} \quad \mathbb{G}_1^{(1,-1;c)}(B_g, 1) = \mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_0^{(b)}(A_g)$$

$$\boxed{\text{z26}} \quad \mathbb{G}_1^{(1,-1;c)}(B_g, 2) = \mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{T}_0^{(b)}(A_g)$$

$$\boxed{\text{z29}} \quad \mathbb{Q}_1^{(c)}(B_u, 1) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z30}} \quad \mathbb{Q}_1^{(1,-1;c)}(B_u, 1) = -\mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{T}_1^{(b)}(A_u)$$

$$\boxed{\text{z31}} \quad \mathbb{Q}_1^{(1,-1;c)}(B_u, 2) = -\frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(B_u, 1)}{2} + \frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(A_u)}{2}$$

$$\boxed{\text{z32}} \quad \mathbb{G}_2^{(1,-1;c)}(B_u, 2) = \frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(B_u, 1)}{2} + \frac{\sqrt{2}\mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(A_u)}{2}$$

- **B;B_001_1** : 'B'-'B' bond-cluster

* bra: $\langle s, \uparrow |, \langle s, \downarrow |$

* ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$

* wyckoff: **1a@1b**

$$\boxed{\text{z9}} \quad \mathbb{Q}_0^{(c)}(A_g) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_0^{(b)}(A_g)$$

$$\boxed{\text{z19}} \quad \mathbb{Q}_1^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z20}} \quad \mathbb{G}_0^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z33}} \quad \mathbb{Q}_1^{(1,-1;c)}(B_u, 2) = -\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(B_u, 1)$$

- **B;B_001_2** : 'B'-'B' bond-cluster

* bra: $\langle s, \uparrow |, \langle s, \downarrow |$

* ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$

* wyckoff: **1a@1h**

$$\boxed{\text{z10}} \quad \mathbb{Q}_0^{(c)}(A_g) = \mathbb{Q}_0^{(a)}(A_g)\mathbb{Q}_0^{(b)}(A_g)$$

$$\boxed{\text{z21}} \quad \mathbb{Q}_1^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 2)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z22}} \quad \mathbb{G}_0^{(1,-1;c)}(A_u) = \mathbb{M}_1^{(1,-1;a)}(B_g, 1)\mathbb{T}_1^{(b)}(B_u, 1)$$

$$\boxed{\text{z34}} \quad \mathbb{Q}_1^{(1,-1;c)}(B_u, 2) = -\mathbb{M}_1^{(1,-1;a)}(A_g)\mathbb{T}_1^{(b)}(B_u, 1)$$

Atomic SAMB

- bra: $\langle s, \uparrow |, \langle s, \downarrow |$

- ket: $|s, \uparrow\rangle, |s, \downarrow\rangle$

$$\boxed{\text{x1}} \quad \mathbb{Q}_0^{(a)}(A_g) = \begin{bmatrix} \frac{\sqrt{2}}{2} & 0 \\ 0 & \frac{\sqrt{2}}{2} \end{bmatrix}$$

$$\boxed{\text{x2}} \quad \mathbb{M}_1^{(1,-1;a)}(A_g) = \begin{bmatrix} 0 & -\frac{\sqrt{2}i}{2} \\ \frac{\sqrt{2}i}{2} & 0 \end{bmatrix}$$

$$\boxed{\text{x3}} \quad \mathbb{M}_1^{(1,-1;a)}(B_g, 1) = \begin{bmatrix} 0 & \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & 0 \end{bmatrix}$$

$$\boxed{\text{x4}} \quad \mathbb{M}_1^{(1,-1;a)}(B_g, 2) = \begin{bmatrix} \frac{\sqrt{2}}{2} & 0 \\ 0 & -\frac{\sqrt{2}}{2} \end{bmatrix}$$

Cluster SAMB

- Site cluster

** Wyckoff: **1e**

$$\boxed{\text{y1}} \quad \mathbb{Q}_0^{(s)}(A_g) = [1]$$

** Wyckoff: **1a**

$$\boxed{\text{y2}} \quad \mathbb{Q}_0^{(s)}(A_g) = [1]$$

- Bond cluster

** Wyckoff: **1a@1h**

$$\boxed{\text{y3}} \quad \mathbb{Q}_0^{(b)}(A_g) = [1]$$

$$\boxed{\text{y4}} \quad \mathbb{T}_1^{(b)}(B_u, 1) = [i]$$

** Wyckoff: **1a@1d**

$$\boxed{\text{y5}} \quad \mathbb{Q}_0^{(b)}(A_g) = [1]$$

$$\boxed{\text{y6}} \quad \mathbb{T}_1^{(b)}(B_u, 1) = [i]$$

** Wyckoff: **1a@1b**

$$\boxed{\text{y7}} \quad \mathbb{Q}_0^{(b)}(A_g) = [1]$$

$$\boxed{\text{y8}} \quad \mathbb{T}_1^{(b)}(B_u, 1) = [i]$$

** Wyckoff: **1a@1c**

$$\boxed{\text{y9}} \quad \mathbb{Q}_0^{(b)}(A_g) = [1]$$

$$\boxed{\text{y10}} \quad \mathbb{T}_1^{(b)}(B_u, 1) = [i]$$

** Wyckoff: **4a@4o**

$$\boxed{\text{y11}} \quad \mathbb{Q}_0^{(b)}(A_g) = \left[\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right]$$

$$\boxed{\text{y12}} \quad \mathbb{T}_0^{(b)}(A_g) = \left[\frac{i}{2}, \frac{i}{2}, \frac{i}{2}, \frac{i}{2} \right]$$

$$\boxed{\text{y13}} \quad \mathbb{Q}_1^{(b)}(A_u) = \left[\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2} \right]$$

$$\boxed{\text{y14}} \quad \mathbb{T}_1^{(b)}(A_u) = \left[\frac{i}{2}, \frac{i}{2}, -\frac{i}{2}, -\frac{i}{2} \right]$$

$$\boxed{\text{y15}} \quad \mathbb{M}_1^{(b)}(B_g, 1) = \left[\frac{i}{2}, -\frac{i}{2}, \frac{i}{2}, -\frac{i}{2} \right]$$

$$\boxed{\text{y16}} \quad \mathbb{Q}_2^{(b)}(B_g, 1) = \left[\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2} \right]$$

$$\boxed{\text{y17}} \quad \mathbb{Q}_1^{(b)}(B_u, 1) = \left[\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2} \right]$$

$$\boxed{\text{y18}} \quad \mathbb{T}_1^{(b)}(B_u, 1) = \left[\frac{i}{2}, -\frac{i}{2}, -\frac{i}{2}, \frac{i}{2} \right]$$

— Site and Bond —

Table 5: Orbital of each site

#	site	orbital
1	A	$ s, \uparrow\rangle, s, \downarrow\rangle$
2	B	$ s, \uparrow\rangle, s, \downarrow\rangle$

Table 6: Neighbor and bra-ket of each bond

#	head	tail	neighbor	head (bra)	tail (ket)
1	A	A	[1]	[s]	[s]
2	A	B	[1]	[s]	[s]
3	B	B	[1]	[s]	[s]

— Site in Unit Cell —

Sites in (conventional) cell (no plus set), SL = sublattice

Table 7: 'A' (#1) site cluster (1a), 2/m

SL	position (<i>s</i>)	mapping
1	[0.00000, 0.00000, 0.00000]	[1,2,3,4]

Table 8: 'B' (#2) site cluster (**1e**), 2/m

SL	position (s)	mapping
1	[0.50000, 0.50000, 0.00000]	[1,2,3,4]

— Bond in Unit Cell —

Bonds in (conventional) cell (no plus set): tail, head = (SL, plus set), (N)D = (non)directional (listed up to 5th neighbor at most)

Table 9: 1-th 'A'-'A' [1] (#1) bond cluster (**1a@1d**), ND, $|\mathbf{v}|=1.0$ (cartesian)

SL	vector (v)	center (c)	mapping	head	tail	R (primitive)
1	[-1.00000, 0.00000, 0.00000]	[0.50000, 0.00000, 0.00000]	[1,-2,-3,4]	(1,1)	(1,1)	[1,0,0]

Table 10: 1-th 'A'-'A' [2] (#2) bond cluster (**1a@1c**), ND, $|\mathbf{v}|=1.0$ (cartesian)

SL	vector (v)	center (c)	mapping	head	tail	R (primitive)
1	[0.00000, 0.00000,-1.00000]	[0.00000, 0.00000, 0.50000]	[1,-2,-3,4]	(1,1)	(1,1)	[0,0,1]

Table 11: 1-th 'A'-'B' [1] (#3) bond cluster (4a@4o), D, $|\mathbf{v}|=0.78102$ (cartesian)

SL	vector ($\mathbf{v}$)	center ($\mathbf{c}$)	mapping	head	tail	$\mathbf{R}$ (primitive)
1	[-0.50000, -0.50000, 0.00000]	[0.25000, 0.25000, 0.00000]	[1]	(1,1)	(1,1)	[0,0,0]
2	[0.50000, -0.50000, 0.00000]	[0.75000, 0.25000, 0.00000]	[2]	(1,1)	(1,1)	[-1,0,0]
3	[0.50000, 0.50000, 0.00000]	[0.75000, 0.75000, 0.00000]	[3]	(1,1)	(1,1)	[-1,-1,0]
4	[-0.50000, 0.50000, 0.00000]	[0.25000, 0.75000, 0.00000]	[4]	(1,1)	(1,1)	[0,-1,0]

Table 12: 1-th 'B'-'B' [1] (#4) bond cluster (1a@1b), ND, $|\mathbf{v}|=1.0$ (cartesian)

SL	vector ($\mathbf{v}$)	center ($\mathbf{c}$)	mapping	head	tail	$\mathbf{R}$ (primitive)
1	[-1.00000, 0.00000, 0.00000]	[0.00000, 0.50000, 0.00000]	[1,-2,-3,4]	(1,1)	(1,1)	[1,0,0]

Table 13: 1-th 'B'-'B' [2] (#5) bond cluster (1a@1h), ND, $|\mathbf{v}|=1.0$ (cartesian)

SL	vector ($\mathbf{v}$)	center ($\mathbf{c}$)	mapping	head	tail	$\mathbf{R}$ (primitive)
1	[0.00000, 0.00000, -1.00000]	[0.50000, 0.50000, 0.50000]	[1,-2,-3,4]	(1,1)	(1,1)	[0,0,1]