

MPG No. 8.3.26 $m'mm$ (m'mm setting) [Type III, orthorhombic] [T tensor]

* Rank 0 tensor. * Rank 1 tensor.

$$\begin{bmatrix} T_x & 0 & 0 \end{bmatrix}$$

$$T_x = \mathbb{T}_x^{(1)}$$

* Rank 2 tensor (s). * Rank 2 tensor (a). * Rank 3 tensor (ss).

$$\begin{bmatrix} T_{xxx} & 0 & 0 \\ T_{yyx} & 0 & 0 \\ T_{zzx} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & T_{zzx} \\ 0 & T_{yyx} & 0 \end{bmatrix}$$

$$T_{xxx} = 3\mathbb{T}_x^{(1)} + 2\mathbb{T}_x^{\alpha(1)}$$

$$T_{yyx} = \mathbb{T}_x^{(1)} - \mathbb{T}_x^{\alpha(1)} + \mathbb{T}_x^{\beta(1)}$$

$$T_{zzx} = \mathbb{T}_x^{(1)} - \mathbb{T}_x^{\alpha(1)} - \mathbb{T}_x^{\beta(1)}$$

* Rank 3 tensor (sa).

$$\begin{bmatrix} 0 & 0 & 0 \\ T_{yyx} & 0 & 0 \\ T_{zzx} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -\frac{T_{zzx}}{2} \\ 0 & -\frac{T_{yyx}}{2} & 0 \end{bmatrix}$$

$$T_{yyx} = 2\mathbb{M}_{yz}^{(1)} + 2\mathbb{T}_x^{(2)}$$

$$T_{zzx} = -2\mathbb{M}_{yz}^{(1)} + 2\mathbb{T}_x^{(2)}$$

$$\frac{T_{zzx}}{2} = -\mathbb{M}_{yz}^{(1)} + \mathbb{T}_x^{(2)}$$

$$\frac{T_{yyx}}{2} = \mathbb{M}_{yz}^{(1)} + \mathbb{T}_x^{(2)}$$

* Rank 3 tensor (a).

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & T_{zzx} \\ 0 & T_{xyy} & 0 \end{bmatrix}$$

$$T_{zzx} = \mathbb{M}_{yz}^{(2)} + \mathbb{T}_x^{(3)}$$

$$T_{xyy} = \mathbb{M}_{yz}^{(2)} - \mathbb{T}_x^{(3)}$$

* Rank 4 tensor (sss). * Rank 4 tensor (ssa). * Rank 4 tensor (aas). * Rank 4 tensor (aaa). * Rank 4 tensor (sa). * Rank 4 tensor (as). * Rank 4 tensor (s). * Rank 4 tensor (a). * Rank 4 tensor (t).