

MPG No. 8.6.125 $m'mm$ (mmm' setting) [Type III, orthorhombic] [T tensor]

* Rank 0 tensor. * Rank 1 tensor.

$$\begin{bmatrix} 0 & 0 & T_z \end{bmatrix}$$

$$T_z = \mathbb{T}_z^{(1)}$$

* Rank 2 tensor (s). * Rank 2 tensor (a). * Rank 3 tensor (ss).

$$\begin{bmatrix} 0 & 0 & T_{xxz} \\ 0 & 0 & T_{yyz} \\ 0 & 0 & T_{zzz} \\ 0 & T_{yyz} & 0 \\ T_{xxz} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$T_{xxz} = \mathbb{T}_z^{(1)} - \mathbb{T}_z^{\alpha(1)} + \mathbb{T}_z^{\beta(1)}$$

$$T_{yyz} = \mathbb{T}_z^{(1)} - \mathbb{T}_z^{\alpha(1)} - \mathbb{T}_z^{\beta(1)}$$

$$T_{zzz} = 3\mathbb{T}_z^{(1)} + 2\mathbb{T}_z^{\alpha(1)}$$

* Rank 3 tensor (sa).

$$\begin{bmatrix} 0 & 0 & T_{xxz} \\ 0 & 0 & T_{yyz} \\ 0 & 0 & 0 \\ 0 & -\frac{T_{yyz}}{2} & 0 \\ -\frac{T_{xxz}}{2} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$T_{xxz} = 2\mathbb{M}_{xy}^{(1)} + 2\mathbb{T}_z^{(2)}$$

$$T_{yyz} = -2\mathbb{M}_{xy}^{(1)} + 2\mathbb{T}_z^{(2)}$$

$$\frac{T_{yyz}}{2} = -\mathbb{M}_{xy}^{(1)} + \mathbb{T}_z^{(2)}$$

$$\frac{T_{xxz}}{2} = \mathbb{M}_{xy}^{(1)} + \mathbb{T}_z^{(2)}$$

* Rank 3 tensor (a).

$$\begin{bmatrix} 0 & T_{yzy} & 0 \\ T_{zxx} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$T_{yzy} = \mathbb{M}_{xy}^{(2)} + \mathbb{T}_z^{(3)}$$

$$T_{zxx} = \mathbb{M}_{xy}^{(2)} - \mathbb{T}_z^{(3)}$$

* Rank 4 tensor (sss). * Rank 4 tensor (ssa). * Rank 4 tensor (aas). * Rank 4 tensor (aaa). * Rank 4 tensor (sa). * Rank 4 tensor (as). * Rank 4 tensor (s). * Rank 4 tensor (a). * Rank 4 tensor (t).