

MPG No. 37.1.151 $\bar{6}m2$ (-62m setting) [Type I, hexagonal] [M tensor]

* Rank 0 tensor. * Rank 1 tensor. * Rank 2 tensor (s). * Rank 2 tensor (a). * Rank 3 tensor (ss). * Rank 3 tensor (sa).

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ M_{yzx} & 0 & 0 \\ 0 & -M_{yzx} & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$M_{yzx} = 3\mathbb{T}_u^{(1)}$$

* Rank 3 tensor (a).

$$\begin{bmatrix} M_{yzx} & 0 & 0 \\ 0 & M_{yzx} & 0 \\ 0 & 0 & M_{xyz} \end{bmatrix}$$

$$M_{yzx} = \mathbb{T}_0^{(1)} - \mathbb{T}_u^{(2)}$$

$$M_{xyz} = \mathbb{T}_0^{(1)} + 2\mathbb{T}_u^{(2)}$$

* Rank 4 tensor (sss).

$$\begin{bmatrix} 0 & 0 & 0 & M_{xxyz} & 0 & 0 \\ 0 & 0 & 0 & -M_{xxyz} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ M_{xxyz} & -M_{xxyz} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & M_{xxyz} \\ 0 & 0 & 0 & 0 & M_{xxyz} & 0 \end{bmatrix}$$

$$M_{xxyz} = \mathbb{M}_{4a}^{(1)}$$

* Rank 4 tensor (ssa).

$$\begin{bmatrix} 0 & 0 & 0 & M_{xxyz} & 0 & 0 \\ 0 & 0 & 0 & -M_{xxyz} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ -M_{xxyz} & M_{xxyz} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -M_{xxyz} \\ 0 & 0 & 0 & 0 & M_{xxyz} & 0 \end{bmatrix}$$

$$M_{xxyz} = 2\mathbb{T}_{3a}^{(1)}$$

* Rank 4 tensor (aas). * Rank 4 tensor (aaa). * Rank 4 tensor (sa).

$$\begin{bmatrix} M_{xxyz} & 0 & 0 \\ -M_{xxyz} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -M_{xxyz} & 0 \end{bmatrix}$$

$$M_{xxyz} = \mathbb{T}_{3a}^{(2)}$$

* Rank 4 tensor (as).

$$\begin{bmatrix} M_{yzxx} & -M_{yzxx} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -M_{yzxx} \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$M_{yzxx} = \mathbb{T}_{3a}^{(3)}$$

* Rank 4 tensor (s).

$$\begin{bmatrix} 0 & 0 & 0 & M_{xxyz} & 0 & 0 & M_{xxzy} & 0 & 0 \\ 0 & 0 & 0 & -M_{xxyz} & 0 & 0 & -M_{xxzy} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ M_{yzxx} & -M_{yzxx} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & M_{yzxx} & 0 & 0 & M_{yzxx} \\ 0 & 0 & 0 & 0 & M_{xxzy} & 0 & 0 & M_{xxzy} & 0 \end{bmatrix}$$

$$M_{xxyz} = \mathbb{M}_{4a}^{(1)} + 2\mathbb{T}_{3a}^{(1)} + \mathbb{T}_{3a}^{(2)}$$

$$M_{xxzy} = \mathbb{M}_{4a}^{(1)} + 2\mathbb{T}_{3a}^{(1)} - \mathbb{T}_{3a}^{(2)}$$

$$M_{yzxx} = \mathbb{M}_{4a}^{(1)} - 2\mathbb{T}_{3a}^{(1)}$$

* Rank 4 tensor (a).

$$\begin{bmatrix} M_{yzxx} & -M_{yzxx} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -M_{yzxx} & 0 & 0 & -M_{yzxx} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$M_{yzxx} = \mathbb{T}_{3a}^{(3)}$$

* Rank 4 tensor (t).

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & M_{yyyy} \\ 0 & 0 & 0 \\ 0 & M_{yyyy} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -M_{yyyy} \\ 0 & 0 & 0 \\ 0 & -M_{yyyy} & 0 \\ 0 & 0 & 0 \\ -M_{yyyy} & 0 & 0 \end{bmatrix}$$

$$M_{yyyy} = -\mathbb{M}_{4a}^{(1)}$$