

MPG No. 8.4.27 $m'm'm$ (m'm'm setting) [Type III, orthorhombic] [M tensor]

* Rank 0 tensor. * Rank 1 tensor.

$$\begin{bmatrix} 0 & 0 & M_z \end{bmatrix}$$

$$M_z = \mathbb{M}_z^{(1)}$$

* Rank 2 tensor (s). * Rank 2 tensor (a). * Rank 3 tensor (ss).

$$\begin{bmatrix} 0 & 0 & M_{xxz} \\ 0 & 0 & M_{yyz} \\ 0 & 0 & M_{zzz} \\ 0 & M_{yyz} & 0 \\ M_{xxz} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$M_{xxz} = \mathbb{M}_z^{(1)} - \mathbb{M}_z^{\alpha(1)} + \mathbb{M}_z^{\beta(1)}$$

$$M_{yyz} = \mathbb{M}_z^{(1)} - \mathbb{M}_z^{\alpha(1)} - \mathbb{M}_z^{\beta(1)}$$

$$M_{zzz} = 3\mathbb{M}_z^{(1)} + 2\mathbb{M}_z^{\alpha(1)}$$

* Rank 3 tensor (sa).

$$\begin{bmatrix} 0 & 0 & M_{xxz} \\ 0 & 0 & M_{yyz} \\ 0 & 0 & 0 \\ 0 & -\frac{M_{yyz}}{2} & 0 \\ -\frac{M_{xxz}}{2} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$M_{xxz} = 2\mathbb{M}_z^{(2)} + 2\mathbb{T}_{xy}^{(1)}$$

$$M_{yyz} = 2\mathbb{M}_z^{(2)} - 2\mathbb{T}_{xy}^{(1)}$$

$$\frac{M_{yyz}}{2} = \mathbb{M}_z^{(2)} - \mathbb{T}_{xy}^{(1)}$$

$$\frac{M_{xxz}}{2} = \mathbb{M}_z^{(2)} + \mathbb{T}_{xy}^{(1)}$$

* Rank 3 tensor (a).

$$\begin{bmatrix} 0 & M_{yzy} & 0 \\ M_{zxx} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$M_{yzy} = \mathbb{M}_z^{(3)} + \mathbb{T}_{xy}^{(2)}$$

$$M_{zxx} = -\mathbb{M}_z^{(3)} + \mathbb{T}_{xy}^{(2)}$$

* Rank 4 tensor (sss). * Rank 4 tensor (ssa). * Rank 4 tensor (aas). * Rank 4 tensor (aaa). * Rank 4 tensor (sa). * Rank 4 tensor (as). * Rank 4 tensor (s). * Rank 4 tensor (a). * Rank 4 tensor (t).