

MPG No. 8.9.128 $m'm'm$ (mm'm' setting) [Type III, orthorhombic] [M tensor]

* Rank 0 tensor. * Rank 1 tensor.

$$\begin{bmatrix} M_x & 0 & 0 \end{bmatrix}$$

$$M_x = \mathbb{M}_x^{(1)}$$

* Rank 2 tensor (s). * Rank 2 tensor (a). * Rank 3 tensor (ss).

$$\begin{bmatrix} M_{xxx} & 0 & 0 \\ M_{yyx} & 0 & 0 \\ M_{zzx} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & M_{zzx} \\ 0 & M_{yyx} & 0 \end{bmatrix}$$

$$M_{xxx} = 3\mathbb{M}_x^{(1)} + 2\mathbb{M}_x^{\alpha(1)}$$

$$M_{yyx} = \mathbb{M}_x^{(1)} - \mathbb{M}_x^{\alpha(1)} + \mathbb{M}_x^{\beta(1)}$$

$$M_{zzx} = \mathbb{M}_x^{(1)} - \mathbb{M}_x^{\alpha(1)} - \mathbb{M}_x^{\beta(1)}$$

* Rank 3 tensor (sa).

$$\begin{bmatrix} 0 & 0 & 0 \\ M_{yyx} & 0 & 0 \\ M_{zzx} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -\frac{M_{zzx}}{2} \\ 0 & -\frac{M_{yyx}}{2} & 0 \end{bmatrix}$$

$$M_{yyx} = 2\mathbb{M}_x^{(2)} + 2\mathbb{T}_{yz}^{(1)}$$

$$M_{zzx} = 2\mathbb{M}_x^{(2)} - 2\mathbb{T}_{yz}^{(1)}$$

$$\frac{M_{zzx}}{2} = \mathbb{M}_x^{(2)} - \mathbb{T}_{yz}^{(1)}$$

$$\frac{M_{yyx}}{2} = \mathbb{M}_x^{(2)} + \mathbb{T}_{yz}^{(1)}$$

* Rank 3 tensor (a).

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & M_{zzz} \\ 0 & M_{xyy} & 0 \end{bmatrix}$$

$$M_{zzz} = \mathbb{M}_x^{(3)} + \mathbb{T}_{yz}^{(2)}$$

$$M_{xyy} = -\mathbb{M}_x^{(3)} + \mathbb{T}_{yz}^{(2)}$$

* Rank 4 tensor (sss). * Rank 4 tensor (ssa). * Rank 4 tensor (aas). * Rank 4 tensor (aaa). * Rank 4 tensor (sa). * Rank 4 tensor (as). * Rank 4 tensor (s). * Rank 4 tensor (a). * Rank 4 tensor (t).