SG No. 2 C_{i}^{1} $P\bar{1}$ [triclinic]

* plus set: +[0, 0, 0]

Table 1: Wyckoff site: `1a`, site symmetry: `-1`

No.	position	mapping
1	[0, 0, 0]	`[1,2]`

Table 2: Wyckoff site: `1b`, site symmetry: `-1`

No.	position	mapping
1	$\left[0, 0, \frac{1}{2}\right]$	`[1,2]`

Table 3: Wyckoff site: `1c`, site symmetry: `-1`

No.	position	mapping
1	$\left[0, \frac{1}{2}, 0\right]$	`[1,2]`

Table 4: Wyckoff site: `1d`, site symmetry: `-1`

No.	position	mapping
1	$\left[\frac{1}{2}, 0, 0\right]$	`[1,2]`

Table 5: Wyckoff site: `1e`, site symmetry: `-1`

No.	position	mapping
1	$\left[\frac{1}{2}, \frac{1}{2}, 0\right]$	`[1,2]`

Table 6: Wyckoff site: `1f`, site symmetry: `-1`

No.	position	mapping
1	$\left[\frac{1}{2}, 0, \frac{1}{2}\right]$	`[1,2]`

Table 7: Wyckoff site: `1g`, site symmetry: `-1`

No.	position	mapping
1	$\left[0, \frac{1}{2}, \frac{1}{2}\right]$	`[1,2]`

Table 8: Wyckoff site: `1h`, site symmetry: `-1`

No.	position	mapping
1	$\left[\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right]$	`[1,2]`

Table 9: Wyckoff site: `2i`, site symmetry: `1`

No.	position	mapping
1	$[x,\, y,\, z]$	`[1]`
2	$[-x,\, -y,\, -z]$	`[2]`