


## CANDIDATE CORRECTIONS TO MANUSCRIPT

- [Page 4]:
  - Figure 1: Phylogeny of metazoan species from which **HIF pathway gene** anchor sequences etc.

Protein sequences from 25 species across 13 animal phyla were used to construct **p-HMMs for HIF- $\alpha$ , EGLN and VHL ortholog mining** across the proteomes of 9 platyhelminth species.

**Priapulida clades with Ecdysozoa, not Lophotrochozoa.**


- [Page 5]:
  - Figure 2: **AMENDED**



- [Page 14]:
  - **For HIF- $\alpha$  ortholog mining**, alignment using MAFFT iterative local-pair refinement was selected to maximise positional homology etc.

- [Page 16]:
  - A single candidate VHL ortholog in the **trematode** *Clonorchis sinensis*.
  - Subsequent maximum-likelihood phylogenetic inference placed the *Clonorchis* sequence **in a phylogenetic tree of high-confidence metazoan VHL sequences** as an isolated branch outside other metazoan VHL clades etc.

- [Page 21]:
  - Figure 11A: **AMENDED**



- Figure 11C: Heatmaps **beside** the module assignments etc.
  - Figure 11D: Temporal trajectories of module expression showing mean FPKM **value** of genes etc.
- [Page 31]: These indirect sources of evidence are critical in a non-model metazoan with no transgenic technologies available, precluding the use of **CRE-reporter** constructs for directly testing the function and mapping the target genes of CREs.
  - [Page 32]: Figure 18: Base-pair resolution view of hypoxia-specific Klf10/11 footprint in **h1SMcG0021003**.