

Dyslipidaemia in Rheumatoid Arthritis: The Role of Inflammation, Drugs, Lifestyle and Genetic Factors

Tracey E. Toms¹, Deborah P. Symmons² and George D. Kitas^{1,2,*}

¹Department of Rheumatology, Dudley Group of Hospitals NHS Trust, Russells Hall Hospital, Dudley, West Midlands, UK; ²Arthritis Research Campaign Epidemiology Unit, Manchester University, Manchester, UK

Abstract: Rheumatoid arthritis (RA) associates with excess cardiovascular morbidity and mortality, resulting in significantly shortened lifespan. Traditional risk factors (e.g. dyslipidaemia and hypertension) and novel risk factors (e.g. systemic inflammation) contribute to the development of cardiovascular disease (CVD) in RA. In the general population, dyslipidaemia has been found to be central to the development of CVD, playing an important role in all stages of atherosclerotic plaque formation. In RA, lipid metabolism may be altered by systemic inflammation, environmental lifestyle factors, drug therapy and several genetic factors. This may result in changes in overall lipid levels, as well as modifications of lipid/lipoprotein structure and function. In this review, we discuss lipid abnormalities specifically in the context of RA and highlight the potential impact of inflammation, genetic factors, lifestyle, and anti-rheumatic drugs on lipid metabolism.

Keywords: Rheumatoid arthritis, dyslipidaemia, inflammation, drugs, genetics, lifestyle, environment.

INTRODUCTION

Rheumatoid arthritis (RA) associates with significantly increased cardiovascular disease (CVD) morbidity and mortality compared to the general population. CVD in RA is multi-factorial arising as a consequence of an interplay between established CVD risk factors (e.g. hypertension, dyslipidaemia, diabetes, smoking) [1], genetic predisposition, and novel disease specific factors, with systemic inflammation being the key player [2]. Several of the traditional risk factors have now been studied in some detail, establishing their role in initiating and perpetuating CVD in RA. Dyslipidaemia is a major factor that requires further investigation specifically in RA.

For decades, abnormalities of the lipid profile such as increased levels of low density lipoproteins (LDL) have been recognised as strong predictors of CVD in the general population. In RA, the lipid profile appears to superficially improve with suppression of components such as LDL and total cholesterol (TC) [3-5]. However, disproportionate suppression of other components of the lipid profile such as high density lipoproteins (HDL), as well intricate modifications of lipids, lipoproteins and enzymes involved in lipid metabolism are likely to increase the risk of CVD in RA [6, 7].

In this review, we discuss lipid abnormalities specifically in the context of RA and highlight the potential impact of inflammation, genetic factors, lifestyle, and anti-rheumatic drugs on lipid metabolism.

PREVALENCE AND ASSOCIATIONS OF DYSLIPIDAEMIA IN RA

Dyslipidaemia in RA is likely to be governed by many factors, including inflammatory disease activity [8], reduced physical activity secondary to pain and disability [9] as well as drug therapy [7]. Although, collectively these factors are likely to exert a significant influence on the lipid profile of RA patients, the additional potential contribution of genetic factors controlling lipid metabolism has not been addressed. This may explain why the prevalence of dyslipidaemia in RA varies between populations.

Several studies have commented on the prevalence of dyslipidaemia in RA. The first study assessed the prevalence of dyslipidaemia in 87 patients with inflammatory arthritis [10]. According to the United States (US) National Cholesterol Education Program (NCEP) guidelines [11], 55% of inflammatory arthritis patients were dyslipidaemic compared with 8% of controls. Unfortunately, these figures are not specific to RA, but encompass a broad spectrum of inflammatory joint disease, (RA, spondyloarthropathies and undifferentiated inflammatory arthritis) exhibiting varying degrees of inflammation and different disease-specific characteristics. The second study, was carried out on 60 RA males patients [12], thus not reflecting the typical disease population. The investigators reported 68% of patients had serum levels of TC, HDL or LDL that would be considered as risk factors for the development of atherosclerosis according to NCEP criteria. In our own series of 400 patients meeting American College of Rheumatology (ACR) criteria for RA, the prevalence of dyslipidaemia according to the NCEP criteria (TC ≥ 6.2 mmol/L or LDL ≥ 4.13 mmol/L or HDL < 1.03 mmol/L or TG ≥ 1.7 mmol/L, or taking lipid lowering therapy) was 51.3% [13].

*Address correspondence to this author at the Department of Rheumatology, Dudley Group of Hospitals NHS Trust, Russells Hall Hospital, Pensnett Road, Dudley, West Midlands, DY1 2HQ, UK; Tel: +44-1384-244842; Fax: +44-1384-244808; E-mail: gd.kitas@dgoh.nhs.uk or g.d.kitas@bham.ac.uk

Dyslipidaemia appears to be present in RA patients with both early [14] and advanced disease [15]. While the inflammatory burden can be partially blamed for dyslipidaemia in early RA, there is also evidence to suggest that lipid profile may already be altered prior to the onset of disease. One retrospective study performed in the Netherlands on 1078 blood donors, identified 79 patients who later developed RA, and compared their lipid profiles with that of controls from blood samples taken at least 10 years before the onset of RA [16]. Patients who later developed RA were found to have lower levels of HDL, and higher levels of TC, TG and apolipoprotein B (ApoB). This suggests that alterations in the lipid profile may render people more susceptible to the development of RA [17], that RA patients are genetically predisposed to the development of dyslipidaemia or that the transcription of these genes is altered by persistent low-grade inflammation in patients with subclinical RA.

ALTERATIONS IN THE LIPID PROFILE IN RA

In 1963, London *et al.* [18] described a negative association between TC levels and RA disease activity, but no data were produced regarding the effects of inflammation on other lipid parameters. Nevertheless, this study acted as a springboard for future studies to evaluate the contribution of dyslipidaemia to the cardiovascular risk seen in RA. Currently, the most widely reported abnormality of the lipid profile associated with active RA is suppression of HDL levels [14, 19-21]. Although a fall in TC levels has been observed [18], the drop is not as substantial as that seen in HDL levels, thus producing an increased atherogenic index (TC:HDL ratio) [14]. The atherogenic index may be even higher if TC levels increase as claimed by several investigators [19, 22]. There is some controversy over the effects of inflammation on TGs and LDL, some studies show that high disease activity associates with higher TGs [19] while other studies suggest that TGs are reduced [21]. Similar to TGs, the precise relationship between LDL and inflammation also remains unclear [19, 23]. Much of this controversy can be explained by small study size and differences in the populations studied.

To date, the National Health and Nutrition Examination Survey (NHANES III) is the largest study to compare lipid profiles of untreated RA patients with a control group [7]. This study recruited 128 patients with RA, but only 104 patients were untreated with disease modifying anti-rheumatic drugs (DMARDs) or glucocorticoids (GCs) and thus included in the analysis. Although all major components of the lipid profile were analysed, only HDL and ApoA-I levels were found to be lower in the RA patients.

There is growing evidence that ratios of lipid components and apolipoproteins have a higher predictive value of first myocardial infarction than individual components of the lipid profile [24]. Ratios of particular interest include: TC:HDL (discussed above), LDL:HDL and Apo B:Apo A-I. In RA, not only are all these 3 ratios increased [14], but they may offer a more reliable way of assessing the lipid profile by overcoming individual fluctuations in lipids occurring as part of disease flares.

HDL and LDL particles are categorised according to their size and density. This is clinically relevant as small dense

LDL particles (LDL3) more readily infiltrate the endothelium and are more prone to oxidative change than their larger counterparts [25, 26], whereas smaller HDL particles (HDL2) are more successful at performing reverse cholesterol transport and thus confer greater cardio-protection [27, 28]. In RA, only 1 small study has assessed the characteristics of lipoprotein sub-fractions [6]. This study, on 31 RA patients, demonstrated significantly higher levels of small dense LDL and lower levels of small HDL particles compared with controls. Interestingly, there were no changes in the levels of basic components of the lipid profile, except LDL levels, which were lower in the RA group.

Lipoprotein (a) [Lp(a)], is a LDL particle in which ApoB-100 is bound to lipoprotein a. Current evidence suggests Lp(a) may be a key factor in the development of atherosclerosis [29-31]. The recognition of Lp(a) as an independent cardiovascular risk factor in the general population [32] has sparked interest into the role of this lipoprotein in RA. Levels of Lp(a) have been found to be increased in RA patients [14, 19, 33]. The significant increases in Lp(a) could be purely as a direct result of inflammation or may be due to increased genetic expression [34]. Irrespective of these findings, it still is unclear whether increased levels of Lp(a) accelerate atherosclerosis in RA. Interestingly, although suppression of inflammation with drug therapy (DMARDs, GCs or biologic agents) can result in the reversal of many of the lipid changes described in active RA, Lp(a) levels appear to remain consistently elevated [33]. Table 1 summarises the findings from all the available studies reporting lipid changes in untreated RA.

RA has been associated with high levels of oxidative stress [35]. The increased oxidative load may contribute to articular tissue damage and perpetuates the inflammatory process [36], but may also be strongly implicated in the pathophysiology of CVD in RA [37, 38]. In RA, it has been shown that LDL within the synovial fluid of an inflamed rheumatoid joint is prone to oxidative modification [39]. More recently it has also been reported that increased levels of oxidised LDL are seen in the plasma of RA patients with ischaemic heart disease [40]. The trigger for LDL oxidation in RA is not fully understood, particularly as a study suggests that inflammation does not play a significant role [41].

In summary, RA patients with active disease develop a pro-atherogenic lipid profile, with low HDL levels and raised atherogenic indices. The additional influence of the other lipid components (LDL and TG), and lipid sub fractions on cardiovascular risk in RA requires further investigation.

RA AND THE METABOLIC SYNDROME

The metabolic syndrome (MetS) is a cluster of traditional cardiovascular risk factors (obesity, glucose intolerance, hypertension, hypertriglyceridaemia, and low levels of HDL), thought to confer cardiovascular risk beyond the sum of its individual components [44]. In RA, several of the individual components of the MetS may be altered through disease specific and inflammation-mediated mechanisms. For example, many RA patients develop rheumatoid cachexia, a condition characterised by an increased fat:muscle mass ratio, while maintaining a constant weight [45]. Thus, it has been suggested that in RA lower cut-off levels should be used to de-

Table 1. Studies of Lipid Profiles in Patients with Untreated Rheumatoid Arthritis

| Study | Study Design | TC | HDL | LDL | TG | Lp(a) | Apo A1 | TC:HDL Ratio | LDL:HDL Ratio | apoB:apo A1 Ratio |
|-----------------|--|-----------------------|----------------------|-----------------------|----------------------|-----------------------------|---------------------|----------------------|----------------------|-----------------------|
| Georgiadis [42] | 40 RA <1 yr 45 controls | ↑* | ↓ | ↑* | ↑* | N/S | ↓* | ↑* | ↑* | N/D |
| Georgiadis [43] | 58 RA <1 yr 63 controls | ↑ | ↓* | ↑ | ↑ | N/S | N/S | ↑ | ↑* | N/S |
| Dursunoglu [19] | 87 female RA 50 controls | N/D | ↓* | ↑ | ↑* | ↑* | N/S | ↑* | ↑* | N/S |
| Choi [7] | 104 RA age >60 4758 controls | N/D | ↓ | N/S | N/D | N/S | ↓ | ↑ | N/S | N/S |
| Yoo [8] | 184 RA 161 controls | ↓* females ↓ males | ↓females ↓* males | ↑* females ↑ males | ↑ females ↑ males | ↑* fe- males ↑* males | ↓females ↓ males | ↑ females ↓ males | ↑ females ↑ males | ↑ females ↑* males |
| Lee [3] | 21 RA 19 controls | ↓ | ↓* | N/D | N/D | ↑ | N/S | N/D | N/S | N/S |
| Park [14] | 42 RA 42 controls | | ↓* | | | ↑* | ↓* | ↑* | ↑* | ↑* |
| Seriolo [4] | 137 RA 78 controls | ↓* | N/D | ↓* | N/D | ↑* | N/D | N/S | N/S | N/D |
| Lakator [22] | 129 RA (77 GC, 52 NSAID) 1374 controls | ↑ | ↓* | ↑* | ↓* | N/S | N/S | N/S | N/S | N/S |
| Svenson [5] | 48 RA 21 sero-ve SA | ↓ | ↓* | ↓* | ↓* | N/S | N/S | N/S | N/S | N/S |

* : significant change, N/D: no difference between RA and controls, N/S: not studied, SA: spondyloarthropathies, RA: Rheumatoid Arthritis, GC: glucocorticoids, NSAID: non steroidal anti inflammatory drugs, TC: total cholesterol, HDL: high density lipoproteins, TG: triglycerides, LDL: low density lipoproteins, ApoA: apolipoprotein A, ApoB: apolipoprotein B, yr: years.

fine obesity using the body mass index [46]. In view of such modifications to the individual components, it is not surprising that the MetS has been found to be highly prevalent in RA [47-49]. As inflammation clearly has a central role in the development of the MetS [50], it would be expected that suppression of the inflammatory burden through the use of disease modifying drugs would result in a reduced prevalence of the MetS. Methotrexate has been reported to have cardioprotective properties [51], and more recently has been shown to associate with a lower prevalence of the MetS [49, 52]. Interestingly, recent data would suggest that the 'protective' effects of methotrexate may be due to a drug specific mechanism rather than a generic anti-inflammatory effect [52, 53]. However, this association needs to be explored further in specifically designed longitudinal studies. Long-term GC use associates with many undesirable side effects, including central obesity [54], dyslipidaemia [55], glucose intolerance [56], and hypertension [57]. However, in RA, long-term GC use has not been found to associate with an increased prevalence of the MetS [53].

The rheumatological community is becoming increasingly aware of the need to address cardiovascular risk in RA patients. Tackling the MetS in RA is complex, requiring both

suppression of systemic inflammation through the use of DMARD (in particular methotrexate) and GC use, as well as aggressive management of each of the individual components of the MetS e.g. by the use of appropriate anti-hypertensive therapy for hypertension and/or statins for dyslipidaemia. This is addressed later in this review. However, the importance of lifestyle modification to include weight control and increased physical activity cannot be over-emphasised: this is at least as important in patients with RA as it is in other at risk groups and in the general population.

INFLAMMATION-MEDIATED MECHANISMS CONTROLLING LIPID METABOLISM

The Effects of Inflammation on HDL Metabolism

HDL confers multiple cardiovascular benefits, through its anti-oxidant properties and its ability to perform reverse cholesterol transport [58, 59]. It is now well recognised that high levels of HDL are cardio-protective, whereas low levels act as an independent cardiovascular risk factor [60-63]. Studies in the general population have provided convincing evidence that increasing HDL levels through drug intervention and

lifestyle modification can slow the progression of atherosclerosis and improve cardiovascular outcomes [64-66].

HDL is of particular interest in the context of active inflammation, as its levels can dramatically fall, to a far greater extent than the changes seen in other components of the lipid profile. In acute inflammation, such as infection (bacterial and viral), HDL levels closely correspond to the degree of inflammatory burden, as levels are rapidly suppressed following the onset of infection and their return to normal appears to shadow the patient's recovery [67]. Although, the majority of studies report normalisation of HDL levels within 4 weeks of the onset of infection [67-69], 1 study has shown that the reduced levels persist up to 6 months post infection [70]. In chronic inflammatory conditions, such as RA, HDL levels are persistently low. However, levels may still fluctuate as a consequence of alterations in disease activity [71], drug usage [72, 73] and physical activity levels [9].

The Effect of Acute Phase Proteins of HDL Metabolism

There is accumulating evidence suggesting that many of the anti-oxidant and anti-atherogenic properties of HDL are lost due to direct and indirect effects of inflammation. The inflammatory process triggers the synthesis of many plasma proteins by the liver [74, 75]. Serum amyloid A (SAA) and C-reactive protein (CRP) are 2 of the key plasma proteins whose production is greatly enhanced during inflammation. The net increase in SAA occurs as a result of increased gene transcription [76], and leads to alterations in HDL composition and function. Large amounts of SAA become bound to HDL (mainly HDL3) following the displacement of ApoA-I and ApoAII [77]. The composition of this remodelled acute phase HDL (AP-HDL) is also altered, becoming depleted in cholesterol esters and laden with TG, free cholesterol and fatty acids [78, 79]. The size of AP-HDL is larger than conventional HDL, although the density remains comparable [80]. The modification of the HDL structure to incorporate SAA directly impacts upon its ability to carry out reverse cholesterol transport, as lecithin-cholesterol acyltransferase (LCAT), an enzyme responsible for the esterification of HDL, requires the presence of ApoA-I to be activated. Alongside this, the HDL/SAA structure has an increased affinity for macrophages and a reduced affinity for hepatocytes compared with the unmodified HDL structure [81]. These changes can be partly attributed to alterations in the number of binding sites for the HDL/SAA complex, as inflammation has been shown to increase the number of HDL/SAA binding sites on macrophages and decrease the number on hepatocytes [81]. However, it has been shown that SAA must constitute more than half of the HDL protein in order for cholesterol efflux to become compromised [82].

Interest has also developed around the role of 2 other acute phase reactants, namely, secretory phospholipase A2 (sPLA2), known to possess the capacity to remodel HDL [83], and ceruloplasmin, a copper transporting protein with pro-oxidant properties [84]. Inflammation induces elevations in the plasma concentrations of sPLA2 and this has been linked to smaller HDL particle size, a reduction in HDL cholesterol and apoA-I levels, and an increase in HDL TGs in transgenic mice [85]. Acute phase HDL is also susceptible to enrichment with ceruloplasmin, resulting in a reduced ability

to protect LDL against oxidative modification [84, 86]. However, this theory is under scrutiny, as other studies conclude that ceruloplasmin exhibits anti-oxidant properties [87]. The discrepancies regarding the properties of ceruloplasmin may be explained by differences in its structure, as the removal of 1 copper atom appears to alter its function from anti-oxidant to pro-oxidant [88].

Transferrin, a plasma protein involved in iron transport, can be found in association with HDL [89]. It is thought that the usual role of this metal binding protein is to protect LDL against oxidation [89]. However, during inflammation, levels of transferrin fall [90] and thus further predispose the host to a proatherogenic environment.

The Effect of Inflammation-Mediated Enzymatic Change on HDL Metabolism

Several enzymes fundamental to the metabolism of HDL are affected by inflammation. Hepatic lipase (HL), an enzyme expressed primarily in the liver, converts larger HDL particles (HDL2) into smaller (HDL3) particles, facilitating cholesterol uptake from cells [91]. This process occurs largely due to the hydrolysis of TGs and phospholipids within HDL2. HL also plays a key role in many other aspects of lipoprotein metabolism, including assisting hepatic uptake of HDL and LDL particles by acting as a ligand [92] and aiding reverse cholesterol transport by promoting uptake of HDL cholesterol esters by the scavenger receptor B1 [93]. Animal [94] and human studies [95-97] have shown HL levels to be reduced by the inflammatory process, thus inhibiting its normal functions and the production of a more pro-atherogenic environment.

Reverse cholesterol transport is a complex cascade of events, which requires the presence of many factors in order for the process to be completed efficiently. Cholesterol ester transfer protein (CETP) plays a pivotal role, providing a pathway for cholesterol esters to be transferred from HDL to lipoproteins rich in ApoB, such as very low density lipoproteins (VLDL) and LDL, allowing cholesterol to ultimately be cleared by the liver [98]. Inflammation has been shown to indirectly impact upon CETP activity by reducing mRNA expression in transgenic mice [99]. However, further studies in this field are required to confirm these findings in the context of chronic inflammation in humans. Phospholipid transfer protein (PLTP) is another key protein required for successful reverse cholesterol transport, whose activity is suppressed by the inflammatory process [100]. The resultant limited activity of PLTP means that essential actions, such as mediating the exchange of cholesterol between TG rich particles and HDL, cannot be carried out. Thus, alterations in PLTP activity can be held partly responsible for lower HDL levels observed during inflammation [101].

Inflammation-mediated variations in the enzymatic content of HDL have also been observed, including reductions of paraoxonase (PON-1) and elevated levels of platelet-activating factor acetylhydrolase (PAF-AH) [86]. Deficiency of PON-1 within HDL renders them susceptible to oxidation and can ultimately convert HDL into a pro-oxidant, pro-inflammatory complex [86, 102]. PAF-AH is found in relation to both LDL and HDL particles. Elevated PAF-AH levels have been observed in a broad spectrum of inflammatory

conditions, including the human immunodeficiency virus (HIV) [103] and RA [104]. However, it remains unclear in humans whether PAF-AH activity is increased during inflammation in both HDL and LDL particles [103, 105]. It is postulated that if PAF-AH activity is increased within HDL particles, this may confer mainly anti-atherogenic properties, as it may protect against the oxidation of LDL, but also produce proatherogenic effects by escalating the production of lysophosphatidylcholine (LPC) [106]. Occasional published reports have demonstrated a negative association between inflammatory burden and PAF-AH levels [107, 108]. However, such studies may lack significant power due to small study size.

It is clear from this evidence that it is not just the reduction in HDL that is an important risk factor for coronary heart disease in heightened inflammatory states, such as RA, but that there are multiple other modifications in its function that may also have an additive effect (Fig. 1).

The Effects of Inflammation on Triglyceride Metabolism

The relative contribution of elevated TG levels to the development of cardiovascular disease remains controversial. Although there is growing evidence that hypertriglyceridaemia associates with an increased CVD risk [60, 109, 110], it remains difficult to classify it as an independent risk factor, as alterations in the levels occur in conjunction with changes in HDL levels [111].

For many decades, studies have demonstrated dramatic elevations in TG levels in response to infection as well as acute and chronic inflammation [19, 68, 112]. Some studies have reported these changes to be specific to the infective organism (e.g. Gram negative bacteria) [112], whilst others report a significant increase irrespective of the factor triggering the immune activation [68, 96]. TGs are primarily transported in VLDL, therefore hypertriglyceridaemia occurs as a result of either overproduction or impaired clearance of VLDL. The inflammatory process interferes with the normal metabolism of TGs through the release of multiple cytokines and alterations in enzymatic activity (see below).

The Effects of Cytokines on TG Metabolism

Numerous cytokines released during inflammation may hinder TG metabolism. TNF- α is released in vast quantities during active inflammation. Elevations in TNF- α levels may disturb lipoprotein metabolism by decreasing lipoprotein lipase (LPL) activity, reducing liver metabolism [113] and modifying the composition of lipoprotein particles [114]. Studies in rats have clearly demonstrated that the administration of exogenous TNF- α results in doubling of baseline plasma TG levels [115]. Subsequent studies in humans have produced data confirming a positive correlation between TNF- α and TG levels. However, much of this data has been generated from studies carried out on healthy subjects [116] and patients without active inflammation [117]. Although studies assessing this correlation on the background of




Fig. (1). The effects of inflammation on the structure, composition and function of lipids.

chronic inflammation are few, the available data confirm a striking positive association in patients with systemic lupus erythematosus (SLE) [118].

In RA, the advent of new biologic therapies specifically designed to target TNF- α has enabled this issue to be addressed from another angle. Interestingly, there is no convincing evidence to support the reverse association, (i.e. that as levels of TNF- α fall or TNF- α is inactivated, levels of TG also reduce). The precise effects of anti-TNF agents on TG levels are an issue of much debate. However, the vast majority of current published data indicates that treatment either does not impact upon TG levels [4, 119, 120] or may actually result in a paradoxical rise [121, 122]. These unexpected findings may be explained by small study size, inadequate adjustment for other potential confounders or by an unknown mechanism. Further large-scale studies are required to elucidate the exact effects of anti-TNF agents on cardiovascular risk factors, such as dyslipidaemia.

Over recent years, interest in the inflammatory cytokine IL-6 has escalated, due to the recognition of its atherogenic properties [123], and its potential as a therapeutic target both in cardiovascular diseases and in RA [124, 125]. A growing bank of evidence indicates that IL-6 levels influence lipoprotein metabolism [126], generating particular effects on the concentrations of TGs and HDL. IL-6 levels correlate positively with TG, and negatively with HDL levels [127, 128]. These findings may be particularly important in the context of the metabolic syndrome (MetS) [129], as emerging research indicates a potential role for IL-6 in the development of the MetS [130, 131]. Despite the strong evidence to support a positive relationship between cytokine exposure and TG levels, there are also a few studies demonstrating a negative association [132].

Drug therapy again offers a further insight in to the relationship between IL-6 and TG levels, as lipid lowering therapies such as statins display a dual mode of action by lowering TG levels and suppressing IL-6 through their anti-inflammatory properties [133]. The future use of new monoclonal antibodies specifically targeting IL-6 [125] in chronic inflammatory diseases such as RA, will allow this issue to be addressed in more detail.

The Effects of Inflammation-Mediated Enzymatic Change on TG Metabolism

LPL is an intravascular enzyme specifically found in endothelial cells. It is a multifunctional enzyme, displaying the ability to mediate lipoprotein uptake and to catalyse the hydrolysis of TGs within circulating VLDL and chylomicrons [134]. The explosive release of cytokines such as TNF- α and IL-6 occurring during inflammation leads to a reduction in the levels of LPL and HL *via* down regulation of gene expression at the transcriptional level [135, 136], which ultimately leads to a reduced clearance of TG-rich particles. The net result of these modifications takes some time to accrue. Thus, such changes only contribute to hypertriglyceridaemia in the setting of persistent chronic inflammation [137].

Peroxisome proliferator-activated receptors (PPAR) are ligand-activated transcription factors, important in several aspects of lipid metabolism. Of particular interest is PPAR-

α , regulating lipid metabolism both at the intracellular and extracellular level [138]. LPL has been shown to activate PPARs, through a preferential action on TG-rich particles (VLDL) [139]. Therefore, inflammation-mediated suppression of LPL levels inhibits PPAR activation and ultimately contributes to hypertriglyceridaemia [136]. The pathway of PPAR activation has long been used as a lipid lowering therapeutic target. Fibrates are PPAR- α agonists, used to treat dyslipidaemia by reducing TG levels and increasing HDL levels [140]. HDL levels increase with fibrate therapy due to PPAR activated up-regulation of ApoA-I and ApoA-II gene transcription [141, 142].

The Effects of Inflammation on LDL Metabolism

Epidemiological studies have shown elevated LDL levels to be one of the strongest predictors of coronary artery disease [143]. To combat this risk, guidelines have been developed, with LDL as the primary focus for lipid lowering therapy [11, 144]. Several large scale studies in the general population have demonstrated that intensive lipid lowering therapy confers greater CV benefits than moderate therapy [145-147]. Furthermore, such studies demonstrate continued clinical benefit even when LDL levels are lowered below the current recommended treatment goals [146]. Over the last 50 years a steady decline in LDL levels has been observed amongst Americans, according to the NHANES [148]. These changes may have occurred secondary to an increased prescription of lipid lowering medication and an overall more aggressive approach to the management of dyslipidaemia.

Inflammation lowers LDL levels [20, 149]. Superficially, this appears to produce a less atherogenic environment. However, by delving deeper it becomes apparent that inflammation mediated structural/oxidative changes may promote atherogenesis *via* the development of pro-inflammatory, atherogenic LDL particles [150, 151]. In fact, persistent inflammation such as that seen in RA, may fuel a vicious circle of oxidation and inflammation (Fig. 2).

Inflammation-Mediated Structural Changes of LDL

Low-density lipoproteins are sub-classified according to size and density. A predominance of small dense LDL particles is associated with a 3-5 times increased risk of CHD (the Quebec cardiovascular study) [152]. Studies in patients affected by inflammatory disorders, including psoriatic arthritis [153], the acquired immunodeficiency syndrome (AIDS) [154] and RA [6] have demonstrated a preponderance of small dense LDL particles compared with control groups. At present, information regarding the precise mechanisms behind these changes is limited and further research in this area is required.

Inflammation-Mediated Oxidative Changes of LDL

During inflammation there is release of reactive oxygen species (ROS) from activated leucocytes. Elevated ROS levels overwhelm the host's usual antioxidant mechanisms, resulting in damage to cells and lipid peroxidation [155]. The alterations in LDL composition have been blamed almost entirely for the accelerated oxidative modification during inflammation. In animal models, inflammation induced by the introduction of lipopolysaccharide (a major component




Fig. (2). The vicious cycle of inflammation and oxidised LDL.

of the cell wall of gram negative bacteria), has been shown to increase markers of lipid peroxidation [156]. Other studies carried out in human subjects with inflammatory disorders have confirmed similar elevations in markers of lipid peroxidation and an increased oxidative susceptibility of LDL [157]. Conflicting results are reported by Paredes *et al.*, who found no differences in lipid peroxidation between patients with active RA and the control group [158]. This discrepancy may be explained by the small study size, the degree of inflammatory burden or by the methods used to identify oxidative change.

The Effects of Acute Phase Proteins on LDL Metabolism

The pro-atherogenic properties of LDL are enhanced by CRP and sPLA2 during inflammation [159]. The inflammation mediated increase in sPLA2 activity produces small dense LDL particles that have a surface layer relatively deficient in phospholipids [160]. Such changes in the outer-most layer of LDL increase its ability to interact with arterial proteoglycans, perhaps further enhancing its uptake into the arterial wall and the formation of atherosclerotic plaques. sPLA2 may also indirectly promote LDL oxidation, through the production of fatty acids. Fatty acids are prone to oxidative change, and may subsequently induce oxidation of LDL [161]. Once oxidised, LDL is readily taken up by macrophages, and an abundance of CRP is recognised to facilitate this process, and promote subsequent foam cell formation [162].

The Effects of Inflammation-Mediated Enzymatic Change on LDL Metabolism

Endothelial lipase (EL) is a newly identified member of the triglyceride lipase family [163]. Apart from its role in HDL homeostasis, EL may enhance lipid uptake into the vascular endothelium *via* its bridging function [164]. Al-

though, it is widely accepted that EL is up-regulated during inflammation [165, 166], the exact effect that this has on LDL is still a matter of debate. One of the earliest studies to address this issue was carried out in LDL receptor deficient mice and demonstrated that hepatic expression of EL leads to a reduction in serum LDL levels [163]. In contrast, a later study in EL knockout mice has reported a massive increase (90%) in LDL cholesterol levels [167]. Two further studies, have also tried to identify the role of EL in LDL metabolism. The first failed to demonstrate any effect of EL on LDL levels in the mouse model [168], while the second demonstrated that EL promotes LDL uptake by macrophages [169].

EFFECTS OF ENVIRONMENTAL LIFESTYLE ON THE LIPID PROFILE

Variations in lifestyle and exposure to environmental factors can significantly alter the lipid profile. In RA, some of these factors may be modified as a consequence of complications of the disease. For example, regular physical activity has been shown to produce a less atherogenic lipid profile [170]. However, many RA patients have a limited capacity for physical activity due to the functional limitations produced by joint deformity and pain, and/or organ involvement (breathlessness from lung disease, or cardiac involvement). In this section we discuss the impact of environmental and lifestyle factors on the lipid profile and address how these are further altered by the presence of RA (see Fig. 3)

Seasonal Variation

Studies have demonstrated both biological variation (i.e. normal day to day variation) [171] and cyclical seasonal variation in plasma lipid and lipoprotein levels [172, 173]. Within patient biological variation is in the order of 6-7% for TC and HDL, 8-10% for LDL and 20-30% for TG [171,




Fig. (3). The potential interplay of lifestyle factors in the development of dyslipidaemia in RA.

174]. The most widely reported seasonal changes are elevations in TC and LDL levels in winter [175], with peak levels being reached during January [173, 176]. Apolipoprotein levels also demonstrate seasonal fluctuations, with ApoA-I and A-II levels markedly elevated in the summer [177]. These findings have been shown to be independent of country of origin, ethnicity, age, sex and baseline lipids. Interestingly LDL demonstrates an increased susceptibility to oxidation during winter (January) and a reduced susceptibility during summer (June/July) [178]. Some studies also report winter elevations in HDL and TG levels [176, 179]. The precise mechanisms controlling these changes remain elusive. However, several hypotheses have been put forward. These include: (a) a haemodilution effect due to secondary mobilisation of fluid from the interstitial to the intravascular compartment due to heat acclimatization [180], occurring during the summer months as a result of a combination of increased environmental temperature and physical activity (b) seasonal variations in physical activity levels [176], although some studies would argue against the second hypothesis, as seasonal variations are still present after adjustment for physical activity [176, 177].

Currently, there are no studies to address the effects of biological/seasonal variation of lipids in RA patients. The lack of studies is not surprising as several other factors associated with disease activity display similar fluctuations, such as morning joint stiffness, and disease activity worse in colder weather. Thus, it would be difficult to distinguish between the effects of the disease itself on lipid parameters and those occurring as a result of 'normal' variation.

Menopause

Female sex hormones have been shown to exert a protective effect against the development of CVD. A hormonal deficiency, such as that observed with the menopause, leads

to an increased risk of CVD [181]. This may be partially explained by accelerated proatherogenic changes in the lipid profile occurring independent of increasing age [182]. Postmenopausal women experience a deterioration in their lipid profile, with significant increases in LDL, TC, TG and HDL3 [182-184], with some studies reporting a worse lipid profile than age-matched male counterparts [185]. It has been suggested that genetic mechanisms that control individual variation of lipids, in particular HDL, may differ between premenopausal and postmenopausal women [186]. Relative correction of the hormonal profile through the use of hormone replacement therapy (HRT) can produce beneficial changes in the lipid profile by elevating HDL2 and reducing LDL levels [187]. However, despite these findings, we do not consistently see a decrease in vascular risk with HRT [188]. In addition, there is some evidence that CRP levels increase following the use of certain HRT products [189, 190].

In RA, postmenopausal status has been associated with significantly higher TC, TG, TC:HDL ratio, ApoB, Lp(a) and LDL:HDL ratio than those observed in premenopausal RA patients [8]. Postmenopausal RA women have also been shown to have higher TG and Lp(a) and a lower TC than healthy postmenopausal women [8].

Physical Activity

Regular long-term exercise promotes many health benefits, ranging from improved bone density to a reduction in cardiovascular risk. Much of the reduction in cardiovascular risk can be attributed to modification of conventional risk factors, including a reduction in blood pressure [191], improvement of the lipid profile [9, 42] and weight reduction.

In the general population, exercise is recommended as a lifestyle change for all patients with dyslipidaemia or estab-

lished cardiovascular disease [11]. Supporting this recommendation are several studies and meta-analyses demonstrating significant improvements in the lipid profile in patients adhering to a regular exercise programme [170, 191, 192]. The most significant changes observed with moderate exercise are elevations in HDL levels, lower TC:HDL ratio and lower TG levels [170, 191, 193, 194]. The scale of change appears to be influenced by the mode of exercise taken, with endurance exercise training proving to be substantially more effective than resistance training [195, 196]. Furthermore, age also seems to influence the degree of alteration in the lipid levels seen following exercise [197]. Older patients have been shown to lower the TC levels to a greater extent, but also appear to more readily enhance their HDL levels [197]. Unfortunately, the majority of exercise studies have been conducted in normolipidaemic subjects. Thus, there is a relative lack of available data on the effects of exercise training in dyslipidaemic patients.

Many of the improvements in the lipid profile require several months for the benefits to accrue. However, there is some evidence to suggest that changes in the lipid profile occur following a single episode of exercise [198]. Single exercise sessions appear to result in rapid increases in HDL levels occurring 18–24 h post-exercise that can persist for up to 72 h [199, 200]. Although the minimum amount of exercise required to improve blood lipids is not known, the United Kingdom government recommendations suggest at least 30 min of continuous exercise 5 days a week are required for a healthy lifestyle. However, continuous exercise is not required for alterations in the lipid levels to occur, as changes have been reported in patients whose exercise time is broken up throughout the day [201].

As HDL is one of the key components of the lipid profile to be modified by exercise, some interest has been generated around the effects of exercise on HDL sub fractions. A large proportion of studies assessing the influence of short term exercise on HDL sub fractions has shown significant increases in both HDL-2 (low density) and HDL-3 (high density) [202, 203]. The relative increases in the HDL sub fractions differ between studies with some demonstrating similar increases in both HDL-2 and HDL-3 [203], and others demonstrating a disproportionate increase in HDL-2, resulting in an increase in the HDL-2/HDL-3 ratio [204]. These discrepancies can be explained by differences in the subjects studied (athletes *vs* healthy men), the number of subjects involved, the mode and length of exercise, and advice regarding other lifestyle modifications. Evidence to support similar changes in HDL sub fractions among subjects adhering to longer-term exercise programmes, or in dyslipidaemic subjects is lacking.

The mechanism by which exercise mediates positive effects on HDL can be partly attributed to the alterations in the enzymes responsible for HDL metabolism [205]. Two of the key enzymes whose function may be altered are LPL and hepatic HL. One study demonstrated large increases in the activity of LPL and a significant reduction in HL activity in endurance trained individuals *vs* sedentary controls [206]. The enhanced enzyme activity of LPL may escalate lipid transfer to HDL, and simultaneous decreases in HL activity may result in reduced clearance of HDL.

In RA, traditionally patients and rheumatology healthcare professionals have been cautious regarding exercise, due to a widely held but unfounded fear that it would trigger aggravation of disease or damage the joints. These preconceptions resulted in exercise restriction and may, in part, account for the inactive lifestyle of many RA patients [207]. Indeed, physical activity has been shown to directly associate with CVD risk factors in patients with RA [208]. However, over recent years well designed physical exercise programmes have been shown not only to be safe in RA but also to promote prolonged improvements in morning stiffness, Stanford arthritis self-efficacy scale (SES) and other disease outcomes [209, 210]. The beneficial effects of exercise on disease parameters and well-being have been observed in all RA patients irrespective of disease duration [211] or activity [212, 213]. Although regular exercise as part of an individually designed programme is now recommended as an integral part of RA treatment, and has been included in the ACR treatment guidelines, there are no specific trials addressing the effects of exercise on CVD risk factors (e.g. dyslipidaemia) in RA. Specific well-designed studies are required to address this issue in RA, as the inflammatory burden in these patients may have already modified conventional cardiovascular risk factors (abnormal lipid profile, changes in body fat: muscle ratio etc), and thus the beneficial effects of exercise on CVD may be even greater than those seen in the general population.

Obesity

The classical pattern of dyslipidaemia in overweight and obese subjects is characterised by elevated levels of TG, TC, ApoB-100 and small LDL particles, and decreases in HDL levels [214, 215]. The dyslipidaemia of obesity has become one of the fundamental criteria for diagnosing the MetS [216–219]. Insulin resistance (IR) is often found in conjunction with both obesity and the MetS, and plays a key role in the development of dyslipidaemia. One of the primary obesity-associated defects in lipid metabolism is the overproduction of VLDL [220]. This phenomenon may be a consequence of a cascade of events occurring in the insulin resistant state, resulting in hepatic steatosis [221]. Furthermore, there is a delayed clearance of VLDL particles, due a reduction of the LDL receptor activity that occurs in conjunction with IR [143]. The low HDL levels seen in obese states are likely to be triggered by several mechanisms. Firstly HDL clearance is enhanced in insulin resistant states, due to stimulation of hepatic lipase and the resultant production of smaller HDL particles [222]. Secondly, transfer of apolipoproteins and phospholipids from TG-rich lipoproteins to HDL is reduced. Thirdly, the delayed clearance of TG-rich lipoproteins facilitates the CETP mediated exchange between cholesterol esters in HDL and triacylglycerols in VLDL [223].

RA associates with profound changes in body composition, partly as a result of the process of rheumatoid cachexia. This is an abnormal metabolic response leading to loss of lean body mass, which, in the case of RA patients, is often replaced by fat [224]. As a result, commonly used body mass index (BMI) thresholds for overweight and obesity in the general population, do not apply in RA [225]. Obesity in RA, has been linked to physical inactivity and worse RA

outcomes [46, 226], but also to a worse CVD risk factor profile [208]. Its exact association with lipid changes has not been studied in RA.

Smoking

Smoking reduces life expectancy, through multiple detrimental effects on health [227]. Much of the associated morbidity and mortality can be attributed to the carcinogens contained within the cigarette smoke, resulting in an increased risk of developing cancer (especially lung cancer) [228] or increased rates of CVD [229]. The mechanism by which smoking promotes CVD still needs to be elucidated. However, it has been suggested that smoking induced alterations in the lipid profile may contribute to this susceptibility [230]. Several large studies confirmed that smoking exposure associates with a pro-atherogenic lipid profile, with elevated TC, LDL, TG levels, and the TC:HDL ratio, and an associated reduction in HDL levels [231-234]. Data produced from such studies has to be interpreted with caution, as it is difficult to control for other lifestyle factors, such as weight changes, diet, alcohol consumption and physical activity. The relationship between smoking and lipid levels appears to be dose dependant [234] and readily reversible on cessation [235]. Cigarette smoke is known to create a pro-oxidative state in the circulation [236], and disturb endothelial function [237]. The smoking-induced oxidative burden induces lipid peroxidation [238], thus further accelerating atherosclerosis.

Smoking has been linked to a 2-4 fold increased risk of developing RA [239], and may also influence the severity of the disease [240], functional status [241] and other parameters such as body weight and muscle mass [242]. However, a recent study has produced conflicting findings suggesting that cigarette smoke does not accelerate RA disease progression. In fact, the study demonstrated that heavy smoking may be associated with reduced radiographic progression and improvement in functional scores [243]. The effect of smoking in RA varies due to the length of exposure, with acute exposure resulting in immuno-stimulatory effects and chronic exposure resulting in immunosuppressive effects [244]. Although, it is clear that smoking may influence the development and pathogenesis of RA, no specific studies have addressed the effects of smoking on lipid metabolism in RA.

Alcohol

Moderate alcohol intake associates with a lower risk of CVD [245-247]. The reduction in CVD risk of moderate drinkers compared with those who abstain has been shown to lie between 10 and 40% [247]. The cardio-protective effect of alcohol has been linked to beneficial changes in both the lipid profile [247-249] and haemostatic parameters [250, 251]. By far the most common and well-reported change in the lipid profile is an increase in HDL concentrations [234, 247, 252, 253]. The precise mechanisms contributing to the quantitative change in HDL are still being scrutinised, however, current proposals include: (a) increased hepatic production or increased transport rate of apoA-I and apoA-II [254, 255] (b) increased cholesterol efflux from macrophages to HDL mediated by ABCA1 [256] (c) alcohol-induced re-

duced activity of CETP [257]. However, a recent study failed to demonstrate any relationship between CETP activity and HDL levels among moderate alcohol consumers compared with abstinent controls [258]. Furthermore, the study demonstrated alcohol-induced alterations in LPL activity.

A degree of controversy exists regarding the relative contributions of HDL subfractions, to the elevation of HDL levels seen with moderate alcohol intake. Virtually all studies demonstrate an increase in both HDL-2 and HDL-3 subfractions [253, 259]. However, the effects on the HDL-2:HDL-3 ratio varies between studies [259, 260]. One study assessing the effects of habitual alcohol consumption has demonstrated that the changes in HDL subfractions may differ between genders, with males experiencing a rise in both HDL-2 and HDL-3, but females only experiencing an isolated significant rise in HDL-2 [253]. A recent study, evaluated the qualitative changes of HDL observed during exposure to varying degrees of alcohol in 279 healthy men [259]. The investigators confirmed that alcohol consumption results in a shift from HDL-3 to predominantly HDL-2a, and a significant phospholipid enrichment of all HDL subfractions.

Hypertriglyceridaemia has long been associated with regular alcohol use [261]. A meta-analysis of 42 studies has confirmed a positive relationship between moderate alcohol consumption and TG levels, reporting a 0.19 mg/dl increase per g of alcohol consumed per day [247]. The impact of alcohol exposure on the other components of the lipid profile including LDL and TC levels have been addressed in comparatively fewer studies. However, the majority of these demonstrate a modest inverse correlation of both TC and LDL with alcohol [248, 262]. LDL particle size may be reduced in patients with alcohol-induced hypertriglyceridaemia, resulting in a subsequent increased susceptibility to oxidation [263].

Potential predisposing factors in the development of RA have been debated for decades [264]. The onset of disease is thought to be the result of a complex interaction of environmental agents with genetic factors [265]. Epidemiological studies indicate that alcohol consumption correlates with a reduced risk of developing RA [266, 267]. Unfortunately, post diagnosis data to assess the effects of alcohol on disease parameters and the lipid profile are lacking. Studies to address this issue are likely to be hampered by ethical and safety issues regarding interactions with disease modifying anti-rheumatic drug (DMARD) therapy, in particular, patients are actively discouraged from drinking alcohol when they are receiving therapy with methotrexate, which is currently by far the most commonly used DMARD for the treatment of RA.

Diet

It is well established that dietary intake has far reaching health implications. An unhealthy diet rich in saturated fats and sugars has been linked to an increased risk of CVD [268], whereas diets high in unsaturated fats and antioxidants such as the Mediterranean diet are linked to a reduced CVD risk [269]. There are multiple individual dietary components that impact upon the lipid profile. Fish oils are rich in omega-3 polyunsaturated fatty acids which lowers plasma TG levels [270, 271], red yeast rice has been shown

to reduce TC and LDL levels [272] and olive oil raises HDL-cholesterol and reduces levels of oxLDL [273]. In RA, data regarding the impact of diet on the lipid profile is sparse. One randomised study on 66 active RA patients, demonstrated that patients treated with a gluten-free vegan diet had lower LDL and oxLDL levels [274].

THE EFFECTS OF DRUGS ON THE LIPID PROFILE

The impact of recent therapeutic advances in the management in RA on cardiovascular risk is not clear. However, several recent studies suggest that therapeutic intervention and control of disease activity may reduce cardiovascular risk [42, 43, 275]. A large cross sectional study (QUEST) of over 4,000 RA patients, suggested a reduced cardiovascular risk with prolonged use of DMARDs, glucocorticoids (GCs) or anti-tumour necrosis factor therapy (anti-TNF) [275]. These findings should be interpreted with caution as the study is limited by its cross sectional design and therefore causality can not be assumed. Despite this, a further prospective study has confirmed significant improvements in cardiovascular risk factors following treatment with methotrexate and GCs [42]. Part of these observed effects may be governed through drug-induced changes in the lipid profile. In this section we will summarise the effects of DMARDs, GCs, and biologic agents (anti-TNF, rituximab and tocilizumab) on the lipid profile in RA.

The use of statins in RA is escalating at an exponential rate due to the increased awareness of the importance of tackling CVD in RA. Statins classically produce a less atherogenic lipid profile by lowering TC, LDL and TG and increasing HDL levels. However, in RA statins have been shown to exert additional anti-inflammatory properties [276], resulting in suppression of disease activity [277]. Thus, it is possible that statins produce their effects on the lipid profile in RA both through the inhibition of 3-hydroxy-3-methylglutaryl coenzyme A and through suppression of the inflammatory load.

The pleiotropic effects of statins are far reaching. Statins have anti-inflammatory properties, which have been shown to have a modest beneficial impact on the clinical signs and symptoms of RA [277, 278] and other forms of inflammatory arthritis [279, 280]. Statin use has been reported to produce a statistically (though not clinically) significant reduction in the number of swollen joints and disease activity scores (DAS) [277, 281]. In addition, suppression of biochemical markers of inflammation have been observed [277, 282], with statin therapy suppressing CRP and ESR levels by as much as 50% and 28%, respectively [277]. This suggests that although statins are unlikely to produce clinically significant anti-rheumatic effects on their own, they may be useful adjuncts to standard anti-rheumatic drug therapy. Also, although they are likely to act primarily act on the lipid profile to reduce CVD risk, in RA they have been shown to reduce other markers of vascular dysfunction, including arterial stiffness [283], and flow mediated dilatation [284]: it remains unclear whether this is mediated through lipid effects, anti-inflammatory effects, or is due to other direct effects on the vasculature. There are however no hard CVD end-point statin trials in RA. In fact, RA patients have been systematically excluded from all large statin trials in the gen-

eral population. Such trials are required in patients with RA due to the multitude of factors that may contribute to vascular damage in this disease, which make it difficult to predict whether the effect of statins will be of the same magnitude as that seen in the general population.

Statins may also improve manifestations of extra-articular disease in RA patients, such as renal dysfunction. Renal dysfunction is very common in RA with around two thirds of all RA patients reported to have a mild renal impairment and a further 10-15% to have moderate renal impairment [285]. The renal dysfunction seen in RA patients may occur as the result of inflammation-mediated vascular damage, drug therapy (e.g. non-steroidal anti-inflammatory drugs), and the presence of other associated comorbidities e.g. glucose intolerance and hypertension. Interestingly, poor renal function has been found to associate with an excess cardiovascular mortality in the general population [286], with end stage kidney disease patients approximately twenty times more likely to die from CVD [287]. Although no studies have directly assessed the effects of statins on renal function in RA, data from studies performed in the general population suggest that statins may slow renal function decline [288, 289]. The diversity of the potential beneficial effects of statins, makes them an attractive choice for both CVD risk reduction and for symptom control in RA [290].

The precise effects of statins on dyslipidaemia, disease activity and severity, as well as hard CVD end-points in RA are currently being investigated in the largest, prospective, double-blind, randomised controlled trial ever conducted in RA (the TRACE RA trial) [291].

Glucocorticoids (GCs)

GCs have long been recognised to have beneficial effects in RA [292, 293]. However, in current rheumatological practice GC use has been curbed due to adverse effects. They are used increasingly as a short term measure to induce a rapid reduction in disease activity while awaiting the effects of slower acting DMARD therapy [294]. The long term use of GCs is controversial due to loss of efficacy over time, and the undesirable side effect profile [293].

For many years it has been widely assumed that GC use associates with a more atherogenic lipid profile (raised TC, TG and HDL levels) regardless of the indication for use [295]. However, many of these early studies were performed on renal transplant patients. Thus, they may have been confounded by other lipid influencing variables, such as the underlying disease process, co-morbid conditions and concomitant medications [296, 297]. Over more recent years, conflicting data has been produced from several prospective studies, indicating that GC may actually improve the lipid profile by increasing HDL and lowering the TC:HDL ratio [298, 299]. In 2005, a large study involving 15,004 participants demonstrated differences in the lipid profile in patients receiving GCs according to their age [300, 301]. In patients over the age of 60, GC use was associated with higher HDL and ApoA-I levels, and lower TC:HDL and ApoB:ApoA-I ratios. No association was noted between GC use and HDL levels, TC levels or TC:HDL ratio among subjects below the age of 60. A potential limitation of this study was the difference in the indication for GC use between the age groups,

and thus the underlying effects of the individual disease processes on the lipid profile. Unsurprisingly, studies assessing the effects of GCs on lipid metabolism in healthy individuals are sparse. A small study on 8 normolipidaemic healthy men treated with prednisolone for 14 days demonstrated a significant increase of VLDL-TG, VLDL-cholesterol, HDL-cholesterol, Apo A-I and ApoE, with normalisation of the lipid profile to baseline levels 2 weeks after discontinuation of prednisolone [7, 301]. A further study on 8 normolipidaemic healthy men examined the relationship between GC exposure and HDL sub fractions [302]. This demonstrated a rapid change in the lipid profile following initiation of 30 mg oral prednisolone, with elevations seen in HDL within 2 days of commencement. Within the HDL subclass, GC exposure induced redistribution of HDL particles, resulting in increased levels of HDL2 and a reduction of HDL3.

A systematic review of the literature was performed to identify if GC use alters the lipid profile in RA. The search criteria included 'rheumatoid arthritis', 'lipid', 'lipids', 'dyslipidaemia', 'steroids', 'prednisolone' and 'glucocorticoids'. Following restriction of the search criteria (age >19, human, English, title/abstract), a total of 10 papers were identified. Of these, only 6 papers mentioned the effects of GCs on the lipid profile in RA. However, the majority did not set out to assess this relationship as their primary objective. Thus, the number of patients included that were actually receiving GCs in the majority of studies was very small. The results of some of the studies may have also been affected by potential confounders, such as concurrent changes in DMARD therapy [303]. The largest and most robust study to comment on the effect of GCs on the lipid profile was reported by Boers *et al.* [304]. This was a randomised controlled trial in which 76 RA patients were randomised to take combination therapy of methotrexate (stopped at 40 weeks), sulphasalazine and a reducing dose of prednisolone (initially 60 mg/day prednisolone, tapered in 6 weekly steps to 7.5 mg/day and stopped after 28 weeks) and 79 RA patients who were allocated to take sulphasalazine alone. Both arms of the study were well matched according to age, disease duration and disease activity. The study reported a significant increase in TC and HDL levels, and a reduction in the TC:HDL ratio in both study arms. However, the greatest improvements were observed in the combination therapy arm, with the corrective effect on the lipid profile occurring far more rapidly. The lipid changes were far more significant in the combination therapy arm up until 28 weeks and then all lipid parameters returned to levels similar to those seen in the sulphasalazine only arm, indicating that the changes in the lipid profile may be more specific to GC use (stopped at 28 weeks) rather than the additive effects of the 3 combination therapy drugs.

In RA, the limited available data addressing the relationship between GC exposure and the lipid profile, demonstrate a corrective effect on the altered lipid profile seen in active RA. The most widely reported changes include an increase in both TC and HDL levels [299, 304]. However, HDL appears to increase at a proportionately higher rate compared with TC, thus generating a more favourable atherogenic index [15]. Much of the witnessed normalisation of TC and HDL levels with GC use in RA has been attributed to the

suppression of disease activity through their anti-inflammatory actions [15, 304]. Data produced regarding changes in TG levels is not so clear and has only been specifically looked at in 3 out of the 6 studies identified. Although, 2 studies have shown an increase in TG levels with GC use [5, 305], another study failed to demonstrate any change [303]. This discrepancy may be attributed to differences in the populations studied including disease characteristics, drug use, and the power of the individual studies. Further large studies are required in order to fully elucidate the effects of GCs on TG levels. However, as part of a recent cross sectional study examining the effects of GCs on the presence of the MetS we have demonstrated that although TG and HDL levels increase with GC use, this does not associate with an increased prevalence of the MetS [53].

All studies demonstrating the effects of glucocorticoids on the lipid profile in patients with RA are summarised in Table 2.

Hydroxychloroquine

Hydroxychloroquine (HCQ) is an anti-malarial medication that also exhibits disease modifying and anti-inflammatory actions when prescribed for RA or SLE. In the 1980's Beynen noted a reduction in cholesterol synthesis with low dose chloroquine [307]. A subsequent study confirmed these initial observations and demonstrated reductions in cholesterol and TG levels in those prescribed chloroquine [5]. In the 1990's Wallace *et al.* carried out a study to assess whether HCQ exhibited similar lipid lowering properties as its parent drug, chloroquine [305]. The study involved 150 patients with RA or SLE who were randomised to take HCQ alone, GCs alone, HCQ and GCs, or no drug. HCQ use resulted in a reduction in cholesterol, LDL and serum TG levels, which appeared to be independent of changes in weight and diet. Later studies demonstrate that HCQ produces a less atherogenic lipid profile, by increasing HDL levels by approximately 15% [308].

The mechanism underlying the lipid lowering effects of HCQ and chloroquine are still disputed. However, some plausible modes of action have been described in studies analysing the interaction of chloroquine. Potential reasons include: (a) *via* the inhibitory actions on VLDL secretion by the liver [309] (b) inhibition of cholesterol synthesis by blockage of a site distal to hydroxymethylglutaryl coenzyme A (HMGCoA) reductase [310], and, (c) inhibition of proteolysis of internalised cholesterol esters leading to increased LDL receptor values [307, 311].

Ciclosporin

Since the first reported use of ciclosporin A (CyA) for the management of RA in 1979 [312], the drug has been used both as monotherapy or in combination with other DMARD therapy, such as methotrexate [313, 314]. Chronic CyA administration has been shown to adversely affect the lipid profile. However, much of this data has been generated from small studies on transplant patients [315, 316]. A study of 65 post renal transplant patients, demonstrated elevations in TG and Lp(a) levels, and suppressed HDL levels in those treated with CyA monotherapy compared with azathioprine and prednisolone [317]. The reduction in HDL levels may be

Table 2. Studies Assessing the Impact of Glucocorticoid Use on the Lipid Profile in Patients with RA

| Study | Number of Patients | Other Drugs Used | Follow Up | Patient Demographics | Lipid Changes |
|---|--|--|---------------------|--|--|
| Peters <i>et al.</i> Ann rheum dis 2007 [15] | 80 RA (35 on pred) | Infliximab | 48 wks | Age 56 DD: 10 yrs Female 77 % | Pred ↑ TC and HDL levels and ↓ TC:HDL ratio |
| Dessein <i>et al.</i> J.Rheumatol 2004 [303] | 92 RA (37 had pred in past) (18 currently on pred ≤ 4mg) No controls | At enrolment 17 patients taking DMARDs by completion 84 patients on DMARDs | 6 mths | Female 80% Age 56 yrs DD: 11 yrs | No changes in LDL, HDL or TGs |
| Boers <i>et al.</i> Ann rheum dis 2003 [304] | 155 early active RA | Patients randomised to either MTX & SLZ & pred (76) or SLZ alone (79) | 56 wks | Age 50 yrs DD: 4 months Female 59% | ↑ TC, HDL and ↓ TC:HDL ratio in both group but greater in com- bination therapy group. |
| Dessein <i>et al.</i> Arthritis res 2002 [306] | 79 RA (10 on pred) 39 age/sex matched OA controls (6 on pred) | Median dose of pred 5 mg | Cross- sectional | Age 52 yrs Female 83% DD: 8.5 yrs | No significant differences between RA patients tak- ing and not taking steroids. |
| Wallace <i>et al.</i> Am J med 1990 [305] | 108 SLE, 47 RA (14 HCQ, 8 pred, 4 HCQ & pred, 21 neither drug) 108 SLE | HCQ | Cross- sectional | Age 46.6 yrs | Pred alone have ↑ TC, HDL and TG but ↓ LDL compared to those not on treatment. (unable to dis- tinguish between effects in RA and SLE patients) |
| Svenson <i>et al.</i> Arch intern med 1987 [5] | 33 (only 4 treated with just pred) | Pred & AZA or Pred & cyclo or Pred alone | 9 mths | Age 49 yrs | ↑ TC, TG, HDL and LDL in all groups including pred only arm. |

TC: total cholesterol, TG: triglycerides, HD: high-density lipoproteins, LDL: low-density lipoproteins, Pred: prednisolone, MTX: methotrexate, SLZ: sulphasalazine, AZA: azathioprine, HCQ: hydroxychloroquine, cyclo: cyclophosphamide, OA: osteoarthritis, SLE: systemic lupus erythematosus, DD: disease duration, RA: rheumatoid arthritis, DMARDs: disease modifying anti-rheumatic drugs, yrs: years, wks: weeks, mths: months.

explained by the inhibitory actions of CyA on ABCA1-mediated lipid efflux [318]. Despite these findings in transplant patients, further studies are required to address the potential effects of CyA on the already altered lipid profile seen in RA.

Gold

Gold is one of the oldest treatments for RA, and was first used in the late 1920s [319]. Its use has diminished over recent years with the advent of newer more effective DMARDs and biologic therapies. Gold therapy has many adverse effects including dermatitis, stomatitis, post injection reactions, haematuria and proteinuria. However, not much has been reported about its effects on the lipid profile. Munro *et al.* reported that gold use may have the net effect of producing a more atherogenic profile by increasing TG levels and reducing HDL levels [308]. No further studies have directly looked at the influence of gold on lipid levels.

Anti-Tumour Necrosis Factor Agents

The identification of TNF- α as a key cytokine in the pathogenesis of RA has resulted in the development of specific biologic therapies designed to target TNF- α with the

net effect of inhibiting its inflammatory properties [320]. The introduction of anti-TNF agents in 2000, has revolutionised the treatment of RA with better disease control and dramatic improvements in quality of life [321]. Despite the overwhelming benefits of anti-TNF therapy, a number of complications and adverse effects have been noted [322]. By far the most common complication of anti-TNF therapy is infection [323]. However, there is expanding evidence to suggest that these agents may also interact with other metabolic parameters, such as the lipid profile [73].

To date, 3 anti-TNF agents have been licensed for use in RA, Infliximab, Adalimumab and Etanercept. The structure and mode of action of the 3 agents varies. Infliximab is a chimeric monoclonal antibody, adalimumab is a fully humanised monoclonal antibody and etanercept is a soluble TNF- α receptor fusion protein. The underlying characteristics of these molecules may be key to their mode of action and effects on the lipid profile.

A systematic review of the literature was carried out to address the effect of the 3 anti-TNF agents on the lipid profile. A PubMed search was conducted with specified search criteria: 'rheumatoid arthritis', 'lipid', 'lipids', 'dyslipidaemia', 'anti-tumour necrosis factor', 'infliximab', 'etanercept'

and 'adalimumab. The limites applied to the search included: age >19, human, English and title/abstract. The search identified a total of 19 papers, of these 16 studies were found to address the impact of anti-TNF agents on the lipid profile. Overall, the studies recruited relatively small populations and had a short duration of follow up, with several studies only looking at a handful of patients over a 6 week period [324-326]. The largest study enrolled 97 RA patients who were prospectively followed up for 1 year [327]. The study was limited by the lack of a control group and only reported changes in HDL and TC. Twelve of the fifteen studies did not include a control group for comparison.

Multiple studies have produced information on the short-term effects of anti-TNF agents on the lipid profile [328-330], but there are relatively few addressing the longer term effects [15, 121]. In the short term studies, duration of treatment ranged from 6 weeks to 6 months, and the majority only assessed the effects of infliximab [122, 328, 329, 331]. Irrespective of their duration, these studies demonstrate similar findings with a universal increase in TC, and a large proportion confirming an increase in HDL, but no overall change in the atherogenic index (TC:HDL ratio). Although TG levels were not looked at in all studies, the available data tends to show an increase in TG levels up until 6 months in patients treated with infliximab [122, 328]. Saiki *et al.*, have shown the most convincing evidence of a relationship between anti-TNF and an increase in TG levels [332]. The study compared TC and TG levels among 32 patients with refractory RA treated with infliximab, to 32 age- and sex-matched control patients with active RA treated with methotrexate over a 6 month period. A significant and persistent elevation in TG levels was observed from 2 weeks in the infliximab arm but no change was seen among those treated with methotrexate, thus indicating that changes in TG levels are likely to be due a drug-specific mechanism rather than a 'blanket' anti-inflammatory effect.

Published data regarding the effects of the other 2 anti-TNF drugs is limited. The short term effects of etanercept and adalimumab have only been studied in 2 studies [4, 330]. However, due to study design and small number of patients enrolled, they only reported on the generic effect of anti-TNF on the lipid profile rather than the effect of the individual drugs. Seriola *et al.*, reported similar findings to the studies performed solely on patients receiving infliximab (increased TC and HDL) [4], whereas the other study reported no effect of anti-TNF on the lipid profile [330]. The effect of etanercept and adalimumab on the lipid profile needs to be addressed in further large scale studies, particularly as differences in their molecular structure, mode of action and half-life may alter how they affect the lipid profile.

The effects of anti-TNF therapy on Lp (a) levels were addressed in only 3 studies [4, 324, 330]. None of the studies demonstrated a significant change in Lp(a) levels. However, this may be the result of a lack of power due to small study size, with the largest recruiting a population of 34 RA patients [4]. Unfortunately, 2 of these studies attempted to look at the effects of all 3 anti-TNF agents, thus making the results difficult to interpret, as Lp(a) levels may be affected differently by each TNF agent e.g. cancelling each other out in the reported results.

A summary of all studies addressing the effects of anti-TNF agents on the lipid profile in patients with RA is shown in Table 3.

Rituximab

Rituximab, an anti-CD20 monoclonal antibody, is used to treat patients with active RA unresponsive to DMARDs and/or 1 anti-TNF agent. Rituximab was originally developed for the treatment of B cell lymphoma, but has since been found to be useful in other conditions including RA, with significant reductions in disease activity and functional improvement [333, 334]. Although rituximab is now widely used, relatively little is known about the effects of the drug on the lipid profile, structure or function. To date, there has been 1 prospective study of 6 RA patients treated with rituximab for 6 months [335]. The investigators report minor lipid changes at 2 weeks. However, the changes in the lipid profile are most likely to have been induced by the concomitant infusions of methylprednisolone (given to reduce side effects) rather than the rituximab per se [336]. The lipid profile returned to baseline and no significant differences were reported after 6 months of treatment. However, this study had a primary objective of assessing endothelial function and not lipid parameters and therefore it was unclear whether potential confounders such as statin use were taken into consideration.

Tocilizumab

Tocilizumab is an interleukin-6 (IL-6) receptor antagonist, offering a different mode of action to all other currently available biologic therapies. Although data from clinical trials have shown tocilizumab to be effective at controlling disease activity in RA [125, 337], concerns have arisen regarding its effects on the lipid profile. Available evidence demonstrates a significant increase in TC, HDL, TG and LDL levels following treatment with tocilizumab [125, 337, 338]. The relative changes in each of these levels and the impact this has on atherogenic indices such as TC:HDL ratio is less clear, with some studies reporting an increase in the TC:HDL ratio [125], while others failed to demonstrate a change [337]. It has been hypothesized that the elevations in lipid levels occurs as part of a 'corrective' phenomenon of treatment by reversing the effects of high-grade inflammation on the lipid profile. Although studies do demonstrate both suppression of inflammatory markers (ESR and CRP) and elevations in lipid levels, no direct correlations have been reported to confirm an association. Further studies are therefore required to not only address this issue, but also to establish the effects of tocilizumab treatment on other lipid parameters such as apolipoproteins, Lp(a), lipoprotein structure and function, which may also confer CVD risk.

GENETIC REGULATION OF LIPID METABOLISM

ATP Binding Cassette Transporter Gene (ABCA1) Polymorphisms

The ATP binding cassette transporter protein is primarily involved in the transport of phospholipids and cholesterol from cells to intracellular acceptors, including lipid-free ApoA1. Polymorphisms of the ABCA1 gene lead to alterations in protein function. In the most severe form, patients

Table 3. The Effects of Anti- Tumour Necrosis Factor Agents on the Lipid Profile in Patients with Rheumatoid Arthritis

| Study | Type of Anti-TNF | Number of Patients | F/U | Changes in Lipid Profile | | | | |
|---|---|--|------------------------------------|--------------------------------------|--------------------------------------|--|-----------------|----------|
| | | | | TC | HDL | TC/HDL | TG | Lp(a) |
| Garces <i>et al.</i> , Ann.Rheum.Dis 2008 | Inflix Etan | 30 RA, 29 AS, 6 PsA Inflix= 44 Etan= 21 | 1yr | ↑ inflix N/C Etan | N/C inflix ↑ Etan | NS NS | NS NS | NS NS |
| Nishida <i>et al.</i> Endocrine J 2008 [327] | Inflix | 97 | 1 yr | ↑ | ↑ | NS | NS | NS |
| Soubrier <i>et al.</i> Joint bone spine 2008 [330] | All | Etan =6 Inflix =11 Adal = 12 | 14 wks | N/C | N/C | N/C | N/C | N/C |
| Oguz <i>et al.</i> Acta Clin Belg 2007 [326] | Inflix | 7 | 9 mths | N/C | N/C | NS | NS | NS |
| Popa <i>et al.</i> Ann Rheum Dis 2007 [73] | Inflix | 55 | 55=6 mths 31=1 yr | ↑ | ↑ | ↑ | NS | NS |
| Peters <i>et al.</i> Ann Rheum Dis 2007 [15] | Inflix | 80 | 48 wks | ↑ at 6 wks, by 48 wks baseline | ↑ at 6 wks, by 48 wks baseline | ↑ at 6 wks, by 48 wks back to baseline | NS | NS |
| Saiki <i>et al.</i> J. Rheumatol 2007 [332] | Inflix | 43 refractory RA 32 age/sex matched RA controls on MTX 32 healthy controls | 6 mths | ↑ from 2 wks | NS | NS | ↑ from 2 wks | NS |
| Tam <i>et al.</i> Clin Rheumatol 2007 [328] | Inflix | 19 | 14 wks | ↑ | ↑ | N/C | ↑ | NS |
| Allanore <i>et al.</i> Clin Chim Acta 2006 [119] | Inflix | 59 refractory RA 56 RA controls not on anti-TNF | 30 wks | ↑ | ↑ | N/C | ↑ | NS |
| Dahlqvist <i>et al.</i> Scand J Rheu- matol 2006 [121] | Inflix (41 also MTx, 13 other DMARD 28 Pred) | 52 RA on infliximab 70 early RA controls | 2 yrs | ↑ initial, then ↓ by 6 month | ↑ initial, then ↓ by 6 month | ↑ | NS | NS |
| Kiortis <i>et al.</i> J Rheumatol 2006 [122] | Inflix | 82 (50 = RA 32 = AS) | 6 mths | ↑ | N/C | N/C | ↑ | NS |
| Seriolo <i>et al.</i> Ann NY Acad Sci 2006 [4] | All (plus MTX & pred) | Etan = 16 Inflix = 14 Adal =4 | 48 wks | ↑ | ↑ | N/C | N/C | N/C |
| Spanakis <i>et al.</i> J Rheumatol 2006 [331] | Inflix | 60 (24=RA 26=AS 10=PsA) | 6 mths | ↑ (1 st month only | ↑ | ↓ (at 3 months) | NS | NS |
| Vis <i>et al.</i> J Rheumatol 2005 [329] | Inflix | 69 | 6 wks | ↑ | ↑ | N/C | NS | NS |

(Table 3) contd....

| Study | Type of Anti-TNF | Number of Patients | F/U | Changes in Lipid Profile | | | | |
|--|------------------|--------------------|-------|--|-----|--------|-----|-------|
| | | | | TC | HDL | TC/HDL | TG | Lp(a) |
| Irace <i>et al.</i> Atherosclero 2004 [325] | Inflix | 10 | 6 wks | ↓ only after 1 st infusion | ↓ | NS | N/C | NS |
| Cauza <i>et al.</i> Wien Klin Wo- chenschr 2002 [324] | Inflix | 7 | 6 wks | N/C | ↓ | NS | ↑ | N/C |

Anti-TNF: anti- tumour necrosis factor, Inflix: infliximab, Etan: etanercept, Adal: adalimumab, MTX: methotrexate, Pred: prednisolone, DMARD: disease modifying anti-rheumatic drugs, RA: rheumatoid arthritis, AS: ankylosing spondylitis, PsA: psoriatic arthritis, wks: weeks, mths: months, N/C: no change, NS: not significant, yrs: years, wks: weeks, mths: months.

develop Tangier disease, characterised by reduced HDL in heterozygotes or clinically absent HDL in homozygotes [339]. To date, a number of single nucleotide polymorphisms (SNPs) have been identified and their association with plasma lipid levels and cardiovascular risk addressed in general population studies [340-342]. Patients carrying the K allele of the rs2230806 (R219K) SNP appear to be significantly protected from developing CVD, and a variety of studies report that it associates with lower TG levels and higher HDL levels [343-345]. Other polymorphisms including the I883M (rs4149313) and V771M (rs2066718) have been associated with an increased CVD risk [341, 346, 347], but demonstrate inconsistent effects on lipid parameters [348, 349]. R1587K (rs2230808) and E1172D (*rs33918808*) have also both been associated with an increased CVD risk [341, 350]. ABCA1 polymorphisms and their association with CVD have not been studied in RA.

Cholesterol Ester Transfer Protein (CETP) TaqIB Polymorphism

The cholesterol ester transfer protein (CETP) is key to the transfer of cholesterol esters and triglycerides between lipoproteins. It aids reverse cholesterol transport and HDL metabolism, by transferring cholesterol esters from HDL to VLDL and LDL. Variations in CETP levels can occur as a result of a SNP in intron 1 of the CETP gene (TaqIB), located on chromosome 16. The presence of a B2 allele confers lower CETP levels than the presence of the more common B1 allele. Thus, patients homozygous for the B2 allele often have significant elevations in their HDL-C levels compared with B1 homozygous patients [351]. Based on these findings, it is rather unsurprising that the B2 allele has also been associated with a lower CVD risk in the general population [352, 353]. A meta-analysis of 113,833 patients has demonstrated further changes associated with inheriting the B2 allele, including lower CETP mass and activity, lower levels of LDL-C, apoB, and TGs and increased levels of ApoAI [354]. Interestingly, the presence of the B2 allele may also enhance clinical benefit from statin therapy among patients with significant coronary artery disease, with a net effect of reducing cardiovascular events [355]. The worldwide frequency of the minor allele (B2 allele) is 42%. This was reported to be identical among East Asian and white populations [354]. The genotypic and allelic frequencies of

the CETP TaqIB and the influence this may have on the lipid profile or CVD risk has not been studied in RA.

Apolipoprotein E (ApoE) Polymorphisms

ApoE is a major component of VLDL and chylomicrons. It is essential for the catabolism of TG-rich lipoprotein constituents, and facilitates the uptake of VLDL and chylomicrons into the liver [356]. ApoE may further regulate lipid metabolism *via* enhancing effects of LPL and HL [357]. Three isoforms of ApoE exist as a result of 2 SNPs, ApoE2, ApoE3 and ApoE4. ApoE3 is considered as the parent form, occurring in more than 60% of the population [358] and the other 2 as mutations. ApoE2 is produced due to a base change (arginine to cysteine) at residue 158, whereas ApoE4 is the consequence of a base change (cysteine to arginine) at residue 112. Such changes alter the structure and function of the protein, resulting in a significantly reduced binding affinity of ApoE2 to the liver receptors [359], a subsequent delay in the removal of dietary fat from the blood [360] and the development of type III hyperlipidaemia [361]. ApoE4 predisposes to atherosclerosis through mechanisms that are still being elucidated [358]. However, some studies indicate that the lipoprotein distribution of ApoE4 differs from the parent form, and that particles containing ApoE4 are prone to catabolism, which ultimately leads to further alterations in lipoprotein metabolism with the end result of increasing LDL, TC and TG levels [362, 363]. ApoE polymorphisms may also partially account for variations observed in atherosclerotic plaques, with differences being observed in carotid intima-media thickness [364]. The prevalence and contribution of the ApoE genotypes to the development of CVD in RA have not yet been studied.

Lipoprotein Lipase (LPL) Polymorphisms

The LPL gene is located on chromosome 8p22 and is responsible for the production of LPL. The enzyme plays a key role in the mobilisation of non-esterified fatty acids and monoglycerol molecules for energy utilisation and storage by cells *via* the hydrolysis of the triacylglycerol component of VLDL and chylomicrons. Thus, a deficiency of this enzyme can result in an accumulation of both plasma chylomicrons and TGs, as seen in the autosomal recessive condition – familial LPL deficiency [365]. To date several SNPs of the

LPL gene have been associated with lipid abnormalities and CVD [366]. Four SNPs including D9N (rs1801177), N291S (rs268), S447X (rs328) and hindIII (rs320) have been shown to significantly alter plasma levels of TG and HDL [366]. Available data, from several independent studies have demonstrated an increased CVD risk with LPL mutations (S447X, *PvuII* (rs285)) [367-369], and this association has recently been confirmed in a large meta-analysis [366]. The HuGE association review and meta-analysis demonstrated an increased odds ratio (OR) for myocardial infarction and coronary stenosis with G188E (OR=2.8), D9N (OR=1.33) and T-93G (OR=1.22). However, the remaining 4 SNPs (N291S, *PvuII*, HindIII, S447X) analysed were not found to associate [366]. The frequency and impact of such genetic polymorphisms have not been studied in RA.

Apolipoprotein (A1/C3/A4/A5) Gene Cluster

The long arm of chromosome 11 is home to the regulatory gene cluster that encodes for proteins ApoA1, ApoC3, ApoA4 and ApoA5. Interestingly, not only have the genes responsible for these proteins been found to be close in their chromosomal proximity, but the proteins produced by each gene have been found to have inter related functions involved in the metabolism of TG-rich lipoproteins and HDL [370, 371], a phenomenon that may be attributed to linkage disequilibrium. ApoA1 is the main protein component of HDL and plays an important regulatory role in reverse cholesterol transport. ApoC3 is found in association with VLDL and HDL, and primarily acts as an inhibitor of LPL [372], thus slowing the rate of catabolism of triglyceride rich particles. Due to the inhibitory effects of this protein an inverse relationship between ApoC3 levels and TG levels exists. ApoA4 is a major constituent of chylomicrons, and is thought to be involved in TG metabolism [373], along with ApoA5 [374-376]. An extensive array of SNPs has been identified within the genes responsible for the production of each of these proteins, which have been linked not only to alterations in lipid metabolism and but also to CVD. For example, around 20 SNPs of the ApoA5 gene have been identified, with around half of these found to be associated with disorders of TG metabolism [377], and others associated with excess CVD risk [374, 378]. ApoC3 M455, ApoA4 T347S, ApoA5 S19W may be particularly interesting in the context of the lipid abnormalities described in RA, but have not been studied yet in this condition. ApoC3 M455 is associated with HDL levels, with the presence of the C allele shown to significantly reduce levels [379]. The Apo A4 S347T is associated with an increased risk of CVD [380], and apo A5 S19W with significantly elevated TG levels [381, 382].

CONCLUSIONS

Dyslipidaemia in RA is the consequence of the complex interaction of multiple factors, with inflammation fundamental to most. Although our overall understanding of the role of inflammation in dyslipidaemia can help us to understand the lipid changes seen in RA, further studies are required specifically in RA as there may be disease specific phenomena occurring. The contribution of genetic factors to dyslipidaemia in RA requires further investigation in large-scale studies, particularly in the context of gene-environment interac-

tions. In addition, the precise contribution of dyslipidaemia to the excess CVD risk seen in RA remains to be elucidated, providing scope for future research.

ACKNOWLEDGEMENTS

This work is supported by an Arthritis Research Campaign Clinical Fellowship grant (grant number 18848 to T.E.T), and an Arthritis research Campaign infrastructure support grant (grant number 17682, given to the Dudley Group of Hospitals NHS Foundation Trust, Department of Rheumatology).

DECLARATION OF INTEREST

The authors declare they have no conflicts of interests.

REFERENCES

- Gazi IF, Boumpas DT, Mikhailidis DP, Ganotakis ES. Clustering of cardiovascular risk factors in rheumatoid arthritis: the rationale for using statins. *Clin Exp Rheumatol* 2007; 25: 102-11.
- Radovits BJ, Popa-Diaconu DA, Popa C, *et al.* Disease activity as a risk factor for myocardial infarction in rheumatoid arthritis. *Ann Rheum Dis* 2009; 68: 1271-6.
- Lee YH, Choi SJ, Ji JD, Seo HS, Song GG. Lipoprotein (a) and lipids in relation to inflammation in rheumatoid arthritis. *Clin Rheumatol* 2000; 19: 324-5.
- Seriolo B, Paolino S, Sulli A, Fasciolo D, Cutolo M. Effects of anti-TNF-alpha treatment on lipid profile in patients with active rheumatoid arthritis. *Ann N Y Acad Sci* 2006; 1069: 414-9.
- Svenson KL, Lithell H, Hallgren R, Vessby B. Serum lipoprotein in active rheumatoid arthritis and other chronic inflammatory arthritides. II. Effects of anti-inflammatory and disease-modifying drug treatment. *Arch Intern Med* 1987; 147: 1917-20.
- Hurt-Camejo E, Paredes S, Masana L, *et al.* Elevated levels of small, low-density lipoprotein with high affinity for arterial matrix components in patients with rheumatoid arthritis: possible contribution of phospholipase A2 to this atherogenic profile. *Arthritis Rheum* 2001; 44: 2761-7.
- Choi HK, Seeger JD. Lipid profiles among US elderly with untreated rheumatoid arthritis--the Third National Health and Nutrition Examination Survey. *J Rheumatol* 2005; 32: 2311-6.
- Yoo HW. Dyslipoproteinemia in patients with active rheumatoid arthritis: Effects of disease activity, sex, and menopausal status on lipid profiles. *J Rheumatol* 2004; 31: 1746-53.
- Kelley GA, Kelley KS, Tran ZV. Exercise, lipids, and lipoproteins in older adults: a meta-analysis. *Prev Cardiol* 2005; 8: 206-14.
- Dessein PH, Joffe BI, Stanwix A, Botha AS, Moomal Z. The acute phase response does not fully predict the presence of insulin resistance and dyslipidemia in inflammatory arthritis. *J Rheumatol* 2002; 29: 462-6.
- Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA* 2001; 285: 2486-97.
- Kavanaugh A. Dyslipoproteinaemia in a subset of patients with rheumatoid arthritis. *Ann Rheum Dis* 1994; 53: 551-2.
- Toms TE, Panoulas VF, Douglas KMJ, *et al.* Statin use in rheumatoid arthritis in relation to actual cardiovascular risk: evidence for substantial under treatment of lipid associated cardiovascular risk? Submitted *Ann Rheum Dis* 2009; doi 10.1136/ard.2009.115717 [Epub ahead of print].
- Park YB, Lee SK, Lee WK, *et al.* Lipid profiles in untreated patients with rheumatoid arthritis. *J Rheumatol* 1999; 26: 1701-4.
- Peters MJ, Vis M, van Halm VP, *et al.* Changes in lipid profile during infliximab and corticosteroid treatment in rheumatoid arthritis. *Ann Rheum Dis* 2007; 66: 958-61.
- van Halm VP, Nielen MM, Nurmohamed MT, *et al.* Lipids and inflammation: serial measurements of the lipid profile of blood donors who later developed rheumatoid arthritis. *Ann Rheum Dis* 2007; 66: 184-8.

- [17] Nurmohamed MT. Atherogenic lipid profiles and its management in patients with rheumatoid arthritis. *Vasc Health Risk Manag* 2007; 3: 845-52.
- [18] London MG, Muriden KD, Hewitt JV. Serum cholesterol in rheumatic diseases. *Br Med J* 1963; 1380-3.
- [19] Dursunoglu D, Evrengul H, Polat B, *et al.* Lp(a) lipoprotein and lipids in patients with rheumatoid arthritis: serum levels and relationship to inflammation. *Rheumatol Int* 2005; 25: 241-5.
- [20] Lazarevic MB, Vitic J, Mladenovic V, Myones BL, Skosey JL, Swedler WI. Dyslipoproteinemia in the course of active rheumatoid arthritis. *Semin Arthritis Rheum* 1992; 22(3): 172-8.
- [21] Rantapaa-Dahlqvist S, Wallberg-Jonsson S, Dahlen G. Lipoprotein (a), lipids, and lipoproteins in patients with rheumatoid arthritis. *Ann Rheum Dis* 1991; 50: 366-8.
- [22] Lakator J, Harray S. Serum total, HDL, LDL chol and Triglyceride levels in patients with rheumatoid arthritis. *Clin Biochem* 1988; 21: 93-5.
- [23] Svenson KL, Lithell H, Hallgren R, Selinus I, Vessby B. Serum lipoprotein in active rheumatoid arthritis and other chronic inflammatory arthritides. I. Relativity to inflammatory activity. *Arch Intern Med* 1987; 147: 1912-6.
- [24] Stampfer MJ, Sacks HS, Salvini S, Willett WC, Hennekens CH. A prospective study of cholesterol, apolipoproteins, and the risk of myocardial infarction. *N Engl J Med* 1991; 325: 373-81.
- [25] de Graaf J, Hak-Lemmers HL, Hectors MP, Demacker PN, Hendriks JC, Stalenhoef AF. Enhanced susceptibility to *in vitro* oxidation of the dense low density lipoprotein subfraction in healthy subjects. *Arterioscler Thromb* 1991; 11: 298-306.
- [26] Gazi IF, Tsimihodimos V, Tselepis AD, Elisaf M, Mikhailidis DP. Clinical importance and therapeutic modulation of small dense low-density lipoprotein particles. *Expert Opin Biol Ther* 2007; 7: 53-72.
- [27] Lamarche B, Moorjani S, Cantin B, Dagenais GR, Lupien PJ, Despres JP. Associations of HDL2 and HDL3 subfractions with ischemic heart disease in men. Prospective results from the Quebec Cardiovascular Study. *Arterioscler Thromb Vasc Biol* 1997; 17: 1098-105.
- [28] Florentin M, Liberopoulos EN, Wierzbicki AS, Mikhailidis DP. Multiple actions of high-density lipoprotein. *Curr Opin Cardiol* 2008; 23: 370-8.
- [29] Loscalzo J. Lipoprotein(a). A unique risk factor for atherothrombotic disease. *Arteriosclerosis* 1990; 10: 672-9.
- [30] Tziomalos K, Athyros VG, Wierzbicki AS, Mikhailidis DP. Lipoprotein a: where are we now? *Curr Opin Cardiol* 2009; 24: 351-7.
- [31] Milionis HJ, Mikhailidis DP, Winder AF. Platelets and lipoprotein (a): a brief overview of their role in the pathogenesis of atherothrombosis. *Platelets* 1999; 10: 277-84.
- [32] Danesh J, Collins R, Peto R. Lipoprotein(a) and coronary heart disease. Meta-analysis of prospective studies. *Circulation* 2000; 102: 1082-5.
- [33] Garcia-Gomez C, Nolla JM, Valverde J, Gomez-Gerique JA, Castro MJ, Pinto X. Conventional lipid profile and lipoprotein(a) concentrations in treated patients with rheumatoid arthritis. *J Rheumatol* 2009; 36: 1365.
- [34] Asanuma Y, Kawai S, Aoshima H, Kaburaki J, Mizushima Y. Serum lipoprotein(a) and apolipoprotein(a) phenotypes in patients with rheumatoid arthritis. *Arthritis Rheum* 1999; 42: 443-7.
- [35] Kamanli A, Naziroglu M, Aydilek N, Hacievliyagil C. Plasma lipid peroxidation and antioxidant levels in patients with rheumatoid arthritis. *Cell Biochem Funct* 2004; 22: 53-7.
- [36] Kajanachumpol S, Vanichapuntu M, Veraseritniyom O, Totemchokchayakarn K, Vatanasuk M. Levels of plasma lipid peroxide products and antioxidant status in rheumatoid arthritis. *Southeast Asian J Trop Med Public Health* 2000; 31: 335-8.
- [37] Heinecke JW. Oxidative stress: new approaches to diagnosis and prognosis in atherosclerosis. *Am J Cardiol* 2003; 91: 12A-16A.
- [38] Pemberton PW, Ahmad Y, Bodill H, *et al.* Biomarkers of oxidant stress, insulin sensitivity and endothelial activation in rheumatoid arthritis: a cross-sectional study of their association with accelerated atherosclerosis. *BMC Res Notes* 2009; 2: 83.
- [39] James MJ, van Reyk D, Rye KA, *et al.* Low density lipoprotein of synovial fluid in inflammatory joint disease is mildly oxidised. *Lipids* 1998; 33: 1115-21.
- [40] Aldred S, Banks M, Kitas G, Griffiths HR. Increased levels of oxidised LDL in the plasma of rheumatoid patients with cardiovascular disease: consequences for monocyte scavenger receptor uptake of LDL. *Rheumatology (Oxford)* 2002; 41: 69.
- [41] Kim SH, Lee CK, Lee EY, *et al.* Serum oxidized low-density lipoproteins in rheumatoid arthritis. *Rheumatol Int* 2004; 24: 230-3.
- [42] Georgiadis AN, Voulgari PV, Argyropoulou MI, *et al.* Early treatment reduces the cardiovascular risk factors in newly diagnosed rheumatoid arthritis patients. *Semin Arthritis Rheum* 2008; 38: 13-9.
- [43] Georgiadis AN, Papavasiliou EC, Lourida ES, *et al.* Atherogenic lipid profile is a feature characteristic of patients with early rheumatoid arthritis: effect of early treatment--a prospective, controlled study. *Arthritis Res Ther* 2006; 8: R82.
- [44] Reilly MP, Rader DJ. The metabolic syndrome: more than the sum of its parts? *Circulation* 2003; 108: 1546-51.
- [45] Roubenoff R, Roubenoff RA, Cannon JG, *et al.* Rheumatoid cachexia: cytokine-driven hypermetabolism accompanying reduced body cell mass in chronic inflammation. *J Clin Invest* 1994; 93: 2379-86.
- [46] Stavropoulos-Kalinglou A, Metsios GS, Panoulas VF, *et al.* Underweight and obese states both associate with worse disease activity and physical function in patients with established rheumatoid arthritis. *Clin Rheumatol* 2009; 28: 439-44.
- [47] Chung CP, Oeser A, Solus JF, *et al.* Prevalence of the metabolic syndrome is increased in rheumatoid arthritis and is associated with coronary atherosclerosis. *Atherosclerosis* 2008; 196: 756-63.
- [48] Karvounaris SA, Sidiropoulos PI, Papadakis JA, *et al.* Metabolic syndrome is common among middle-to-older aged Mediterranean patients with rheumatoid arthritis and correlates with disease activity: a retrospective, cross-sectional, controlled, study. *Ann Rheum Dis* 2007; 66: 28-33.
- [49] Zonana-Nacach A, Santana-Sahagun E, Jimenez-Balderas FJ, Camargo-Coronel A. Prevalence and factors associated with metabolic syndrome in patients with rheumatoid arthritis and systemic lupus erythematosus. *J Clin Rheumatol* 2008; 14: 74-7.
- [50] Tracy RP. Inflammation, the metabolic syndrome and cardiovascular risk. *Int J Clin Pract Suppl* 2003; 134: 10-7.
- [51] Choi HK, Hernan MA, Seeger JD, Robins JM, Wolfe F. Methotrexate and mortality in patients with rheumatoid arthritis: a prospective study. *Lancet* 2002; 359: 1173-7.
- [52] Toms TE, Panoulas VF, Douglas KMJ, Kitas GD. Methotrexate therapy associates with a reduced prevalence of the metabolic syndrome in rheumatoid arthritis patients over the age of 60: More than just an anti-inflammatory effect? A cross sectional study. *Arthritis Res Ther* 2009; 11: R110.
- [53] Toms TE, Panoulas VF, Douglas KM, Griffiths HR, Kitas GD. Lack of association between glucocorticoid use and presence of the metabolic syndrome in patients with rheumatoid arthritis: a cross sectional study. *Arthritis Res Ther* 2008; 10: R145.
- [54] Fraser R, Ingram MC, Anderson NH, Morrison C, Davies E, Connell JM. Cortisol effects on body mass, blood pressure, and cholesterol in the general population. *Hypertension* 1999; 33: 1364-8.
- [55] el Shaboury AH, Hayes TM. Hyperlipidaemia in asthmatic patients receiving long-term steroid therapy. *Br Med J* 1973; 2: 85-6.
- [56] Panthakalam S, Bhatnagar D, Klimiuk P. The prevalence and management of hyperglycaemia in patients with rheumatoid arthritis on corticosteroid therapy. *Scott Med J* 2004; 49: 139-41.
- [57] Panoulas VF, Douglas KM, Stavropoulos-Kalinoglou A, *et al.* Long-term exposure to medium-dose glucocorticoid therapy associates with hypertension in patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2008; 47: 72-5.
- [58] Brewer HB Jr, Rader DJ. HDL: structure, function and metabolism. *Prog Lipid Res* 1991; 30: 139-44.
- [59] Zhang B, Saku K, Ohta T. *In vivo* metabolism of HDL, apo A-I, and Lp A-I, and function of HDL--a clinical perspective. *J Atheroscler Thromb* 2000; 7: 59-66.
- [60] Assmann G, Schulte H, von Eckardstein A, Huang Y. High-density lipoprotein cholesterol as a predictor of coronary heart disease risk. The PROCAM experience and pathophysiological implications for reverse cholesterol transport. *Atherosclerosis* 1996; 124 (Suppl: S11): S20.
- [61] Majeed F, Miller M. Low high-density lipoprotein cholesterol: an important consideration in coronary heart disease risk assessment. *Curr Opin Endocrinol Diabetes Obes* 2008; 15: 175-81.
- [62] Goldbourt U, Yaari S, Medalie JH. Isolated low HDL cholesterol as a risk factor for coronary heart disease mortality. A 21-year fol-

- low-up of 8000 men. *Arterioscler Thromb Vasc Biol* 1997; 17: 107-13.
- [63] Gordon T, Castelli WP, Hjortland MC, Kannel WB, Dawber TR. High density lipoprotein as a protective factor against coronary heart disease. The Framingham Study. *Am J Med* 1977; 62: 707-14.
- [64] Robins SJ, Collins D, Wittes JT, *et al.* Relation of gemfibrozil treatment and lipid levels with major coronary events: VA-HIT: a randomized controlled trial. *JAMA* 2001; 285: 1585-91.
- [65] Kalogirou M, Tsimihodimos V, Gazi I, *et al.* Effect of ezetimibe monotherapy on the concentration of lipoprotein subfractions in patients with primary dyslipidaemia. *Curr Med Res Opin* 2007; 23: 1169-76.
- [66] Rubins HB, Robins SJ, Collins D, *et al.* Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group. *N Engl J Med* 1999; 341: 410-8.
- [67] Alvarez C, Ramos A. Lipids, lipoproteins, and apoproteins in serum during infection. *Clin Chem* 1986; 32(1 Pt 1): 142-5.
- [68] Sammalcorpi K, Valtonen V, Kerttula Y, Nikkila E, Taskinen MR. Changes in serum lipoprotein pattern induced by acute infections. *Metabolism* 1988; 37: 859-65.
- [69] van Leeuwen HJ, Heezius EC, Dallinga GM, van Strijp JA, Verhoef J, van Kessel KP. Lipoprotein metabolism in patients with severe sepsis. *Crit Care Med* 2003; 31: 1359-66.
- [70] Rodriguez RJJ, Iglesias CG, Vazquez M, *et al.* Variation in plasma lipid and lipoprotein concentrations in community-acquired pneumonia a six-month prospective study. *Eur J Clin Chem Clin Biochem* 1996; 34: 245-9.
- [71] Vottery R, Saigal R, Singhal N, Gupta BS. Lipid profile in rheumatoid arthritis and its relation to disease activity. *J Assoc Physicians India* 2001; 49: 1188-90.
- [72] Park YB, Choi HK, Kim MY, *et al.* Effects of antirheumatic therapy on serum lipid levels in patients with rheumatoid arthritis: a prospective study. *Am J Med* 2002; 113: 188-93.
- [73] Popa C, van den Hoogen FH, Radstake TR, *et al.* Modulation of lipoprotein plasma concentrations during long-term anti-TNF therapy in patients with active rheumatoid arthritis. *Ann Rheum Dis* 2007; 66: 1503-7.
- [74] Malle E, Steinmetz A, Raynes JG. Serum amyloid A (SAA): an acute phase protein and apolipoprotein. *Atherosclerosis* 1993; 102: 131-46.
- [75] Pepys MB, Baltz ML. Acute phase proteins with special reference to C-reactive protein and related proteins (pentaxins) and serum amyloid A protein. *Adv Immunol* 1983; 34: 141-212.
- [76] Zahedi K, Gonnerman WA, Debeer FC, *et al.* Major acute-phase reactant synthesis during chronic inflammation in amyloid-susceptible and -resistant mouse strains. *Inflammation* 1991; 15: 1-14.
- [77] Benditt EP, Eriksen N. Amyloid protein SAA is associated with high density lipoprotein from human serum. *Proc Natl Acad Sci USA* 1977; 74: 4025-8.
- [78] Cabana VG, Lukens JR, Rice KS, Hawkins TJ, Getz GS. HDL content and composition in acute phase response in three species: triglyceride enrichment of HDL a factor in its decrease. *J Lipid Res* 1996; 37: 2662-74.
- [79] Pruzanski W, Stefanski E, de Beer FC, de Beer MC, Ravandi A, Kuksis A. Comparative analysis of lipid composition of normal and acute-phase high density lipoproteins. *J Lipid Res* 2000; 41: 1035-47.
- [80] Clifton PM, Mackinnon AM, Barter PJ. Effects of serum amyloid A protein (SAA) on composition, size, and density of high density lipoproteins in subjects with myocardial infarction. *J Lipid Res* 1985; 26: 1389-98.
- [81] Kisilevsky R, Subrahmanyam L. Serum amyloid A changes high density lipoprotein's cellular affinity. A clue to serum amyloid A's principal function. *Lab Invest* 1992; 66: 778-85.
- [82] Banka CL, Yuan T, de Beer MC, Kindy M, Curtiss LK, de Beer FC. Serum amyloid A (SAA): influence on HDL-mediated cellular cholesterol efflux. *J Lipid Res* 1995; 36: 1058-65.
- [83] Rye KA, Duong MN. Influence of phospholipid depletion on the size, structure, and remodeling of reconstituted high density lipoproteins. *J Lipid Res* 2000; 41: 1640-50.
- [84] Ehrenwald E, Chisolm GM, Fox PL. Intact human ceruloplasmin oxidatively modifies low density lipoprotein. *J Clin Invest* 1994; 93: 1493-501.
- [85] Tietge UJ, Maugeais C, Lund-Katz S, Grass D, Debeer FC, Rader DJ. Human secretory phospholipase A2 mediates decreased plasma levels of HDL cholesterol and apoA-I in response to inflammation in human apoA-I transgenic mice. *Arterioscler Thromb Vasc Biol* 2002; 22: 1213-8.
- [86] Van Lenten BJ, Hama SY, de Beer FC, *et al.* Anti-inflammatory HDL becomes pro-inflammatory during the acute phase response. Loss of protective effect of HDL against LDL oxidation in aortic wall cell cocultures. *J Clin Invest* 1995; 96: 2758-67.
- [87] Samokyszyn VM, Miller DM, Reif DW, Aust SD. Inhibition of superoxide and ferritin-dependent lipid peroxidation by ceruloplasmin. *J Biol Chem* 1989; 264: 21-6.
- [88] Chisolm GM, III, Hazen SL, Fox PL, Cathcart MK. The oxidation of lipoproteins by monocytes-macrophages. Biochemical and biological mechanisms. *J Biol Chem* 1999; 274: 25959-62.
- [89] Kunitake ST, Jarvis MR, Hamilton RL, Kane JP. Binding of transition metals by apolipoprotein A-I-containing plasma lipoproteins: inhibition of oxidation of low density lipoproteins. *Proc Natl Acad Sci USA* 1992; 89: 6993-7.
- [90] Ritchie RF, Palomaki GE, Neveux LM, Navolotskaia O, Ledue TB, Craig WY. Reference distributions for the negative acute-phase serum proteins, albumin, transferrin and transthyretin: a practical, simple and clinically relevant approach in a large cohort. *J Clin Lab Anal* 1999; 13: 273-9.
- [91] Bush J, Richardson J, Cardelli J. Molecular cloning and characterization of the full-length cDNA encoding the developmentally regulated lysosomal enzyme beta-glucosidase in *Dictyostelium discoideum*. *J Biol Chem* 1994; 269: 1468-76.
- [92] Zambon A, Deeb SS, Bensadoun A, Foster KE, Brunzell JD. *In vivo* evidence of a role for hepatic lipase in human apoB-containing lipoprotein metabolism, independent of its lipolytic activity. *J Lipid Res* 2000; 41: 2094-9.
- [93] Fluiter K, van Berkel TJ. Scavenger receptor B1 (SR-B1) substrates inhibit the selective uptake of high-density-lipoprotein cholesteryl esters by rat parenchymal liver cells. *Biochem J* 1997; 326 (Pt 2): 515-9.
- [94] Kawakami M, Murase T, Itakura H, Yamada N, Ohsawa N, Takaku F. Lipid metabolism in endotoxic rats: decrease in hepatic triglyceride lipase activity. *Microbiol Immunol* 1986; 30: 849-54.
- [95] Levy E, Gurbindo C, Lacaille F, Paradis K, Thibault L, Seidman E. Circulating tumor necrosis factor- α levels and lipid abnormalities in patients with cystic fibrosis. *Pediatr Res* 1993; 34: 162-6.
- [96] Grunfeld C, Pang M, Doerrler W, Shigenaga JK, Jensen P, Feingold KR. Lipids, lipoproteins, triglyceride clearance, and cytokines in human immunodeficiency virus infection and the acquired immunodeficiency syndrome. *J Clin Endocrinol Metab* 1992; 74: 1045-52.
- [97] Feingold KR, Pollock AS, Moser AH, Shigenaga JK, Grunfeld C. Discordant regulation of proteins of cholesterol metabolism during the acute phase response. *J Lipid Res* 1995; 36: 1474-82.
- [98] Tall AR. Plasma cholesteryl ester transfer protein. *J Lipid Res* 1993; 34: 1255-74.
- [99] Masucci-Magoulas L, Moulin P, Jiang XC, *et al.* Decreased cholesteryl ester transfer protein (CETP) mRNA and protein and increased high density lipoprotein following lipopolysaccharide administration in human CETP transgenic mice. *J Clin Invest* 1995; 95: 1587-94.
- [100] Jiang XC, Bruce C. Regulation of murine plasma phospholipid transfer protein activity and mRNA levels by lipopolysaccharide and high cholesterol diet. *J Biol Chem* 1995; 270: 17133-8.
- [101] Jiang XC, Bruce C, Mar J, *et al.* Targeted mutation of plasma phospholipid transfer protein gene markedly reduces high-density lipoprotein levels. *J Clin Invest* 1999; 103: 907-14.
- [102] Deakin S, Moren X, James RW. HDL oxidation compromises its influence on paraoxonase-1 secretion and its capacity to modulate enzyme activity. *Arterioscler Thromb Vasc Biol* 2007; 27: 1146-52.
- [103] Khovidhunkit W, Memon RA, Shigenaga JK, *et al.* Plasma platelet-activating factor acetylhydrolase activity in human immunodeficiency virus infection and the acquired immunodeficiency syndrome. *Metabolism* 1999; 48: 1524-31.

- [104] Dulioust A, Hilliquin P, Menkes CJ, Benveniste J, Arnoux B. Paf-acether acetylhydrolase activity is increased in patients with rheumatic diseases. *Scand J Rheumatol* 1992; 21: 161-4.
- [105] Memon RA, Fuller J, Moser AH, Feingold KR, Grunfeld C. *In vivo* regulation of plasma platelet-activating factor acetylhydrolase during the acute phase response. *Am J Physiol* 1999; 277(1 Pt 2): R94-103.
- [106] Quinn MT, Parthasarathy S, Steinberg D. Lysophosphatidylcholine: a chemotactic factor for human monocytes and its potential role in atherogenesis. *Proc Natl Acad Sci USA* 1988; 85: 2805-9.
- [107] Tetta C, Bussolino F, Modena V, *et al.* Release of platelet-activating factor in systemic lupus erythematosus. *Int Arch Allergy Appl Immunol* 1990; 91: 244-56.
- [108] Tselepis AD, Elisaf M, Besis S, Karabina SA, Chapman MJ, Siomopoulou A. Association of the inflammatory state in active juvenile rheumatoid arthritis with hypo-high-density lipoproteinemia and reduced lipoprotein-associated platelet-activating factor acetylhydrolase activity. *Arthritis Rheum* 1999; 42: 373-83.
- [109] Bansal S, Buring JE, Rifai N, Mora S, Sacks FM, Ridker PM. Fasting compared with nonfasting triglycerides and risk of cardiovascular events in women. *CMAJ* 2007; 298: 309-16.
- [110] Sattar N, Packard CJ, Petrie JR. The end of triglycerides in cardiovascular risk assessment? Rumours of death are greatly exaggerated. *BMJ* 1998; 317: 553-54.
- [111] Gotto AM, Jr. Triglyceride as a risk factor for coronary artery disease. *Am J Cardiol* 1998; 82: 22Q-25Q.
- [112] Gallin JI, Kaye D, O'Leary WM. Serum lipids in infection. *N Engl J Med* 1969; 281: 1081-6.
- [113] McDevitt H, Munson S, Ettinger R, Wu A. Multiple roles for tumor necrosis factor- α and lymphotoxin α/β in immunity and autoimmunity. *Arthritis Res* 2002; 4 (Suppl 3): S141-S152.
- [114] Memon RA, Holleran WM, Moser AH, *et al.* Endotoxin and cytokines increase hepatic sphingolipid biosynthesis and produce lipoproteins enriched in ceramides and sphingomyelin. *Arterioscler Thromb Vasc Biol* 1998; 18: 1257-65.
- [115] Feingold KR, Soued M, Staprans I, *et al.* Effect of tumor necrosis factor (TNF) on lipid metabolism in the diabetic rat. Evidence that inhibition of adipose tissue lipoprotein lipase activity is not required for TNF-induced hyperlipidemia. *J Clin Invest* 1989; 83: 1116-21.
- [116] Skoog T, Dichtl W, Boquist S, *et al.* Plasma tumour necrosis factor- α and early carotid atherosclerosis in healthy middle-aged men. *Eur Heart J* 2002; 23: 376-83.
- [117] Mizia-Stec K, Zahorska-Markiewicz B, Mandecki T, *et al.* Hyperlipidaemias and serum cytokines in patients with coronary artery disease. *Acta Cardiol* 2003; 58: 9-15.
- [118] Svenungsson E, Fei GZ, Jensen-Ustad K, de Faire U, Hamsten A, Frostegard J. TNF- α : a link between hypertriglyceridaemia and inflammation in SLE patients with cardiovascular disease. *Lupus* 2003; 12: 454-61.
- [119] Allanore Y, Kahan A, Sellam J, Ekindjian OG, Borderie D. Effects of repeated infliximab therapy on serum lipid profile in patients with refractory rheumatoid arthritis. *Clin Chim Acta* 2006; 365: 143-8.
- [120] Del Porto F, Lagana B, Nofroni I, Tinti F, Mitterhofer AP, D'Amelio R. Effects of tumour necrosis factor α blockade on lipid profile in active rheumatoid arthritis. *Rheumatology (Oxford)* 2007; 46: 1626-7.
- [121] Dahlqvist SR, Engstrand S, Berglin E, Johnson O. Conversion towards an atherogenic lipid profile in rheumatoid arthritis patients during long-term infliximab therapy. *Scand J Rheumatol* 2006; 35: 107-11.
- [122] Kiortsis DN, Mayrdis AK, Filippatos TD, Vasakos S, Nikas SN, Drosos AA. Effects of infliximab treatment on lipoprotein profile in patients with rheumatoid arthritis and ankylosing spondylitis. *J Rheumatol* 2006; 33: 921-3.
- [123] Yudkin JS, Kumari M, Humphries SE, Mohamed-Ali V. Inflammation, obesity, stress and coronary heart disease: is interleukin-6 the link? *Atherosclerosis* 2000; 148: 209-14.
- [124] Kapoor S. Interleukin-6 antagonists for the management of hypertension. *Hypertension* 2007; 49: e18.
- [125] Smolen JS, Beaulieu A, Rubbert-Roth A, *et al.* Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (OPTION study): a double-blind, placebo-controlled, randomised trial. *Lancet* 2008; 371: 987-97.
- [126] Glund S, Krook A. Role of interleukin-6 signalling in glucose and lipid metabolism. *Acta Physiol (Oxf)* 2008; 192: 37-48.
- [127] Fernandez-Real JM, Broch M, Vendrell J, Richart C, Ricart W. Interleukin-6 gene polymorphism and lipid abnormalities in healthy subjects. *J Clin Endocrinol Metab* 2000; 85: 1334-9.
- [128] Mendall MA, Patel P, Asante M, *et al.* Relation of serum cytokine concentrations to cardiovascular risk factors and coronary heart disease. *Heart* 1997; 78: 273-7.
- [129] Sutherland JP, McKinley B, Eckel RH. The metabolic syndrome and inflammation. *Metab Syndr Relat Disord* 2004; 2: 82-104.
- [130] Hamid YH, Rose CS, Urhammer SA, *et al.* Variations of the interleukin-6 promoter are associated with features of the metabolic syndrome in Caucasian Danes. *Diabetologia* 2005; 48: 251-60.
- [131] Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 2001; 286: 327-34.
- [132] Sweep CG, Hermus RM, van der Meer MJ, *et al.* Chronic intraperitoneal infusion of low doses of tumor necrosis factor α in rats induces a reduction in plasma triglyceride levels. *Cytokine* 1992; 4: 561-7.
- [133] Nawawi H, Osman NS, Annur R, Khalid BA, Yusoff K. Soluble intercellular adhesion molecule-1 and interleukin-6 levels reflect endothelial dysfunction in patients with primary hypercholesterolaemia treated with atorvastatin. *Atherosclerosis* 2003; 169: 283-91.
- [134] Mead JR, Irvine SA, Ramji DP. Lipoprotein lipase: structure, function, regulation, and role in disease. *J Mol Med* 2002; 80: 753-69.
- [135] Zechner R, Newman TC, Sherry B, Cerami A, Breslow JL. Recombinant human cachectin/tumor necrosis factor but not interleukin-1 α downregulates lipoprotein lipase gene expression at the transcriptional level in mouse 3T3-L1 adipocytes. *Mol Cell Biol* 1988; 8: 2394-401.
- [136] Ziouzenkova O, Plutzky J. Lipolytic PPAR activation: new insights into the intersection of triglycerides and inflammation? *Curr Opin Clin Nutr Metab Care* 2004; 7: 369-375.
- [137] Grunfeld C, Gulli R, Moser AH, Gavin LA, Feingold KR. Effect of tumor necrosis factor administration *in vivo* on lipoprotein lipase activity in various tissues of the rat. *J Lipid Res* 1989; 30: 579-85.
- [138] Gervois P, Torra IP, Fruchart JC, Staels B. Regulation of lipid and lipoprotein metabolism by PPAR activators. *Clin Chem Lab Med* 2000; 38(1):3-11.
- [139] Ziouzenkova O, Perrey S, Asatryan L, *et al.* Lipolysis of triglyceride-rich lipoproteins generates PPAR ligands: evidence for an anti-inflammatory role for lipoprotein lipase. *Proc Natl Acad Sci USA* 2003; 100(5):2730-35.
- [140] Auwerx J, Schoonjans K, Fruchart JC, Staels B. Regulation of triglyceride metabolism by PPARs: fibrates and thiazolidinediones have distinct effects. *J Atheroscler Thromb* 1996; 3(2):81-89.
- [141] Vu-Dac N, Schoonjans K, Laine B, Fruchart JC, Auwerx J, Staels B. Negative regulation of the human apolipoprotein A-I promoter by fibrates can be attenuated by the interaction of the peroxisome proliferator-activated receptor with its response element. *J Biol Chem* 1994; 269(49): 31012-8.
- [142] Vu-Dac N, Schoonjans K, Kosykh V, *et al.* Fibrates increase human apolipoprotein A-II expression through activation of the peroxisome proliferator-activated receptor. *J Clin Invest* 1995; 96: 741-50.
- [143] Howard BV, Robbins DC, Sievers ML, *et al.* LDL cholesterol as a strong predictor of coronary heart disease in diabetic individuals with insulin resistance and low LDL: the strong heart study. *Arterioscler Thromb Vasc Biol* 2000; 20: 830-5.
- [144] Wood DA. JBS 2: Joint British Societies' guidelines on prevention of cardiovascular disease in clinical practice. *Heart* 2005; 91 (Suppl 5): v1-52.
- [145] Cannon CP, Steinberg BA, Murphy SA, Mega JL, Braunwald E. Meta-analysis of cardiovascular outcomes trials comparing intensive vs moderate statin therapy. *J Am Coll Cardiol* 2006; 48(3): 438-45.
- [146] LaRosa JC, Grundy SM, Waters DD, *et al.* Intensive lipid lowering with atorvastatin in patients with stable coronary disease. *N Engl J Med* 2005; 352: 1425-35.
- [147] Wenger NK, Lewis SJ, Welty FK, Herrington DM, Bittner V. Beneficial effects of aggressive low-density lipoprotein cholesterol lowering in women with stable coronary heart disease in the Treating to New Targets (TNT) study. *Heart* 2008; 94: 434-9.

- [148] Carroll MD, Lacher DA, Sorlie PD, *et al.* Trends in serum lipids and lipoproteins of adults, 1960-2002. *JAMA* 2005; 294(14): 1773-81.
- [149] Floris-Moore M, Howard AA, Lo Y, Schoenbaum EE, Arnsten JH, Klein RS. Hepatitis C infection is associated with lower lipids and high-sensitivity C-reactive protein in HIV-infected men. *AIDS Patient Care STDS* 2007; 21(7): 479-91.
- [150] Khan M, Pelengaris S, Cooper M, Smith C, Evan G, Betteridge J. Oxidised lipoproteins may promote inflammation through the selective delay of engulfment but not binding of apoptotic cells by macrophages. *Atherosclerosis* 2003; 171(1): 21-9.
- [151] Khouidhunkit W, Memon RA, Feingold KR, Grunfield C. Infection and inflammation induced proatherogenic changes of lipoproteins. *J Infect Dis* 2000; 181: s462-72.
- [152] Rizzo M, Berneis K. Low-density lipoprotein size and cardiovascular risk assessment. *QJM* 2006; 99(1): 1-14.
- [153] Jones SM, Harris CP, Lloyd J, Stirling CA, Reckless JP, McHugh NJ. Lipoproteins and their subfractions in psoriatic arthritis: identification of an atherogenic profile with active joint disease. *Ann Rheum Dis* 2000; 59(11): 904-9.
- [154] Feingold KR, Krauss RM, Pang M, Doerrler W, Jensen P, Grunfeld C. The hypertriglyceridemia of acquired immunodeficiency syndrome is associated with an increased prevalence of low density lipoprotein subclass pattern B. *J Clin Endocrinol Metab* 1993; 76(6): 1423-7.
- [155] Halliwell B. Free radicals, antioxidants, and human disease: curiosity, cause, or consequence? *Lancet* 1994; 344(8924): 721-4.
- [156] Memon RA, Staprans I, Noor M, *et al.* Infection and inflammation induce LDL oxidation *in vivo*. *Arterioscler Thromb Vasc Biol* 2000; 20(6): 1536-42.
- [157] Orem A, Yandi YE, Vanizor B, Cimsit G, Uydu HA, Malkoc M. The evaluation of autoantibodies against oxidatively modified low-density lipoprotein (LDL), susceptibility of LDL to oxidation, serum lipids and lipid hydroperoxide levels, total antioxidant status, antioxidant enzyme activities, and endothelial dysfunction in patients with Behcet's disease. *Clin Biochem* 2002; 35(3): 217-24.
- [158] Paredes S, Girona J, Hurt-Camejo E, *et al.* Antioxidant vitamins and lipid peroxidation in patients with rheumatoid arthritis: association with inflammatory markers. *J Rheumatol* 2002; 29(11): 2271-7.
- [159] Karabina SA, Brocheriou I, Le Naour G, *et al.* Atherogenic properties of LDL particles modified by human group X secreted phospholipase A2 on human endothelial cell function. *FASEB J* 2006; 20(14): 2547-9.
- [160] Hurt-Camejo E, Camejo G, Sartipy P. Phospholipase A2 and small, dense low-density lipoprotein. *Curr Opin Lipidol* 2000; 11(5): 465-71.
- [161] Leitinger N, Watson AD, Hama SY, *et al.* Role of group II secretory phospholipase A2 in atherosclerosis: 2. Potential involvement of biologically active oxidized phospholipids. *Arterioscler Thromb Vasc Biol* 1999; 19(5): 1291-8.
- [162] Chang MK, Binder CJ, Torzewski M, Witztum JL. C-reactive protein binds to both oxidized LDL and apoptotic cells through recognition of a common ligand: Phosphorylcholine of oxidized phospholipids. *Proc Natl Acad Sci USA* 2002; 99(20): 13043-8.
- [163] Jaye M, Lynch KJ, Krawiec J, *et al.* A novel endothelial-derived lipase that modulates HDL metabolism. *Nat Genet* 1999; 21(4): 424-8.
- [164] Fuki IV, Blanchard N, Jin W, *et al.* Endogenously produced endothelial lipase enhances binding and cellular processing of plasma lipoproteins *via* heparan sulfate proteoglycan-mediated pathway. *J Biol Chem* 2003; 278(36): 34331-8.
- [165] Badellino KO, Wolfe ML, Reilly MP, Rader DJ. Endothelial lipase is increased *in vivo* by inflammation in humans. *Circulation* 2008; 117(5): 678-85.
- [166] Paradis ME, Badellino KO, Rader DJ, *et al.* Endothelial lipase is associated with inflammation in humans. *J Lipid Res* 2006; 47(12): 2808-13.
- [167] Ishida T, Choi S, Kundu RK, *et al.* Endothelial lipase is a major determinant of HDL level. *J Clin Invest* 2003; 111(3): 347-55.
- [168] Ma K, Cilingiroglu M, Orvos JD, Ballantyne CM, Marian AJ, Chan L. Endothelial lipase is a major genetic determinant for high-density lipoprotein concentration, structure, and metabolism. *Proc Natl Acad Sci USA* 2003; 100(5): 2748-53.
- [169] Yasuda T, Hirata K, Ishida T, *et al.* Endothelial lipase is increased by inflammation and promotes LDL uptake in macrophages. *J Atheroscler Thromb* 2007; 14(4): 192-201.
- [170] Kelly GA, Kelly KS, Tran ZV. Exercise, lipids and lipoproteins in older adults: a meta-analysis. *Prev Cardiol* 2005; 8(4): 206-14.
- [171] Marcovina SM, Gaur VP, Albers JJ. Biological variability of cholesterol, triglyceride, low- and high-density lipoprotein cholesterol, lipoprotein(a), and apolipoproteins A-I and B. *Clin Chem* 1994; 40(4): 574-8.
- [172] Gordon DJ, Trost DC, Hyde J, *et al.* Seasonal cholesterol cycles: the lipid research clinics coronary primary prevention trial placebo group. *Circulation* 1987; 76(6): 1224-31.
- [173] Gordon DJ, Hyde J, Trost DC, *et al.* Cyclic seasonal variation in plasma lipid and lipoprotein levels: the lipid research clinics coronary primary prevention trial placebo group. *J Clin Epidemiol* 1988; 41(7): 679-89.
- [174] Smith SJ, Cooper GR, Myers GL, Sampson EJ. Biological variability in concentrations of serum lipids: sources of variation among results from published studies and composite predicted values. *Clin Chem* 1993; 39(6): 1012-22.
- [175] Hadaegh F, Harati H, Zabetian A, Azizi F. Seasonal variability of serum lipids in adults: Tehran Lipid and Glucose Study. *Med J Malaysia* 2006; 61(3): 332-8.
- [176] Ockene IS, Chiriboga DE, Stanek EJ, III, *et al.* Seasonal variation in serum cholesterol levels: treatment implications and possible mechanisms. *Arch Intern Med* 2004; 164(8): 863-70.
- [177] Buxtorf JC, Baudet MF, Martin C, Richard JL, Jacotot B. Seasonal variations of serum lipids and apoproteins. *Ann Nutr Metab* 1988; 32(2): 68-74.
- [178] Riemersma RA, Wilson R, Payne JA, Shepherd MJ. Seasonal variation in copper-mediated low density lipoprotein oxidation *in vitro* is related to varying plasma concentration of oxidised lipids in the summer and winter. *Free Radic Res* 2003; 37(3): 341-7.
- [179] Woodhouse PR, Khaw KT. Seasonal variation of risk factors for cardiovascular disease and diet in older adults. *Int J Circumpolar Health* 2000; 59(3-4): 204-9.
- [180] Kristal-Boneh E, Froom P, Harari G, Shapiro Y, Green MS. Seasonal changes in red blood cell parameters. *Br J Haematol* 1993; 85(3): 603-7.
- [181] Heller RF, Jacobs HS. Coronary heart disease in relation to age, sex and the menopause. *Br Med J* 1978; 1: 472-4.
- [182] Stevenson JC, Crook D, Goddard IF. Influence of age and menopause on serum lipids and lipoproteins in healthy women. *Atherosclerosis* 1993; 98: 83-90.
- [183] Hjortland MC, MnNamara PM, Kannel WB. Some atherogenic concomitants of menopause: the Framingham Study. *Am J Epidemiol* 1976; 103: 304-11.
- [184] Wu Z, Wu X, Zhang Y. Relationship of menopausal status and sex hormones to serum lipids and blood pressure. *Int J Epidemiol* 1990; 19(2): 297-302.
- [185] Shaw LJ, Bairey Merz CN, Pepine CJ, *et al.* Insights from the NHLBI-sponsored Women's Ischaemia Syndrome Evaluation (WISE) study; Part I: gender differences in traditional and novel risk factors, symptom evaluation, and gender-optimised diagnostic strategies. *J Am Coll Cardiol* 2006; 47: S4-S20.
- [186] Middleberg RPS, Spector TD, Swaminathan R, Sneider H. Genetic and environmental influences on lipids, lipoproteins, and apolipoproteins. Effects of the menopause. *Arterioscler Thromb Vasc Biol* 2002; 22: 1142-7.
- [187] Pickar JH, Wild RA, Walsh B, Hirvonen E, Lobo RA. Effects of different hormone replacement regimens on postmenopausal women with abnormal lipid levels. *Menopause Study Group. Climacteric* 1998; 1(1): 26-32.
- [188] Mosca L, Collins P, Herrington DM, *et al.* Hormone replacement therapy and cardiovascular disease: a statement for healthcare professionals from the American Heart Association. *Circulation* 2001; 104(4): 499-503.
- [189] Silvestri A, Gebara O, Vitale C, *et al.* Increased levels of C-reactive protein after oral hormone replacement therapy may not be related to an increased inflammatory response. *Circulation* 2003; 107(25): 3165-9.
- [190] Vitale C, Gebara O, Mercuro G, *et al.* Value of C-reactive protein levels and IL-6 in predicting events levels in women at increased cardiovascular risk. *Maturitas* 2005; 50(4): 239-46.
- [191] Cox KL, Burke V, Morton AR, Gillam HF, Beilin LJ, Puddey IB. Long-term effects of exercise on blood pressure and lipids in

- healthy women aged 40-65 years: The Sedentary Women Exercise Adherence Trial (SWEAT). *J Hypertens* 2001; 19(10): 1733-43.
- [192] Boardley D, Fahlman M, Topp R, Morgan AL, McNevin N. The impact of exercise training on blood lipids in older adults. *Am J Geriatr Cardiol* 2007; 16(1): 30-5.
- [193] Sugiura H, Sugiura H, Kajima K, Mirbod SM, Iwata H, Matsuoka T. Effects of long-term moderate exercise and increase in number of daily steps on serum lipids in women: randomised controlled trial [ISRCTN21921919]. *BMC Womens Health* 2002; 2(1): 3.
- [194] Thompson PD, Cullinane EM, Sady SP, *et al.* Modest changes in high-density lipoprotein concentration and metabolism with prolonged exercise training. *Circulation* 1988; 78(1): 25-34.
- [195] Durstine JL, Haskell WL. Effects of exercise training on plasma lipids and lipoproteins. *Exerc Sport Sci Rev* 1994; 22: 477-521.
- [196] Halverstadt A, Phares DA, Wilund KR, Goldberg AP, Hagberg JM. Endurance exercise training raises high-density lipoprotein cholesterol and lowers small low-density lipoprotein and very low-density lipoprotein independent of body fat phenotypes in older men and women. *Metabolism* 2007; 56(4): 444-50.
- [197] Ainslie PN, Reilly T, Maclaren DP, Campbell IT. Changes in plasma lipids and lipoproteins following 10-days of prolonged walking: influence of age and relationship to physical activity level. *Ergonomics* 2005; 48(11-14): 1352-64.
- [198] Tsekouras YE, Yanni AE, Bougatsas D, Kavouras SA, Sidossis LS. A single bout of brisk walking increases basal very low-density lipoprotein triacylglycerol clearance in young men. *Metabolism* 2007; 56(8): 1037-43.
- [199] Crouse SF, O'Brien BC, Grandjean PW, Lowe RC, Rohack JJ, Green JS. Effects of training and a single session of exercise on lipids and apolipoproteins in hypercholesterolemic men. *J Appl Physiol* 1997; 83(6):2019-28.
- [200] Gordon PM, Goss FL, Visich PS, *et al.* The acute effects of exercise intensity on HDL-C metabolism. *Med Sci Sports Exerc* 1994; 26(6): 671-7.
- [201] Ebisu T. Splitting the distance of endurance running on cardiovascular endurance and blood lipids. *Jpn J Physical Ed* 1995; 30(1): 37-43.
- [202] Sgouraki E, Tsopanakis A, Tsopanakis C. Acute exercise: response of HDL-C, LDL-C lipoproteins and HDL-C subfractions levels in selected sport disciplines. *J Sports Med Phys Fitness* 2001; 41(3): 386-91.
- [203] Park DH, Ransone JW. Effects of submaximal exercise on high-density lipoprotein-cholesterol subfractions. *Int J Sports Med* 2003; 24(4): 245-51.
- [204] Fernandez-Pardo J, Rubies-Prat J, Pedro-Botet J, *et al.* High density lipoprotein subfractions and physical activity: changes after moderate and heavy exercise training. *Rev Esp Fisiol* 1991; 47(4): 181-6.
- [205] Rashid S, Genest J. Effect of obesity on high-density lipoprotein metabolism. *Obesity (Silver Spring)* 2007; 15(12): 2875-88.
- [206] Herbert PN, Bernier DN, Cullinane EM, Edelstein L, Kantor MA, Thompson PD. High-density lipoprotein metabolism in runners and sedentary men. *JAMA* 1984; 252(8): 1034-7.
- [207] Ekdahl C, Broman G. Muscle strength, endurance, and aerobic capacity in rheumatoid arthritis: a comparative study with healthy subjects. *Ann Rheum Dis* 1992; 51(1): 35-40.
- [208] Metsios GS, Stavropoulos-Kalinglou A, Panoulas VF, *et al.* Association of physical inactivity with increased cardiovascular risk in patients with rheumatoid arthritis. *Eur J Cardiovascular Prev Rehab* 2009; 16(2): 188-94.
- [209] de Jong Z, Munneke M, Zwinderman AH, *et al.* Is a long-term high-intensity exercise program effective and safe in patients with rheumatoid arthritis? Results of a randomized controlled trial. *Arthritis Rheum* 2003; 48(9): 2415-24.
- [210] Lineker SC, Bell MJ, Wilkins AL, Badley EM. Improvements following short term home based physical therapy are maintained at one year in people with moderate to severe rheumatoid arthritis. *J Rheumatol* 2001; 28(1): 165-8.
- [211] Hakkinen A, Hakkinen K, Hannonen P. Effects of strength training on neuromuscular function and disease activity in patients with recent-onset inflammatory arthritis. *Scand J Rheumatol* 1994; 23(5): 237-42.
- [212] Van den Ende CH, Hazes JM, le Cessie S, *et al.* Comparison of high and low intensity training in well controlled rheumatoid arthritis. Results of a randomised clinical trial. *Ann Rheum Dis* 1996; 55(11): 798-805.
- [213] Van den Ende CH, Breedveld FC, le Cessie S, Dijkmans BA, de Mug AW, Hazes JM. Effect of intensive exercise on patients with active rheumatoid arthritis: a randomised clinical trial. *Ann Rheum Dis* 2000; 59(8): 615-21.
- [214] Miller WM, Nori-Janosz KE, Lillystone M, Yanez J, McCullough PA. Obesity and lipids. *Curr Cardiol Rep* 2005; 7(6): 465-70.
- [215] Pelkonen R, Nikkila EA, Koskinen S, Penttinen K, Sarna S. Association of serum lipids and obesity with cardiovascular mortality. *Br Med J* 1977; 2(6096): 1185-7.
- [216] Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA* 2001; 285(19): 2486-97.
- [217] Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998; 15(7): 539-53.
- [218] Alberti KG, Zimmet P, Shaw J. The metabolic syndrome--a new worldwide definition. *Lancet* 2005; 366(9491): 1059-62.
- [219] Balkau B, Charles MA. Comment on the provisional report from the WHO consultation. *Diabet Med* 1999; 16: 442-3.
- [220] Howard BV, Ruotolo G, Robbins DC. Obesity and dyslipidemia. *Endocrinol Metab Clin North Am* 2003; 32(4): 855-67.
- [221] Adiels M, Taskinen MR, Packard C, *et al.* Overproduction of large VLDL particles is driven by increased liver fat content in man. *Diabetologia* 2006; 49(4): 755-65.
- [222] Frenais R, Nazih H, Ouguerram K, *et al.* *In vivo* evidence for the role of lipoprotein lipase activity in the regulation of apolipoprotein AI metabolism: a kinetic study in control subjects and patients with type II diabetes mellitus. *J Clin Endocrinol Metab* 2001; 86(5): 1962-7.
- [223] Van Gaal LF, Mertens IL, De Block CE. Mechanisms linking obesity with cardiovascular disease. *Nature* 2006; 444(7121): 875-80.
- [224] Metsios GS, Stavropoulos-Kalinglou A, Koutedakis Y, *et al.* A new resting energy expenditure prediction equation for patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2008; 47(4): 500-6.
- [225] Stavropoulos-Kalinoglou A, Metsios GS, Koutedakis Y, *et al.* Redefining overweight and obesity in rheumatoid arthritis patients. *Ann Rheum Dis* 2007; 66(10): 1316-21.
- [226] Stavropoulos-Kalinglou A, Metsios GS, Panoulas VF, *et al.* Associations of obesity with modifiable risk factors for the development of cardiovascular disease in patients with rheumatoid arthritis. *Ann Rheum Dis* 2009; 68(2): 242-5.
- [227] Skurnik Y, Shoenfeld Y. Health effects of cigarette smoking. *Clin Dermatol* 1998; 16(5): 545-56.
- [228] Das SK. Harmful health effects of cigarette smoking. *Mol Cell Biochem* 2003; 253(1-2): 159-65.
- [229] Wilson PW. Smoking, smoking cessation, and risk of cardiovascular disease. *Curr Treat Options Cardiovasc Med* 2006; 8(4): 276-81.
- [230] Gofman JW, Lindgren FT, Strisower B, De Lalla O, Glazier F, Tamplin A. Cigarette smoking, serum lipoproteins, and coronary heart disease. *Geriatrics* 1955; 10(8): 349-54.
- [231] Handa K, Tanaka H, Shindo M, Kono S, Sasaki J, Arakawa K. Relationship of cigarette smoking to blood pressure and serum lipids. *Atherosclerosis* 1990; 84(2-3): 189-93.
- [232] Imamura H, Tanaka K, Hirae C, *et al.* Relationship of cigarette smoking to blood pressure and serum lipids and lipoproteins in men. *Clin Exp Pharmacol Physiol* 1996; 23(5): 397-02.
- [233] Imamura H, Miyamoto N, Uchida K, Teshima K, Masuda Y, Kobata D. Cigarette smoking, blood pressure and serum lipids and lipoproteins in middle-aged women. *J Physiol Anthropol Appl Human Sci* 2001; 20(1): 1-6.
- [234] Whitehead TP, Robinson D, Allaway SL. The effects of cigarette smoking and alcohol consumption on blood lipids: a dose-related study on men. *Ann Clin Biochem* 1996; 33 (Pt 2): 99-106.
- [235] Moffatt RJ. Effects of cessation of smoking on serum lipids and high density lipoprotein-cholesterol. *Atherosclerosis* 1988; 74(1-2): 85-9.
- [236] Bernhard D, Wang XL. Smoking, oxidative stress and cardiovascular diseases--do anti-oxidative therapies fail? *Curr Med Chem* 2007; 14(16): 1703-12.
- [237] Tanriverdi H, Evrengul H, Kuru O, *et al.* Cigarette smoking induced oxidative stress may impair endothelial function and coro-

- nary blood flow in angiographically normal coronary arteries. *Circ J* 2006; 70(5): 593-9.
- [238] Lykkesfeldt J. Malondialdehyde as biomarker of oxidative damage to lipids caused by smoking. *Clin Chim Acta* 2007; 380(1-2): 50-8.
- [239] Albano SA, Santana-Sahagun E, Weisman MH. Cigarette smoking and rheumatoid arthritis. *Semin Arthritis Rheum* 2001; 31(3): 146-59.
- [240] Harrison BJ. Influence of cigarette smoking on disease outcome in rheumatoid arthritis. *Curr Opin Rheumatol* 2002; 14(2): 93-7.
- [241] Metsios GS, Stavropoulos-Kalinoglou A, Nevill AM, Douglas KM, Koutedakis Y, Kitis GD. Smoking significantly increases basal metabolic rate in patients with rheumatoid arthritis. *Ann Rheum Dis* 2007; 67(1): 70-3.
- [242] Stavropoulos-Kalinglou A, Metsios GS, Panoulas VF, *et al*. Cigarette smoking associates with body weight and muscle mass of patients with rheumatoid arthritis: a cross sectional, observational study. *Arthritis Res Ther* 2008; 10(3): R59.
- [243] Finckh A, Dehler S, Costenbader KH, Gabay C. Cigarette smoking and radiographic progression in rheumatoid arthritis. *Ann Rheum Dis* 2007; 66(8): 1066-71.
- [244] Sopori MM. Effects of cigarette smoke on the immune system. *Nat Rev Immunol* 2002; 2(5): 372-7.
- [245] Maclure M. Demonstration of deductive meta-analysis: ethanol intake and risk of myocardial infarction. *Epidemiol Rev* 1993; 15(2): 328-51.
- [246] Moore RD, Pearson TA. Moderate alcohol consumption and coronary artery disease. A review. *Medicine (Baltimore)* 1986; 65(4): 242-67.
- [247] Rimm EB, Williams P, Fosher K, Criqui M, Stampfer MJ. Moderate alcohol intake and lower risk of coronary heart disease: meta-analysis of effects on lipids and haemostatic factors. *BMJ* 1999; 319(7224):1523-8.
- [248] Handa K, Sasaki J, Saku K, Kono S, Arakawa K. Alcohol consumption, serum lipids and severity of angiographically determined coronary artery disease. *Am J Cardiol* 1990; 65(5): 287-9.
- [249] Kolovou GD, Salpea KD, Anagnostopoulou KK, Mikhailidis DP. Alcohol use, vascular disease, and lipid-lowering drugs. *J Pharmacol Exp Ther* 2006; 18(1): 1-7.
- [250] van de WA, van Golde PM, Kraaijenhagen RJ, den Borne PA, Bouma BN, Hart HC. Acute inhibitory effect of alcohol on fibrinolysis. *Eur J Clin Invest* 2001; 31(2): 164-70.
- [251] van Golde PM, Hart HC, Kraaijenhagen RJ, Bouma BN, van de WA. Regular alcohol intake and fibrinolysis. *Neth J Med* 2002; 60(7): 285-8.
- [252] Koppes LL, Twisk JW, Van Mechelen W, Snel J, Kemper HC. Cross-sectional and longitudinal relationships between alcohol consumption and lipids, blood pressure and body weight indices. *J Stud Alcohol* 2005; 66(6): 713-21.
- [253] Rossouw JE, Lai-Tung MT, Jooste PL, Weight MJ, Benade AJ. Alcohol intake in relation to lipids, lipoproteins and blood pressure. *S Afr Med J* 1992; 82(4): 246-50.
- [254] De Oliveira E Silva ER, Foster D, McGee HM, *et al*. Alcohol consumption raises HDL cholesterol levels by increasing the transport rate of apolipoproteins A-I and A-II. *Circulation* 2000; 102(19):2347-52.
- [255] Moore RD, Smith CR, Kwieterovich PO, Pearson TA. Effect of low-dose alcohol use vs abstinence on apolipoproteins A-I and B. *Am J Med* 1988; 84(5): 884-90.
- [256] Beulens JW, Sierksma A, van Tol A, *et al*. Moderate alcohol consumption increases cholesterol efflux mediated by ABCA1. *J Lipid Res* 2004; 45(9): 1716-23.
- [257] Hannuksela MY, Marcel YL, Kesaniemi YA, Savolainen MJ. Reduction in the concentration and activity of plasma cholesteryl ester transfer protein by alcohol. *J Lipid Res* 1997; 33: 737-44.
- [258] Nishiwaki M, Ishikawa T, Ito T, *et al*. Effects of alcohol on lipoprotein lipase, hepatic lipase, cholesteryl ester transfer protein, and lecithin:cholesterol acyltransferase in high-density lipoprotein cholesterol elevation. *Atherosclerosis* 1994; 111(1): 99-109.
- [259] Schafer C, Parlesak A, Eckoldt J, *et al*. Beyond HDL-cholesterol increase: phospholipid enrichment and shift from HDL3 to HDL2 in alcohol consumers. *J Lipid Res* 2007; 48(7): 1550-8.
- [260] Diehl AK, Fuller JH, Mattock MB, Salter AM, el Gohari R, Keen H. The relationship of high density lipoprotein subfractions to alcohol consumption, other lifestyle factors, and coronary heart disease. *Atherosclerosis* 1988; 69(2-3): 145-53.
- [261] Ostrander LD, Jr., Lamphiear DE, Block WD, Johnson BC, Ravenscroft C, Epstein FH. Relationship of serum lipid concentrations to alcohol consumption. *Arch Intern Med* 1974; 134(3): 451-6.
- [262] Frimpong NA, Lapp JA. Effects of moderate alcohol intake in fixed or variable amounts on concentration of serum lipids and liver enzymes in healthy young men. *Am J Clin Nutr* 1989; 50(5): 987-91.
- [263] Ayaori M, Ishikawa T, Yoshida H, *et al*. Beneficial effects of alcohol withdrawal on LDL particle size distribution and oxidative susceptibility in subjects with alcohol-induced hypertriglyceridemia. *Arterioscler Thromb Vasc Biol* 1997; 17(11): 2540-7.
- [264] Marcolongo R. The influence of genetic and environmental factors on the prevalence of rheumatoid arthritis and rheumatoid factor. *Reumatismo* 1971; 23(5): 241-51.
- [265] Klareskog L, Padyukov L, Lorentzen J, Alfredsson L. Mechanisms of disease: genetic susceptibility and environmental triggers in the development of rheumatoid arthritis. *Nat Clin Pract Rheumatol* 2006; 2(8): 425-33.
- [266] Hazes JM, Dijkmans BA, Vandenbroucke JP, de Vries RR, Cats A. Lifestyle and the risk of rheumatoid arthritis: cigarette smoking and alcohol consumption. *Ann Rheum Dis* 1990; 49(12): 980-2.
- [267] Voigt LF, Koepsell TD, Nelson JL, Dugowson CE, Daling JR. Smoking, obesity, alcohol consumption, and the risk of rheumatoid arthritis. *Epidemiology* 1994; 5(5): 525-32.
- [268] Iqbal R, Anand S, Ounpuu S, *et al*. Dietary patterns and the risk of acute myocardial infarction in 52 countries: results of the INTERHEART study. *Circulation* 2008; 118(19): 1913-4.
- [269] Huang CL, Sumpoi BE. Olive oil, the mediterranean diet, and cardiovascular health. *J Am Coll Surg* 2008; 207(3): 407-16.
- [270] Harris WS, Connor WE, Illingworth DR, Rothrock DW, Foster DM. The effects of fish oil on VLDL triglyceride kinetics in humans. *J Lipid Res* 1990; 31(9): 1549-58.
- [271] Tziomalos K, Athyros VG, Mikhailidis DP. Fish oils and vascular disease prevention: an update. *Curr Med Chem* 2007; 14(24): 2622-8.
- [272] Heber D, Yip I, Ashley JM, Elashoff DA, Elashoff RM, Go VL. Cholesterol-lowering effects of a proprietary Chinese red-yeast-rice dietary supplement. *Am J Clin Nutr* 1999; 69(2): 231-6.
- [273] Covas MI, Nyyssonen K, Poulsen HE, *et al*. The effects of polyphenols in olive oil on heart disease risk factors: a randomized trial. *Ann Intern Med* 2006; 145(5): 333-41.
- [274] Elkan AC, Sjoberg B, Kolsrud B, Ringertz B, Hafstrom I, Frostegard J. Gluten-free vegan diet induces decreased LDL and oxidized LDL levels and raised atheroprotective natural antibodies against phosphorylcholine in patients with rheumatoid arthritis: a randomized study. *Arthritis Res Ther* 2008; 10(2): R34.
- [275] Naranjo A, Sokka T, Descalzo MA, *et al*. Cardiovascular disease in patients with rheumatoid arthritis: results from the QUEST-RA study. *Arthritis Res Ther* 2008; 10(2): R30.
- [276] Toms TE, Panoulas VF, Stavropoulos-Kalinglou A, Kitis GD. Cardiovascular drugs in rheumatoid arthritis: killing two birds with one stone? *Immunol, Endocr Metab Agents - Med Chem* 2008; 8(3): 259-74.
- [277] McCarey DW, McInnes IB, Madhok R, *et al*. Trial of Atorvastatin in Rheumatoid Arthritis (TARA): double-blind, randomised placebo-controlled trial. *Lancet* 2004; 363(9426): 2015-21.
- [278] Jansen TL. Atorvastatin for chronic synovitis due to massive intra-articular cholesterol monohydrate deposition in long-standing rheumatoid arthritis. *Rheumatology (Oxford)* 2006; 45(12): 1577-8.
- [279] ten Cate R, Nibbeling PH, Bredius RG. Therapy-refractory systemic juvenile idiopathic arthritis successfully treated with statins. *Rheumatology (Oxford)* 2004; 43(7): 934-5.
- [280] Abud-Mendoza C, de la Fuente H, Cuevas-Orta E, Baranda L, Cruz-Rizo J, Gonzalez-Amaro R. Therapy with statins in patients with refractory rheumatic diseases: a preliminary study. *Lupus* 2003; 12(8): 607-11.
- [281] Okamoto H, Koizumi K, Kamitsuiji S, *et al*. Beneficial action of statins in patients with rheumatoid arthritis in a large observational cohort. *J Rheumatol* 2007; 34(5): 964-8.
- [282] Charles-Schoeman C, Khanna D, Furst DE, *et al*. Effects of high-dose atorvastatin on antiinflammatory properties of high density lipoprotein in patients with rheumatoid arthritis: a pilot study. *J Rheumatol* 2007; 34(7):1459-64.
- [283] Van Doornum S, McColl G, Wicks IP. Accelerated atherosclerosis: an extraarticular feature of rheumatoid arthritis? *Arthritis Rheum* 2002; 46(4): 862-73.

- [284] Hermann F, Forster A, Chenevard R, *et al.* Simvastatin improves endothelial function in patients with rheumatoid arthritis. *J Am Coll Cardiol* 2005; 45(3): 461-4.
- [285] Daoussis D, Panoulas VF, Antonopoulos I, *et al.* Cardiovascular risk factors and not disease activity, severity or therapy associate with renal dysfunction in patients with rheumatoid arthritis. *Ann Rheum Dis* 2009; doi: 10.1136/ard.2008.105049 [Epub a head of print].
- [286] Di Angelantonio E, Danesh J, Eiriksdottir G, Gudnason V. Renal function and risk of coronary heart disease in general populations: new prospective study and systematic review. *PLOS Med* 2007; 4(9): 1497-1507.
- [287] Efstratiadis G, Tziomalos K, Mikhailidis DP, Athyros VG, Hatzitolios A. Atherogenesis in renal patients: A model of vascular disease? *Curr Vasc Pharmacol* 2008; 6(2): 93-107.
- [288] Shepherd J, Kastelein JJ, Bittner V, *et al.* Effect of intensive lipid lowering with atorvastatin on renal function in patients with coronary heart disease: the Treating to New Targets (TNT) study. *Clin J Am Soc Nephrol* 2007; 2(6): 1131-9.
- [289] Athyros VG, Kakafika AI, papageorgiou AA, *et al.* Statin-induced increase in HDL-C and renal function in coronary heart disease patients. *Open Cardiovasc Med J* 2007; 1: 8-14.
- [290] Nurmohamed MT, Dijkmans BA. Dyslipidaemia, statins and rheumatoid arthritis. *Ann Rheum Dis* 2009; 68(4): 453-5.
- [291] Available from: www.dgoh.nhs.uk/tracera/ 2009.
- [292] EMPIRE Rheumatism Council: multi-centre controlled trial comparing cortisone acetate and acetyl salicylic acid in the long-term treatment of rheumatoid arthritis; results of three years' treatment. *Ann Rheum Dis* 1957; 16: 277-89.
- [293] Kirwan JR. The effect of glucocorticoids on joint destruction in rheumatoid arthritis. The Arthritis and Rheumatism Council Low-Dose Glucocorticoid Study Group. *N Engl J Med* 1995; 333(3): 142-6.
- [294] Bijlsma JW, Van Everdingen AA, Huisman M, De Nijs RN, Jacobs JW. Glucocorticoids in rheumatoid arthritis: effects on erosions and bone. *Ann N Y Acad Sci* 2002; 966: 82-90.
- [295] Maxwell SR, Moots RJ, Kendall MJ. Corticosteroids: do they damage the cardiovascular system? *Postgrad Med J* 1994; 70(830): 863-70.
- [296] Bagdade J, Casaretto A, Albers J. Effects of chronic uremia, hemodialysis, and renal transplantation on plasma lipids and lipoproteins in man. *J Lab Clin Med* 1976; 87(1): 38-48.
- [297] Cattran DC, Steiner G, Wilson DR, Fenton SA. Hyperlipidemia after renal transplantation: natural history and pathophysiology. *Ann Intern Med* 1979; 91(4): 554-9.
- [298] Zimmerman J, Fainaru M, Eisenberg S. The effects of prednisolone therapy on plasma lipoproteins and apolipoproteins: a prospective study. *Metabolism* 1984; 33: 521-6.
- [299] Ettinger WH, Klinefelter HF, Kwiterovich PO. Effect of short-term, low-dose corticosteroids on plasma lipoprotein lipids. *Atherosclerosis* 1987; 63(2-3): 167-72.
- [300] Choi HK, Seeger JD. Glucocorticoid use and serum lipid levels in US adults: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum* 2005; 53(4): 528-35.
- [301] Ettinger WH, Jr., Hazzard WR. Prednisone increases very low density lipoprotein and high density lipoprotein in healthy men. *Metabolism* 1988; 37(11): 1055-8.
- [302] Taskinen MR, Kuusi T, Yki-Jarvinen H, Nikkila EA. Short-term effects of prednisone on serum lipids and high density lipoprotein subfractions in normolipidemic healthy men. *J Clin Endocrinol Metab* 1988; 67(2): 291-9.
- [303] Dessein PH, Joffe BI, Stanwix AE, Christian BF, Veller M. Glucocorticoids and insulin sensitivity in rheumatoid arthritis. *J Rheum* 2004; 31(5): 867-74.
- [304] Boers M, Nurmohamed MT, Doelman CJ, *et al.* Influence of glucocorticoids and disease activity on total and high density lipoprotein cholesterol in patients with rheumatoid arthritis. *Ann Rheum Dis* 2003; 62(9): 842-5.
- [305] Wallace DJ, Metzger AL, Stecher VJ, Turnbull BA, Kern PA. Cholesterol-lowering effect of hydroxychloroquine in patients with rheumatic disease: reversal of deleterious effects of steroids on lipids. *Am J Med* 1990; 89(3): 322-6.
- [306] Dessein PH, Stanwix AE, Joffe BI. Cardiovascular risk in rheumatoid arthritis vs osteoarthritis: acute phase response related decreased insulin sensitivity and high-density lipoprotein cholesterol as well as clustering of metabolic syndrome features in rheumatoid arthritis. *Arthritis Res* 2002; 4(5): R5.
- [307] Beynen AC. Can chloroquine be of value in the treatment of hypercholesterolaemia. *Artery* 1986; 13: 340-51.
- [308] Munro R, Morrison E, McDonald AG, Hunter JA, Madhok R, Capell HA. Effect of disease modifying agents on the lipid profiles of patients with rheumatoid arthritis. *Ann Rheum Dis* 1997; 56(6): 374-7.
- [309] Rustan AC, Nossen JO, Tefre T, Drevon CA. Inhibition of very-low-density lipoprotein secretion by chloroquine, verapamil and monensin takes place in the Golgi complex. *Biochim Biophys Acta* 1987; 930(3): 311-9.
- [310] Chen HW, Leonard DA. Chloroquine inhibits cyclization of squalene oxide to lanosterol in mammalian cells. *J Biol Chem* 1984; 259(13): 8156-62.
- [311] Goldstein JL, Brunschede GY, Brown MS. Inhibition of proteolytic degradation of low density lipoprotein in human fibroblasts by chloroquine, concanavalin A, and Triton WR 1339. *J Biol Chem* 1975; 250(19): 7854-62.
- [312] Hermann B, Muller W. Die therapie der chronischen polyarthritis mit cyclosporin A: einem new en immunosuppressum (the therapy immunosuppressive). *Akt Rheumatol* 1979; 4: 173-86.
- [313] Farooqui AN, Ahmad SL, Mansuri FA. Efficacy of cyclosporin-A in refractory rheumatoid arthritis. *J Coll Physicians Surg Pak* 2004; 14(3): 139-41.
- [314] Gerards AH, Landewe RB, Prins AP, *et al.* Cyclosporin A monotherapy vs cyclosporin A and methotrexate combination therapy in patients with early rheumatoid arthritis: a double blind randomised placebo controlled trial. *Ann Rheum Dis* 2003; 62(4): 291-6.
- [315] Harris KP, Russell GI, Parvin SD, Veitch PS, Walls J. Alterations in lipid and carbohydrate metabolism attributable to cyclosporin A in renal transplant recipients. *Br Med J (Clin Res Ed)* 1986; 292(6512): 16.
- [316] Schorn TF, Kliem V, Bojanovski M, *et al.* Impact of long-term immunosuppression with cyclosporin A on serum lipids in stable renal transplant recipients. *Transpl Int* 1991; 4(2): 92-5.
- [317] Hilbrands LB, Demacker PN, Hoitsma AJ. Cyclosporin and serum lipids in renal transplant recipients. *Lancet* 1993; 341(8847): 765-6.
- [318] Le Goff W, Peng DQ, Settle M, Brubaker G, Morton RE, Smith JD. Cyclosporin A traps ABCA1 at the plasma membrane and inhibits ABCA1-mediated lipid efflux to apolipoprotein A-I. *Arterioscler Thromb Vasc Biol* 2004; 24(11): 2155-61.
- [319] Forestier F, De Seze S. History of the discovery of gold salts action in rheumatoid polyarthritis. *Rev Prat* 1992; 42(14): 1804-6.
- [320] Feldman M. Development of anti-tumour necrosis therapy for rheumatoid arthritis. *Nat Rev Immunol* 2007; 2(5): 364-71.
- [321] Kievit W, Adang EM, Fransen J, *et al.* The effectiveness and medication costs of three anti-TNF agents[alpha] in the treatment of rheumatoid arthritis from prospective clinical practice data. *Ann Rheum Dis* 2008; 67: 1229-34.
- [322] Glare J. Meta-analysis: serious adverse effects of anti-tumour necrosis therapies. *JAMA* 2006; 295: 2275-85.
- [323] Branten AJW. Risk of infectious complications during anti-tumour necrosis factor alpha therapy. *The Netherlands J Med* 2008; 66(2): 50-2.
- [324] Cauza E, Cauza K, Hanusch-Enserer U, Etemad M, Dunky A, Kostner K. Intravenous anti TNF-alpha antibody therapy leads to elevated triglyceride and reduced HDL-cholesterol levels in patients with rheumatoid and psoriatic arthritis. *Wien Klin Wochenschr* 2002; 114: 1004-7.
- [325] Irace C, Mancuso G, Fiaschi E, Madaia A, Sesti G, Gnasso A. Effect of anti TNF alpha therapy on arterial diameter and shear wall stress and HDL cholesterol. *Atherosclerosis* 2004; 177(1): 113-8.
- [326] Oguz FM, Oguz A, Uzunlulu M. The effect of infliximab treatment on insulin resistance in patients with rheumatoid arthritis. *Acta Clin Belg* 2007; 62(4): 218-22.
- [327] Nishida K, Okada Y, Nawata M, Saito K, Tanaka Y. Induction of hyperadiponectinemia following long-term treatment of patients with rheumatoid arthritis with infliximab (IFX), an anti-TNF-alpha antibody. *Endocr J* 2008; 55(1): 213-6.
- [328] Tam LS, Tomlinson B, Chu TT, Li TK, Li EK. Impact of TNF inhibition on insulin resistance and lipids levels in patients with rheumatoid arthritis. *Clin Rheumatol* 2007; 26(9): 1495-8.
- [329] Vis M, Nurmohamed MT, Wolbink G, *et al.* Short term effects of infliximab on the lipid profile in patients with rheumatoid arthritis. *J Rheumatol* 2005; 32(2): 252-5.

- [330] Soubrier M, Jouanel P, Mathieu S, *et al.* Effects of anti-tumor necrosis factor therapy on lipid profile in patients with rheumatoid arthritis. *Joint Bone Spine* 2008; 75(1): 22-4.
- [331] Spanakis E, Sidiropoulos P, Papadakis J, *et al.* Modest but sustained increase of serum high density lipoprotein cholesterol levels in patients with inflammatory arthritides treated with infliximab. *J Rheumatol* 2006; 33(12): 2440-6.
- [332] Saiki O, Takao R, Naruse Y, Kuhara M, Imai S, Uda H. Infliximab but not methotrexate induces extra high levels of VLDL-triglyceride in patients with rheumatoid arthritis. *J Rheumatol* 2007; 34(10): 1997-2004.
- [333] Cohen SB, Emery P, Greenwald MW, *et al.* Rituximab for rheumatoid arthritis refractory to anti-tumor necrosis factor therapy: results of a multicenter, randomized, double-blind, placebo-controlled, phase III trial evaluating primary efficacy and safety at twenty-four weeks. *Arthritis Rheum* 2006; 54(9): 2793-806.
- [334] Keystone E, Emery P, Peterfy CG, *et al.* Rituximab inhibits structural joint damage in patients with rheumatoid arthritis with an inadequate response to tumour necrosis factor inhibitor therapies. *Ann Rheum Dis* 2009; 68(2): 216-21.
- [335] Gonzalez-Juanatey C, Llorca J, Vazquez-Rodriguez TR, Diaz-Varela N, Garcia-Quiroga H, Gonzalez-Gay MA. Short-term improvement of endothelial function in rituximab-treated rheumatoid arthritis patients refractory to tumor necrosis factor alpha blocker therapy. *Arthritis Rheum* 2008; 59(12): 1821-4.
- [336] Sandoo A, Toms T, Zanten JV, Carroll D, Kitas GD. Short-term effects of rituximab on flow-mediated dilatation may be mediated by intravenous glucocorticoids: comment on the article by Gonzalez-Juanatey, *et al.* *Arthritis Rheum* 2009; 61(6): 854-5.
- [337] Nishimoto N, Hashimoto J, Miyasaka N, *et al.* Study of active controlled monotherapy used for rheumatoid arthritis, an IL-6 inhibitor (SAMURAI): evidence of clinical and radiographic benefit from an x ray reader-blinded randomised controlled trial of tocilizumab. *Ann Rheum Dis* 2007; 66(9): 1162-7.
- [338] Maini RN, Taylor PC, Szechinski J, *et al.* Double-blind randomized controlled clinical trial of the interleukin-6 receptor antagonist, tocilizumab, in European patients with rheumatoid arthritis who had an incomplete response to methotrexate. *Arthritis Rheum* 2006; 54(9): 2817-29.
- [339] Bertolini S, Pisciotto L, Seri M, *et al.* A point mutation in ABCA1 gene in a patient with severe premature coronary heart disease and mild clinical phenotype of Tangier disease. *Atherosclerosis* 2001; 154(3): 599-605.
- [340] Frikke-Schmidt R, Nordestgaard BG, Schnohr P, Steffensen R, Tybjaerg-Hansen A. Mutation in ABCA1 predicted risk of ischemic heart disease in the Copenhagen City Heart Study Population. *J Am Coll Cardiol* 2005; 46(8): 1516-20.
- [341] Frikke-Schmidt R, Nordestgaard BG, Jensen GB, Steffensen R, Tybjaerg-Hansen A. Genetic variation in ABCA1 predicts ischemic heart disease in the general population. *Arterioscler Thromb Vasc Biol* 2008; 28(1): 180-6.
- [342] Iatan I, Alrasadi K, Ruel I, Alwaili K, Genest J. Effect of ABCA1 mutations on risk for myocardial infarction. *Curr Atheroscler Rep* 2008; 10(5): 413-26.
- [343] Bertolini S, Pisciotto L, Di Scala L, *et al.* Genetic polymorphisms affecting the phenotypic expression of familial hypercholesterolemia. *Atherosclerosis* 2004; 174(1): 57-65.
- [344] Evans D, Beil FU. The association of the R219K polymorphism in the ATP-binding cassette transporter 1 (ABCA1) gene with coronary heart disease and hyperlipidaemia. *J Mol Med* 2003; 81(4): 264-70.
- [345] Srinivasan SR, Li S, Chen W, Boerwinkle E, Berenson GS. R219K polymorphism of the ABCA1 gene and its modulation of the variations in serum high-density lipoprotein cholesterol and triglycerides related to age and adiposity in white vs black young adults. The Bogalusa heart study. *Metabolism* 2003; 52(7): 930-4.
- [346] Andrikovics H, Pongracz E, Kalina E, *et al.* Decreased frequencies of ABCA1 polymorphisms R219K and V771M in Hungarian patients with cerebrovascular and cardiovascular diseases. *Cerebrovasc Dis* 2006; 21(4): 254-9.
- [347] Nebel A, Croucher PJ, El Mokhtari NE, Flachsbarth F, Schreiber S. Common coding polymorphisms in the ABCA1 gene and risk of early-onset coronary heart disease in northern Germany. *Atherosclerosis* 2007; 193(2): 458-60.
- [348] Frikke-Schmidt R, Nordestgaard BG, Jensen GB, Tybjaerg-Hansen A. Genetic variation in ABC transporter A1 contributes to HDL cholesterol in the general population. *J Clin Invest* 2004; 114(9): 1343-53.
- [349] Kakko S, Kelloniemi J, von Rohr P, *et al.* ATP-binding cassette transporter A1 locus is not a major determinant of HDL-C levels in a population at high risk for coronary heart disease. *Atherosclerosis* 2003; 166(2): 285-90.
- [350] Brousseau ME, Bodzioch M, Schaefer EJ, *et al.* Common variants in the gene encoding ATP-binding cassette transporter 1 in men with low HDL cholesterol levels and coronary heart disease. *Atherosclerosis* 2001; 154(3): 607-11.
- [351] Boekholdt SM, Sacks FM, Jukema JW, *et al.* Cholesterol ester transfer protein TaqIB variant, high density lipoprotein cholesterol levels, cardiovascular risk and efficacy of Pravastatin treatment. Individual patient meta-analysis of 13677 subjects. *Circulation* 2005; 111: 278-87.
- [352] Ordovas JM, Cupples LA, Corella D, *et al.* Association of cholesterol ester transfer protein-TaqIB polymorphism with variations in lipoprotein subclasses and coronary heart disease risk: the Framingham study. *Arterioscler Thromb Vasc Biol* 2000; 20: 1323-9.
- [353] Brousseau ME, O'Connor JJ Jr, Ordovas JM, *et al.* Cholesterol ester transfer protein TaqIB2B2 genotype is associated with higher HDL cholesterol and lowers risk of coronary heart disease end points in men with HDL deficiency: Veterans affairs HDL cholesterol intervention trial. *Arterioscler Thromb Vasc Biol* 2002; 22(1148): 1154.
- [354] Thompson A, Angelantonio ED, Sarwar N, *et al.* Association of cholesteryl ester transfer protein genotypes with CETP mass and activity, lipid levels, and coronary risk. *JAMA* 2008; 299(23): 2777-88.
- [355] Carlquist JF, Muhlestein JB, Horne BD, *et al.* The cholesteryl ester transfer protein TaqIB gene polymorphism predicts clinical benefit of statin therapy in patients with significant coronary artery disease. *Am Heart J* 2003; 146(6): 1007-14.
- [356] Shelbourne F, Hanks J, Meyers W, Quarfordt S. Effect of apoproteins on hepatic uptake of triglyceride emulsion in the rat. *J Clin Invest* 1980; 65: 652-8.
- [357] Goldberg IJ, Le NA, Paterniti JR, Ginsberg HN, Lindgren FT, Brown WV. Lipoprotein metabolism during acute inhibition of hepatic triglyceride lipase in the cynomolgus monkey. *J Clin Invest* 1982; 70: 1184-92.
- [358] Eichner JE, Dunn ST, Perveen G, Thompson DM, Stewart KE, Stroehla BC. Apolipoprotein E polymorphism and cardiovascular disease: A HuGE review. *Am J Epidemiol* 2002; 155(6): 487-95.
- [359] Siest G, Pillot T, Regis-Bailly A, *et al.* Apolipoprotein E: an important gene and protein to follow in laboratory medicine. *Clin Chem* 1995; 41(8 Pt 1): 1068-86.
- [360] Weintraub MS, Eisenberg S, Breslow JL. Dietary fat clearance in normal subjects is regulated by genetic variation in apolipoprotein E. *J Clin Invest* 1987; 80(6): 1571-7.
- [361] Utermann G, Hees M, Steinmetz A. Polymorphism of apolipoprotein E and occurrence of dysbetalipoproteinemia in man. *Nature* 1977; 269(604): 607.
- [362] Davignon J, Gregg RE, Sing CF. Apolipoprotein E polymorphism and atherosclerosis. *Arteriosclerosis* 1988; 8(1): 1-21.
- [363] Utermann G. Apolipoprotein E polymorphism in health and disease. *Am Heart J* 1987; 113(433): 440.
- [364] Cattin L, Fiscaro M, Tonizzo M, *et al.* Polymorphism of the apolipoprotein E gene and early carotid atherosclerosis defined by ultrasonography in asymptomatic adults. *Arterioscler Thromb Vasc Biol* 1997; 17(1): 91-4.
- [365] Hooper AJ, Crawford GM, Brisbane JM, *et al.* Familial lipoprotein lipase deficiency caused by known (G188E) and novel (W394X) LPL gene mutations. *Ann Clin Biochem* 2008; 45: 102-5.
- [366] Sagoo GS, Tatt I, Salanti G, *et al.* Seven lipoprotein lipase gene polymorphisms, lipid fractions and coronary disease: a HuGE association review and meta-analysis. *Am J Epidemiol* 2008; 168(11): 1233-46.
- [367] Burdon KP, Langefeld CD, Beck SR, *et al.* Association of genes of lipid metabolism with measures of subclinical cardiovascular disease in the Diabetes Heart Study. *J Med Genet* 2005; 42(9): 720-4.
- [368] Morabia A, Cayanis E, Costanza MC, *et al.* Association between lipoprotein lipase (LPL) gene and blood lipids: a common variant for a common trait? *Genet Epidemiol* 2003; 24(4): 309-21.
- [369] Saha N, Low PS, Kamboh MI. Association of two polymorphisms in the lipoprotein lipase gene with coronary heart disease. *Am J Hum Genet* 1998; (Suppl. 1): A209.

- [370] Talmud PJ, Hawe E, Martin S, *et al.* Relative contribution of variation within the APOC3/A4/A5 gene cluster in determining plasma triglycerides. *Hum Mol Genet* 2002; 4:3039-46.
- [371] Wang QF, Liu X, O'Connell J, *et al.* Haplotypes in the ApoA1-C3-A4-A5 gene cluster affect plasma lipids in both humans and baboons. *Hum Mol Genet* 2004; 13(10):1049-56.
- [372] Wang CS, McConathy WJ, Kloer HU, Alaupovic P. Modulation of lipoprotein lipase activity by apolipoproteins. Effect of apolipoprotein C-III. *J Clin Invest* 1985; 75: 384-90.
- [373] Herron KL, Lofgren IE, Adiconis X, Ordovas JM, Fernandez ML. Associations between plasma lipid parameters and ApoC3 and ApoA4 genotypes in a healthy population are independent of dietary cholesterol intake. *Atherosclerosis* 2006; 184(1): 113-20.
- [374] Shanker J, Perumal G, Rao VS, *et al.* Genetic studies on the ApoA1-C3-A5 gene cluster in Asian Indians with premature coronary disease. *Lipids Health Dis* 2008; 19(7): 33.
- [375] Vaessen SFC, Schaap FG, Kuivenhoven JA, *et al.* Apolipoprotein A-V, triglycerides and risk of coronary disease: the prospective Epic-Norfolk population study. *J Lipid Res* 2006; 47: 2064-70.
- [376] Yu Y, Xue L, Zhao CY. Study of the polymorphism in the apolipoprotein A5 gene in patients with premature coronary heart disease. *Beijing Da Xue Xue Bao* 2007; 39: 576-80.
- [377] Pennacchio LA, Rubin EM. Apolipoprotein A5, a newly identified gene that affects plasma triglyceride levels in humans and mice. *Arterioscler Thromb Vasc Biol* 2003; 23: 529-34.
- [378] Szalai C, Keszei M, Duba J, *et al.* Polymorphism in the promoter region of the apolipoprotein A5 gene is associated with an increased susceptibility for coronary artery disease. *Atherosclerosis* 2004; 173: 109-14.
- [379] Fiegenbaum M, de Andrade FM, Hutz MH. Association between plasma lipid parameters and APOC3 genotypes in Brazilian subjects: effect of gender, smoking and APOE genotypes. *Clin Chim Acta* 2007; 380(1-2): 175-81.
- [380] van der Net JB, Janssens AC, Defesche JC, Kastelein JJ, Sijbrands EJ, Steyerberg EW. Usefulness of genetic polymorphisms and conventional risk factors to predict coronary heart disease in patients with familial hypercholesterolemia. *Am J Cardiol* 2009; 103(3): 375-80.
- [381] Martin-Campos JM, Rico N, Bonet R, Mayoral C, Ordonez-Llanos J, Blanco-Vaca F. Apolipoprotein A5 S19W may play a role in dysbetalipoproteinemia in patients with the Apo E2/E2 genotype. *Clin Chem* 2006; 52(10): 1974-5.
- [382] Martinelli N, Trabetti E, Bassi A, *et al.* The -1131 T>C and S19W APOA5 gene polymorphisms are associated with high levels of triglycerides and apolipoprotein C-III, but not with coronary artery disease: an angiographic study. *Atherosclerosis* 2007; 191(2): 409-17.