

An update on the cytokine network in rheumatoid arthritis

Pierre Miossec

Purpose of review

To update the knowledge accumulated on the contribution of cytokines to rheumatoid arthritis and related animal models. Publications from the end of 2002 and 2003 period were analyzed for a selection.

Recent findings

A better understanding of the clinical results with tumor necrosis factor- α inhibitors has come from studies in treated patients. The expected effect of infliximab on the apoptosis of cells expressing tumor necrosis factor- α was not observed in synovium biopsy specimens. The mode of action of tumor necrosis factor- α on bone destruction has been clarified in gene-defective mice. Tumor necrosis factor- α acts through osteoclasts—an effect that is inhibited with osteoprotegerin. New interleukin-1 inhibitors with a potential for increased efficacy, such as interleukin-1 trap, have been manufactured and are now being tested in rheumatoid arthritis. The list of cytokines of interest for therapeutic intervention has been growing rapidly. The results with animal models have provided clues to control arthritis with natural interleukin-18 inhibitors, such as interleukin-18 BP. Additional results have been accumulated that indicate the contribution of T cell subsets in inflammation and destruction through the production of interleukin-17. Synergistic interactions with other cytokines are critical in the interleukin-17 tuning effects. Macrophage inhibitory factor was described many years ago. Its comeback is based on properties of synovocyte activation and proliferation.

Summary

Such findings are critical for a better understanding of response heterogeneity in patients treated with the cytokine inhibitors now on the market. New therapeutic approaches are being planned from these results.

Keywords

rheumatoid arthritis, inflammation, destruction, cytokines, treatment

Current Opinion in Rheumatology 2004, 16:218–222

Abbreviations

IFN	interferon
IL	interleukin
MIF	migration inhibitory factor
MMP	matrix metalloproteinases
RA	rheumatoid arthritis
TLR	toll-like receptor
TNF	tumor necrosis factor

© 2004 Lippincott Williams & Wilkins
1040–8711

Introduction

In recent years, a large body of results has clearly indicated that cytokines contribute to joint diseases [1••]. The role of cytokines in chronic inflammation has been clarified with clinical results obtained with inhibitors. From early results obtained in rheumatoid arthritis (RA), more recent studies have shown that the same cytokines are also involved in other types of arthritis, such as ankylosing spondylitis, and in many inflammatory diseases. The nonspecific effect of cytokines is indicated because the same findings have been obtained in diseases with different anatomic localizations.

However, the situation is far from being completely understood. Additional mechanisms have been identified to explain the efficacy and failures as well as the side effects. Some 20 years ago, the picture could be simplified with the identification of only two major cytokines, tumor necrosis factor (TNF)- α and interleukin (IL)-1, in the understanding of chronic inflammation. Today the list of cytokines, chemokines, and other factors reaches more than 100, making the choice of which one to target more difficult. At the same time, it remains to be understood how the inhibition of a single cytokine can still be clinically effective.

Here we will review the literature on the cytokine network in RA and related models published during the last part of 2002 and 2003. We will focus first on novel understandings obtained from clinical results with TNF- α and IL-1 inhibitors. Then we will consider cytokines, which could also become treatment targets.

Curr Opin Rheumatol 16:218–222. © 2004 Lippincott Williams & Wilkins.

Department of Immunology and Rheumatology, Hôpital Edouard Herriot, Lyon, France

Correspondence to Pierre Miossec, MD, PhD, Department of Immunology and Rheumatology, Hôpital Edouard Herriot, 69437 Lyon, Cedex 03, France
Tel: 334 72 11 74 87; Fax: 334 72 11 74 29; e-mail: miossec@univ-lyon1.fr

Effects of tumor necrosis factor- α , interleukin-1 inhibitors, and interleukin-6 inhibitors

Clinical results with tumor necrosis factor- α inhibitors

The mode of action of infliximab, a chimeric monoclonal antibody, could differ from that of etanercept, a soluble p75 TNF- α receptor. It was suggested that monoclonal antibodies could recognize membrane-bound TNF- α , leading to the destruction of the TNF- α -expressing cells through classic pathway complement activation. This would not be the case with etanercept, able to bind only soluble TNF- α . However, when synovial biopsy specimens were compared before and at day 28 of infliximab treatment, an increase in the number of apoptotic cells defined with the TUNEL assay could not be detected. Examination showed a reduced number of cells in the biopsy specimens from treated patients, indicating an effect on cell migration [2•].

Various cytokines have been shown to amplify the production of TNF- α . Among these, IL-7 is a cytokine produced by synoviocytes, which was found to stimulate TNF- α production by synovial fluid mononuclear cells and very potently that of IFN- γ and TNF- α by synovial fluid CD4+ T cells [3•].

The mode of action of TNF- α on joint destruction has been clarified using transgenic mice that express human TNF- α . In this model, mice have a severe and destructive arthritis. They were crossed with c-fos-deficient mice, which completely lack osteoclasts and thus have osteopetrosis. In the resulting mutant mice, a TNF- α -dependent arthritis developed in the absence of osteoclasts. However, despite the presence of severe inflammatory changes, these mice were fully protected against bone destruction. These results indicate that TNF- α -dependent bone erosion is mediated by osteoclasts. Therefore, in addition to the use of anti-inflammatory therapies targeting cytokines, osteoclast inhibition could be beneficial for RA treatment [4••]. Indeed, treatment of human TNF- α transgenic mice with osteoprotegerin completely blocked TNF- α -mediated bone loss by increasing bone mineral density and bone volume. Moreover, bone formation was dramatically increased [5•].

Improvement of interleukin-1 inhibitors

Whereas the result with TNF- α inhibitors are in line with those obtained in animal models, the results with IL-1 inhibition have showed rather inferior effects. It was suggested that the lower effects of IL-1Ra in the clinic could be explained by its short half-life *in vivo* and by the need for a permanent presence of IL-1Ra in the circulation at high concentrations. Among other ways to inhibit IL-1 with endogenous inhibitors, soluble IL-1 receptors could be of interest. This applies not so much to the type I receptor as to the type II IL-1 receptor. The

type I membrane-bound receptor is able to transduce the IL-1 driven signal, whereas the type II does not do so but is released as a soluble form, acting as a natural inhibitor. To lead to signal activation and transduction, the type I membrane receptor needs to be associated with the IL-1 receptor accessory protein. Such an association is not observed with IL-1Ra, explaining its lack of agonistic effect. On the basis of these findings, the administration of the sIL-1 receptor accessory protein showed a positive protective effect on joint destruction in the collagen arthritis model. To go further, a combination of both the soluble type I IL-1R and the sIL-1AcP led to the production of an IL-1 trap [6•]. Such a complex showed an increased affinity for IL-1. Its use in the collagen arthritis model led to a protective effect [7••]. It remains to be seen whether this complex will show the same efficacy in RA trials.

The IL-1Ra, as used as a treatment today, is a member of a growing family of endogenous IL-1 inhibitors. In particular, various intracellular forms of IL-1Ra have been isolated. These variants can still have a regulatory effect directly inside the cell, without interaction at the level of the membrane IL-1 receptor. Such a regulatory effect was indicated when mice transgenic for the intracellular IL-1Ra type I were protected from collagen-induced arthritis [8•].

In the spontaneous inflammatory arthritis of K/BxN T cell receptor transgenic mice, the effector phase of the disease is induced by anti-glucose 6 phosphate isomerase antibodies. It should be mentioned that the applications of these new concepts to human RA might be difficult because the same pathogenic antibodies in the mouse were not observed in various series of human RA [9•]. The roles of TNF- α and IL-1 have been evaluated by transferring arthrogenic serum into a panel of genetically deficient mice [10•]. To get the severe destructive arthritis, IL-1 proved to be absolutely necessary. TNF- α was also required, but less than IL-1, because disease developed in some of the TNF- α -deficient mice. However, there was no evidence of a role for TNF- α in bone destruction.

The administration of a toll-like receptor (TLR)-4 ligand, lipopolysaccharide, combined with arthrogenic serum (defined above) in IL-1 receptor-deficient mice resulted in acute joint swelling, but not in MyD88-deficient mice. Benign arthritis was observed in TLR-4 mutant mice [11•]. These results indicated the contribution of innate immunity via TLR-4. Because MyD88 is a common pathway for IL-1 and TLR-4 signaling, TLR-4 pathway may bypass the direct contribution of IL-1 in chronic joint inflammation.

Interleukin-1 is a critical factor for the activation of matrix metalloproteinases (MMP) in cartilage. Bcl-3 is a

member of the NF- κ B family and a known regulator of NF- κ B. It appears to be critical in the IL-1-induced MMP-1 transcription in chondrocytes, through an interaction with NF- κ B [12]. ESE-1 is another transcription factor expressed in RA synovium. It is specifically induced in synovial fibroblasts, chondrocytes, osteoblasts, and monocytes/macrophages by IL-1 β , TNF- α , or LPS. This induction is related to the translocation of the NF- κ B family members p50 and p65 to the nucleus and the transactivation of the ESE-1 promoter via a high-affinity NF- κ B binding site [13].

Synovium in RA is associated with neoangiogenesis, as observed in tumor growth and invasiveness. Using various IL-1 deficient mice, it was observed that IL-1 β and, to a lesser extent, IL-1 α , were required for *in vivo* angiogenesis and invasiveness of different tumor cells. Conversely, these effects were reduced with IL-1Ra [14••].

Interleukin-6 inhibition

More recently, IL-6 has been the target of RA treatment with an anti-IL-6 receptor antibody. Such treatment reduced VEGF production in RA patient serum. IL-6 acted synergistically with IL-1 β or TNF- α to induce VEGF production by RA synoviocytes. Such synergy with IL-1 β or TNF- α may be the mechanism by which IL-6 blockade effectively suppresses VEGF production in synovial fibroblasts [15].

More cytokines as therapeutic targets

Interleukin-17

Interleukin-17 is a T cell cytokine produced in the context of RA synovium. Although its effects alone are rather limited, IL-17 appears to be a potent actor in joint inflammation and destruction through additive and synergistic interactions with other proinflammatory cytokines [16•]. Such enhancing effects have been demonstrated mainly for TNF- α and IL-1. In addition, IL-17 increases IL-1 and TNF- α production by monocytes. Conversely, IL-17 production was found to be increased by monocyte products such as the new IL-23 [17•]. Accordingly, T cells through the production of IL-17 may regulate the proinflammatory status.

The production of IL-17 from activated T cells is required for the spontaneous development of destructive arthritis in mice deficient in IL-1Ra [18•]. IL-17 is part of the T cell-derived cytokines other than RANKL such as granulocyte-macrophage colony-stimulating factor and IFN- γ , which have powerful regulatory effects on osteoclastogenesis [19••].

Previously, IL-17 was shown to be a nitric oxide-producing cytokine. Such an effect leads to an inhibitory effect on proteoglycan synthesis [20]. In synoviocytes, IL-17

showed additive effects with IL-1 β and TNF- α on the expression of TSG-6, a hyaluronan-binding protein [21].

T cells regulate the expression of MMPs in human osteoblasts, including MMP-13. Under conditions of chronic inflammation, multiple T cell cytokines, including IL-17, synergize to induce high levels of MMP-13 via a mechanism that is dependent on activated p38 MAP kinase and is suppressed by activated ERK-1/2 [22].

Interleukin-18

IL-18 is one of the major cytokines, which induce a Th1 profile. Its effect is mediated by IL-12. Stimulation with IL-12 induces the β chain of the IL-18 receptor. This makes functional the IL-18 receptor, which is composed of the constitutive α chain and the regulated β chain. On mononuclear synovium cells, both receptors are expressed. Synoviocytes do not respond to IL-18 because the β chain is not expressed and cannot be induced by IL-12 [23•]. In RA synovium, expression of IL-18 has been associated with an increased expression of TNF α and IL-1, indicating the link between IL-18 and inflammation [24•]. IL-18 was found to be a chemoattractant for T cells and RA synovial CD4+ T cells responded to IL-18 by adopting a polarized morphology. Injection of IL-18 into mouse footpad led to the local accumulation of inflammatory cells [25•].

Such results suggest the control of IL-18 as treatment of joint inflammation. The high levels of IL-18 expression in RA joints are in contrast with the reduced IL-18 expression in RA peripheral blood mononuclear cells. Such low levels were increased by treatment with steroids [26].

The action of IL-18 is regulated by the endogenous release of IL-18BP, which binds to IL-18 with a high affinity as a decoy receptor. Intra-articular over-expression of IL-18BP significantly reduced incidence of collagen-induced arthritis in treated knee joints. Affected knee joints of IL-18BP-treated mice showed less severe arthritis, characterized by a reduction of inflammation and destruction of bone and cartilage. Local intra-articular IL-18BP treatment in both knees provided additional protection against CIA incidence and severity in distal paws, suggesting a systemic effect [27••]. Treatment of human synoviocytes with IFN γ increased IL-18BP, indicating a negative feedback loop via IL-18BP, which may limit IL-18 biologic activity in arthritis [28].

New interleukin-12 family members

IL-12 has been characterized as a major cytokine leading to the Th1 pathway at least in part through the production of IFN γ by T cells and NK cells. IL-18 acts in synergy with IL-12 to further increase such effects. Additional members of the IL-12 family can also contribute. IL-12 is composed of two subunits p35 and p40 whereas

the new IL-23 is made of the same p40 subunit combined with a new p19 subunit. It appears that IL-23 is in fact the key cytokine for the IL-12 effect. This was first shown for animal models of brain inflammation [29••]. Specific IL-23 inhibitors were able to reduce brain inflammation first thought to be related to an IL-12 effect. This case of mistaken identity appears to result from the use of inhibitors acting both on IL-12 and IL-23. As indicated above, IL-23 has been characterized as a potent inducer of IL-17 production by CD4 T cells [17•].

Macrophage migration inhibitory factor

Macrophage migration inhibitory factor (MIF) is a pro-inflammatory cytokine that may be of critical importance in the pathogenesis of RA [30••]. In particular, MIF may regulate RA synovial hyperplasia by acting on synovio-cytes directly and via an involvement in the effects of IL-1 β and TNF α . In addition, the effects of MIF on synovio-cyte activation are independent of NF- κ B, and dependent on ERK MAP kinase [31]. In MIF knockout mice, synovio-cyte proliferation was decreased. The decrease in the severity of antigen-induced arthritis was associated with an increase in p53 expression and apoptosis in synovium without an obvious effect on proliferation. These results indicate a role for MIF in the regulation of p53 expression and p53-mediated events in the inflamed synovium [32]. These data suggest an important therapeutic potential for MIF antagonism in RA.

Such a link with joint inflammation has been linked to MIF gene heterogeneity. In patients with systemic-onset juvenile idiopathic arthritis, the -173 single-nucleotide G-to-C polymorphism of the MIF gene was associated with higher levels of MIF in serum and synovial fluid and moreover with a poor outcome [33].

Conclusion

Inhibition of cytokines has led to an important step in the treatment of chronic inflammation. Their nonspecific effects, once considered to be a limitation, allowed the use of the same inhibitors in different diseases. Today their mode of action, as well as that of their inhibitors, remains to be clarified to explain heterogeneity in patient response and tolerance. The identification of new therapeutic targets research may allow better understanding and treatment of these complex diseases.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- Of special interest
- Of outstanding interest

- 1 Firestein GS: Evolving concepts of rheumatoid arthritis. *Nature* 2003, **423**:356–361.
- A very good review on the pathogenesis of rheumatoid arthritis with a personal historical perspective. In particular, the various theories used over the years are described with their pitfalls.

- 2 Smeets TJ, Kraan MC, van Loon ME, et al.: Tumor necrosis factor alpha blockade reduces the synovial cell infiltrate early after initiation of treatment, but apparently not by induction of apoptosis in synovial tissue. *Arthritis Rheum* 2003, **48**:2155–2162.
- An *in vivo* demonstration indicating that infliximab may not act through synovial cell apoptosis. Such mechanism was proposed to differentiate the mode of action of monoclonal antibodies from that of soluble receptors.
- 3 van Roon JA, Glaudemans KA, Bijlsma JW, et al.: Interleukin 7 stimulates tumour necrosis factor alpha and Th1 cytokine production in joints of patients with rheumatoid arthritis. *Ann Rheum Dis* 2003, **62**:113–119.
- A study indicating the proinflammatory contribution of IL-7.
- 4 Redlich K, Hayer S, Ricci R, et al.: Osteoclasts are essential for TNF-alpha-mediated joint destruction. *J Clin Invest* 2002, **110**:1419–1427.
- The demonstration of the critical role of osteoclasts in TNF-induced joint destruction using c-fos-deficient mice. The demonstration was made with c-fos-deficient mice, which lack osteoclasts. These mice still had arthritis but without bone destruction.
- 5 Schett G, Redlich K, Hayer S, et al.: Osteoprotegerin protects against generalized bone loss in tumor necrosis factor-transgenic mice. *Arthritis Rheum* 2003, **48**:2042–2051.
- A study on the effect of osteoprotegerin on joint-associated bone destruction.
- 6 Dinarello CA: Setting the cytokine trap for autoimmunity. *Nat Med* 2003, **9**:20–22.
- An overview on the new tools for cytokine inhibition.
- 7 Economides AN, Carpenter LR, Rudge JS, et al.: Cytokine traps: multi-component, high-affinity blockers of cytokine action. *Nat Med* 2003, **9**:47–52.
- This is the first report showing the proof of principle indicating the interest of the cytokine trap concept for treatment. In particular, IL-1 trap showed a powerful effect in the collagen arthritis model. Studies in human RA are in progress and should clarify the contribution of IL-1 in RA.
- 8 Palmer G, Talabot-Ayer D, Szalay-Quinodoz L, et al.: Mice transgenic for intracellular interleukin-1 receptor antagonist type 1 are protected from collagen-induced arthritis. *Eur J Immunol* 2003, **33**:434–440.
- A mouse study showing the biologic function of intracellular forms of IL-1Ra.
- 9 Matsumoto I, Lee DM, Goldbach-Mansky R, et al.: Low prevalence of antibodies to glucose-6-phosphate isomerase in patients with rheumatoid arthritis and a spectrum of other chronic autoimmune disorders. *Arthritis Rheum* 2003, **48**:944–954.
- A clinical study showing the lack of association between RA and antibodies to glucose-6-phosphate isomerase. The direct role of such antibodies as observed in the KRN model is not similarly observed in human RA.
- 10 Ji H, Pettit A, Ohmura K, et al.: Critical roles for interleukin 1 and tumor necrosis factor alpha in antibody-induced arthritis. *J Exp Med* 2002, **196**:77–85.
- In the KRN model of antibody-induced arthritis, the respective roles of TNF and IL-1 were shown using various deficient mice. The contribution of IL-1 was more critical than that of TNF.
- 11 Choe JY, Crain B, Wu SR, et al.: Interleukin 1 receptor dependence of serum transferred arthritis can be circumvented by toll-like receptor 4 signaling. *J Exp Med* 2003, **197**:537–542.
- This study provides arguments for the contribution of innate immunity in arthritis. Activation through TLR-4 was able to act as with IL-1. This is explained by the use of common transduction pathways for TLR-4 and IL-1.
- 12 Elliott SF, Coon CI, Hays E, et al.: Bcl-3 is an interleukin-1-responsive gene in chondrocytes and synovial fibroblasts that activates transcription of the matrix metalloproteinase 1 gene. *Arthritis Rheum* 2002, **46**:3230–3239.
- 13 Grall F, Gu X, Tan L, et al.: Responses to the proinflammatory cytokines interleukin-1 and tumor necrosis factor alpha in cells derived from rheumatoid synovium and other joint tissues involve nuclear factor kappaB-mediated induction of the Ets transcription factor ESE-1. *Arthritis Rheum* 2003, **48**:1249–1260.
- 14 Voronov E, Shouval DS, Krelin Y, et al.: IL-1 is required for tumor invasiveness and angiogenesis. *Proc Natl Acad Sci U S A* 2003, **100**:2645–2650.
- Given that the RA synovium has been compared to a situation with tumor-like properties, the effect of IL-1 on tumor invasiveness and new blood vessel formation provides arguments for the inhibition of IL-1 in RA inflammation.
- 15 Nakahara H, Song J, Sugimoto M, et al.: Anti-interleukin-6 receptor antibody therapy reduces vascular endothelial growth factor production in rheumatoid arthritis. *Arthritis Rheum* 2003, **48**:1521–1529.
- 16 Miossec P: Interleukin-17 in rheumatoid arthritis: if T cells were to contribute to inflammation and destruction through synergy. *Arthritis Rheum* 2003, **48**:594–601.
- This review of IL-17 in arthritis focuses on the interaction between a T cell product such as IL-17 and monocyte products such as TNF and IL-1. Knowledge of such

synergistic interactions may provide a better understanding of the local cytokine situation in the RA joint. These same concepts may apply to therapeutic modulation.

- 17 Aggarwal S, Ghilardi N, Xie MH, et al.: Interleukin-23 promotes a distinct CD4 • T cell activation state characterized by the production of interleukin-17. *J Biol Chem* 2003, 278:1910–1914.

The new IL-23 is shown to interact with T cells leading to the production of IL-17, which in turn acts on proinflammatory cytokine production by monocytes.

- 18 Nakae S, Saijo S, Horai R, et al.: IL-17 production from activated T cells is • required for the spontaneous development of destructive arthritis in mice deficient in IL-1 receptor antagonist. *Proc Natl Acad Sci U S A* 2003, 100: 5986–5990.

This paper demonstrates the role of T cells through IL-17 production in joint inflammation. Mice deficient in IL-1Ra show a destructive arthritis only when synergistic signals are provided by T cells through IL-17.

- 19 Udagawa N, Kotake S, Kamatani N, et al.: The molecular mechanism of os- • teoclastogenesis in rheumatoid arthritis. *Arthritis Res* 2002, 4:281–289.

This is a very good review of osteoclastogenesis in RA, focusing on the contribution of various cytokines and their producing cell subsets.

- 20 Pacquelet S, Presle N, Boileau C, et al.: Interleukin 17, a nitric oxide-pro- • ducing cytokine with a peroxynitrite-independent inhibitory effect on proteoglycan synthesis. *J Rheumatol* 2002, 29:2602–2610.

- 21 Kehlen A, Pachnio A, Thiele K, Langner J: Gene expression induced by inter- • leukin-17 in fibroblast-like synoviocytes of patients with rheumatoid arthritis: upregulation of hyaluronan-binding protein TSG-6. *Arthritis Res Ther* 2003, 5:R186–R192.

- 22 Rifas L, Arackal S: T cells regulate the expression of matrix metalloproteinase • in human osteoblasts via a dual mitogen-activated protein kinase mechanism. *Arthritis Rheum* 2003, 48:993–1001.

- 23 Kawashima M, Miossec P: Heterogeneity of response of rheumatoid syno- • vium cell subsets to interleukin-18 in relation to differential interleukin-18 receptor expression. *Arthritis Rheum* 2003, 48:631–637.

The discrepancy between results on the effect of IL-18 on synoviocytes is related to the lack of functional IL-18 receptors on these cells.

- 24 Joosten LA, Radstake TR, Lubberts E, et al.: Association of interleukin-18 • expression with enhanced levels of both interleukin-1beta and tumor necrosis factor alpha in knee synovial tissue of patients with rheumatoid arthritis. *Arthritis Rheum* 2003, 48:339–347.

This clinical study indicates the link between the expression of IL-18 and that of IL-1 and TNF in sections of RA synovium. As such, these results suggest the control of IL-18 during chronic inflammation.

- 25 Komai-Koma M, Gracie JA, Wei XQ, et al.: Chemoattraction of human T cells • by IL-18. *J Immunol* 2003, 170:1084–1090.

This paper shows an additional effect of IL-18 on T cells. Combined with IL-12, IL-18 is a potent inducer of interferon- γ . A chemotactic effect on T cells both *in vitro* and *in vivo* is shown.

- 26 Moller B, Kukoc-Zivojnov N, Koyama N, et al.: Prednisolone induces interleu- • kin-18 expression in mononuclear blood and myeloid progenitor cells. *Inflamm Res* 2002, 51:457–463.

- 27 Smeets RL, van de Loo FA, Arntz OJ, et al.: Adenoviral delivery of IL-18 bind- • ing protein C ameliorates collagen-induced arthritis in mice. *Gene Ther* 2003, 10:1004–1011.

This preclinical study shows the interest in controlling joint inflammation by the administration of IL-18 binding protein, which is a natural decoy inhibitor of IL-18. Protection was observed in the arthritis model through adenoviral delivery. It remains to be calculated how much would be needed to obtain the same results in human RA.

- 28 Moller B, Paulukat J, Nold M, et al.: Interferon-gamma induces expression of • interleukin-18 binding protein in fibroblast-like synoviocytes. *Rheumatology (Oxford)* 2003, 42:442–445.

- 29 Cua DJ, Sherlock J, Chen Y, et al.: Interleukin-23 rather than interleukin-12 is • the critical cytokine for autoimmune inflammation of the brain. *Nature* 2003, 421:744–748.

This is the first paper to differentiate the role of IL-23 from that of IL-12 in experimental encephalomyelitis. Clarification was made possible through the use of both IL-12 and IL-23 specific inhibitors. Because this is a general question, these results are also of interest in the field of arthritis.

- 30 Morand EF, Bucala R, Leech M: Macrophage migration inhibitory factor: an • emerging therapeutic target in rheumatoid arthritis. *Arthritis Rheum* 2003, 48:291–299.

A very good review of the role of MIF in RA. Details on the role of MIF in synovial cell proliferation and p53 regulation are described in [31] and [32].

- 31 Lacey D, Sampey A, Mitchell R, et al.: Control of fibroblast-like synovial cell • proliferation by macrophage migration inhibitory factor. *Arthritis Rheum* 2003, 48:103–109.

- 32 Leech M, Lacey D, Xue JR, et al.: Regulation of p53 by macrophage migration • inhibitory factor in inflammatory arthritis. *Arthritis Rheum* 2003, 48:1881–1889.

- 33 De Benedetti F, Meazza C, Vivarelli M, et al.: Functional and prognostic rel- • evance of the -173 polymorphism of the macrophage migration inhibitory factor gene in systemic-onset juvenile idiopathic arthritis. *Arthritis Rheum* 2003, 48:1398–1407.