#### C.01.01.Z1 – Biblioteca Simplificada de Gás Ideal

Aplicação em FTHA – Finite Time Heat Addition Otto Engine Model

Prof. C. Naaktgeboren, PhD



https://github.com/CNThermSci/ApplThermSci Compiled on 2020-09-09 06h15m33s UTC





$$Pv = RT$$

$$P\bar{v} = \bar{R}T$$
  $\rightarrow$ 





$$Pv = RT$$

$$P = \frac{RT}{v}$$

$$P\bar{v} = \bar{R}T$$

$$P = rac{RT}{ar{v}}$$
 —





$$Pv = RT$$

$$P = \frac{RT}{v}$$

$$T = \frac{Pv}{R}$$

$$P\bar{v} = \bar{R}T$$

$$P = rac{ar{R}T}{ar{v}}$$
 —

$$T = \frac{P\bar{\nu}}{\bar{R}}$$





$$Pv = RT$$

$$P = \frac{RT}{v}$$

$$T = \frac{Pv}{R}$$

$$RT$$

$$P\bar{v} = \bar{R}T$$
  $\rightarrow$ 
 $P = \frac{\bar{R}T}{\bar{v}}$   $\rightarrow$ 
 $T = \frac{P\bar{v}}{\bar{R}}$   $\rightarrow$ 
 $\bar{v} = \frac{\bar{R}T}{\bar{R}}$   $\therefore$ 





$$Pv = RT$$
  $P\bar{v} = \bar{R}T$   $\neg$ 

$$P = \frac{RT}{v}$$
  $P = \frac{\bar{R}T}{\bar{v}}$   $\neg$ 

$$T = \frac{Pv}{R}$$
  $T = \frac{P\bar{v}}{\bar{R}}$   $\neg$ 

$$v = \frac{RT}{P}$$
  $\bar{v} = \frac{\bar{R}T}{P}$   $\vdots$ 

Cada equação com forma nas bases mássica, e molar, com  $R = \bar{R}/M$  — armazenar  $\bar{R}$  e M!





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1},$$

$$T_{min} \leqslant T \leqslant T_{max}$$
  $\rightarrow$ 





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1},$$

$$\bar{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3,$$

$$T_{min} \leqslant T \leqslant T_{max}$$
 -

$$T_{min} \leqslant T \leqslant T_{max}$$
  $\neg$ 





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{c}_v(T) = \bar{c}_p(T) - \bar{R} = \sum_{i=1}^4 b_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max}$$





$$ar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} - ar{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3, \qquad T_{min} \leqslant T \leqslant T_{max} - ar{c}_v(T) = ar{c}_p(T) - ar{R} = \sum_{i=1}^4 b_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} - ar{c}_v(T) = a_1 - ar{R}.$$





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \overline{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \overline{c}_p(T) = \bar{c}_p(T) - \bar{R} = \sum_{i=1}^4 b_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \overline{c}_p(T) = a_1 - \bar{R}, \qquad b_{i>1} = a_{i>1} \qquad \vdots$$

Armazenar  $a_i$ ,  $T_{min}$  e  $T_{max}$  e saber as conversões (i)  $a_i \to b_i$  e (ii)  $\bar{c}_{p,v}(T) \to c_{p,v}(T)$ 





