C.01.01.Z1 – Biblioteca Simplificada de Gás Ideal

Aplicação em FTHA – Finite Time Heat Addition Otto Engine Model

Prof. C. Naaktgeboren, PhD



https://github.com/CNThermSci/ApplThermSci Compiled on 2020-09-09 07h14m55s UTC





$$Pv = RT$$

$$P\bar{v} = \bar{R}T$$
 \rightarrow





$$Pv = RT$$

$$P = \frac{RT}{v}$$

$$P\bar{v} = \bar{R}T$$

$$P = rac{RT}{ar{v}}$$
 —





$$Pv = RT$$

$$P = \frac{RT}{v}$$

$$T = \frac{Pv}{R}$$

$$P\bar{v} = \bar{R}T$$

$$P = rac{ar{R}T}{ar{v}}$$
 —

$$T = \frac{P\bar{\nu}}{\bar{R}}$$





$$Pv = RT$$

$$P = \frac{RT}{v}$$

$$T = \frac{Pv}{R}$$

$$RT$$

$$P\bar{v} = \bar{R}T$$
 \rightarrow
 $P = \frac{\bar{R}T}{\bar{v}}$ \rightarrow
 $T = \frac{P\bar{v}}{\bar{R}}$ \rightarrow
 $\bar{v} = \frac{\bar{R}T}{\bar{R}}$ \therefore





$$Pv = RT$$
 $P\bar{v} = \bar{R}T$ \neg

$$P = \frac{RT}{v}$$
 $P = \frac{\bar{R}T}{\bar{v}}$ \neg

$$T = \frac{Pv}{R}$$
 $T = \frac{P\bar{v}}{\bar{R}}$ \neg

$$v = \frac{RT}{P}$$
 $\bar{v} = \frac{\bar{R}T}{P}$ \vdots

Cada equação com forma nas bases mássica, e molar, com $R = \bar{R}/M$ — armazenar \bar{R} e M!





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1},$$

$$T_{min} \leqslant T \leqslant T_{max}$$
 \rightarrow





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1},$$

$$\bar{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3,$$

$$T_{min} \leqslant T \leqslant T_{max}$$
 -

$$T_{min} \leqslant T \leqslant T_{max}$$
 \neg





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{c}_v(T) = \bar{c}_p(T) - \bar{R} = \sum_{i=1}^4 b_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max}$$





$$ar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} - ar{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3, \qquad T_{min} \leqslant T \leqslant T_{max} - ar{c}_v(T) = ar{c}_p(T) - ar{R} = \sum_{i=1}^4 b_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} - ar{c}_v(T) = a_1 - ar{R}.$$





$$\bar{c}_p(T) = \sum_{i=1}^4 a_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \overline{c}_p(T) = a_1 + a_2 T + a_3 T^2 + a_4 T^3, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \overline{c}_p(T) = \bar{c}_p(T) - \bar{R} = \sum_{i=1}^4 b_i T^{i-1}, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \overline{c}_p(T) = a_1 - \bar{R}, \qquad b_{i>1} = a_{i>1} \qquad \vdots$$

Armazenar a_i , T_{min} e T_{max} e saber as conversões (i) $a_i \to b_i$ e (ii) $\bar{c}_{p,v}(T) \to c_{p,v}(T)$





Modelo de $\bar{c}_p(T)$ Polinomial: $\bar{u}(T)$

$$ar{u}(T) = \int_{T_{ref}}^T ar{c}_{\scriptscriptstyle \mathcal{V}}(T) \, dT = \int_{T_{ref}}^T \sum_{i=1}^4 b_i T^{i-1} \, dT, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad -$$





Modelo de $\bar{c}_n(T)$ Polinomial: $\bar{u}(T)$

$$ar{u}(T) = \int_{T_{vorf}}^T ar{c}_{\scriptscriptstyle \mathcal{V}}(T) \, dT = \int_{T_{vorf}}^T \sum_{i=1}^4 b_i T^{i-1} \, dT,$$

$$ar{u}(T) = \left(b_1 T + rac{b_2 T^2}{2} + rac{b_3 T^3}{3} + rac{b_4 T^4}{4}
ight)_{T_{ref}}^T, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$T_{min} \leqslant T \leqslant T_{max}$$
 -

$$\leqslant T \leqslant T_{max}$$





Modelo de $\bar{c}_n(T)$ Polinomial: $\bar{u}(T)$

$$ar{u}(T) = \int_{T_{ref}}^T ar{c}_{v}(T) dT = \int_{T_{ref}}^T \sum_{i=1}^4 b_i T^{i-1} dT, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{u}(T) = \left(b_1 T + \frac{b_2 T^2}{2} + \frac{b_3 T^3}{3} + \frac{b_4 T^4}{4}\right)_{T=1}^T, \qquad T_{min} \leqslant T \leqslant T_{max}$$

Armazenar T_{ref} ,





Modelo de $\bar{c}_p(T)$ Polinomial: $\bar{u}(T)$

$$ar{u}(T) = \int_{T_{ref}}^T ar{c}_v(T) \, dT = \int_{T_{ref}}^T \sum_{i=1}^4 b_i T^{i-1} \, dT, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad -$$

$$\bar{u}(T) = \left(b_1 T + \frac{b_2 T^2}{2} + \frac{b_3 T^3}{3} + \frac{b_4 T^4}{4}\right)_{T=1}^T, \qquad T_{min} \leqslant T \leqslant T_{max}$$

- Armazenar T_{ref} ,
- Compor eficientemente a soma de produtos, e





Modelo de $\bar{c}_p(T)$ Polinomial: $\bar{u}(T)$

$$ar{u}(T) = \int_{T_{min}}^T ar{c}_v(T) dT = \int_{T_{min}}^T \sum_{i=1}^4 b_i T^{i-1} dT, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{u}(T) = \left(b_1 T + \frac{b_2 T^2}{2} + \frac{b_3 T^3}{3} + \frac{b_4 T^4}{4}\right)_{T=1}^T, \qquad T_{min} \leqslant T \leqslant T_{max}$$

- Armazenar T_{ref} ,
- Compor eficientemente a soma de produtos, e
- Saber as conversões $\bar{u}(T) \rightarrow u(T)$.





$$ar{h}(T) = \int_{T_{ref}}^T ar{c}_p(T) dT + ar{R}T_{ref} = \int_{T_{ref}}^T \sum_{i=1}^4 a_i T^{i-1} dT + ar{R}T_{ref}, \quad T_{min} \leqslant T \leqslant T_{max}$$





$$ar{h}(T) = \int_{T_{ref}}^T ar{c}_p(T) dT + ar{R}T_{ref} = \int_{T_{ref}}^T \sum_{i=1}^4 a_i T^{i-1} dT + ar{R}T_{ref}, \quad T_{min} \leqslant T \leqslant T_{max} \qquad -$$

$$\bar{h}(T) = \left(a_1 T + \frac{a_2 T^2}{2} + \frac{a_3 T^3}{3} + \frac{a_4 T^4}{4}\right)_{T_{ref}}^T + \bar{R} T_{ref}, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \therefore$$





$$\bar{h}(T) = \int_{T_{ref}}^{T} \bar{c}_{p}(T) dT + \bar{R}T_{ref} = \int_{T_{ref}}^{T} \sum_{i=1}^{4} a_{i}T^{i-1} dT + \bar{R}T_{ref}, \quad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{h}(T) = \left(a_{1}T + \frac{a_{2}T^{2}}{2} + \frac{a_{3}T^{3}}{3} + \frac{a_{4}T^{4}}{4}\right)_{T}^{T} + \bar{R}T_{ref}, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\therefore$$

• Compor eficientemente a soma de produtos, e





$$ar{h}(T) = \int_{T_{ref}}^{T} ar{c}_p(T) dT + ar{R}T_{ref} = \int_{T_{ref}}^{T} \sum_{i=1}^{4} a_i T^{i-1} dT + ar{R}T_{ref}, \quad T_{min} \leqslant T \leqslant T_{max}$$

• Saber as conversões
$$\bar{h}(T) \rightarrow h(T)$$
.





Modelo de
$$\bar{c}_p(T)$$
 Polinomial: $\bar{s}^{\circ}(T)$

$$ar{s}^{\circ}(T) = \int_0^T rac{ar{c}_p(T)}{T} \, dT = \int_0^{T_{ref}} rac{ar{c}_p(T)}{T} \, dT + \int_{T_{ref}}^T rac{ar{c}_p(T)}{T} \, dT, \quad T_{min} \leqslant T \leqslant T_{max}$$





$$\bar{s}^{\circ}(T) = \int_0^T \frac{\bar{c}_p(T)}{T} dT = \int_0^{T_{ref}} \frac{\bar{c}_p(T)}{T} dT + \int_{T_{ref}}^T \frac{\bar{c}_p(T)}{T} dT, \quad T_{min} \leqslant T \leqslant T_{max} \qquad -$$

$$ar{s}^{\circ}(T) = ar{s}_{ref}^{\circ} + \int_{T_{ref}}^{T} \sum_{i=1}^{4} a_i T^{i-2} dT$$
 $T_{min} \leqslant T \leqslant T_{max}$





$$ar{s}^{\circ}(T) = \int_{0}^{T} rac{ar{c}_{p}(T)}{T} dT = \int_{0}^{T_{ref}} rac{ar{c}_{p}(T)}{T} dT + \int_{T_{ref}}^{T} rac{ar{c}_{p}(T)}{T} dT, \quad T_{min} \leqslant T \leqslant T_{max}$$

$$- ar{s}^{\circ}(T) = ar{s}^{\circ}_{ref} + \int_{T_{ref}}^{T} \sum_{i=1}^{4} a_{i} T^{i-2} dT \qquad T_{min} \leqslant T \leqslant T_{max} \qquad - ar{s}^{\circ}_{ref} + ar{s}^{\circ}_{ref} +$$

$$\int_{T_{ref}} i=1$$

$$\int_{T_{ref}} a_{1}T^{2} = a_{1}T^{3} \setminus_{T}^{T}$$

$$ar{s}^{\circ}(T) = ar{s}_{ref}^{\circ} + \left(a_1 \ln(T) + a_2 T + rac{a_3 T^2}{2} + rac{a_4 T^3}{3}
ight)_T^T$$
, $T_{min} \leqslant T \leqslant T_{max}$.





$$\bar{s}^{\circ}(T) = \int_{0}^{T} \frac{\bar{c}_{p}(T)}{T} dT = \int_{0}^{T_{ref}} \frac{\bar{c}_{p}(T)}{T} dT + \int_{T_{ref}}^{T} \frac{\bar{c}_{p}(T)}{T} dT, \quad T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{s}^{\circ}(T) = \bar{s}^{\circ}_{ref} + \int_{T_{ref}}^{T} \sum_{i=1}^{4} a_{i} T^{i-2} dT$$

$$T_{min} \leqslant T \leqslant T_{max}$$

$$\bar{s}^{\circ}(T) = \bar{s}^{\circ}_{ref} + \left(a_{1} \ln(T) + a_{2} T + \frac{a_{3} T^{2}}{2} + \frac{a_{4} T^{3}}{3}\right)_{T}^{T}, \qquad T_{min} \leqslant T \leqslant T_{max}$$

$$\therefore$$

• Armazenar \bar{s}_{ref}° , compor eficientemente a soma de produtos, e





$$\bar{s}^{\circ}(T) = \int_{0}^{T} \frac{\bar{c}_{p}(T)}{T} dT = \int_{0}^{T_{ref}} \frac{\bar{c}_{p}(T)}{T} dT + \int_{T_{ref}}^{T} \frac{\bar{c}_{p}(T)}{T} dT, \quad T_{min} \leqslant T \leqslant T_{max} \qquad \neg$$

$$\bar{s}^{\circ}(T) = \bar{s}^{\circ}_{ref} + \int_{T_{ref}}^{T} \sum_{i=1}^{4} a_{i} T^{i-2} dT \qquad \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \neg$$

$$\bar{s}^{\circ}(T) = \bar{s}^{\circ}_{ref} + \left(a_{1} \ln(T) + a_{2} T + \frac{a_{3} T^{2}}{2} + \frac{a_{4} T^{3}}{3} \right)_{T}^{T}, \qquad T_{min} \leqslant T \leqslant T_{max} \qquad \therefore$$

- Armazenar \bar{s}_{ref}° , compor eficientemente a soma de produtos, e
- Saber as conversões $\bar{s}^{\circ}(T) \to s^{\circ}(T)$.





