COMP 3721

Tutorial 1

- Xiao Huiru
- hxiaoaf@connect.ust.hk
- No fixed Office Hour
- Feel free to contact with me if any questions

- Tutorials:
 - Solutions will be posted on tutorial webpage after each tutorial.

• Languages and Regular Expressions

- Determine whether each of the following is true or false.
 - a) $\emptyset \in \emptyset$
 - b) $\emptyset \subset \emptyset$
 - c) $\{a, b\} \subseteq \{a, b, \{a, b\}\}$
 - d) $\{a,b\} \in \{a,b,\{a,b\}\}$

- 1) Determine whether each of the following is true or false.
 - a) $\emptyset \in \emptyset$ False. Empty set contains no elements.
 - b) $\emptyset \subset \emptyset$
 - c) $\{a, b\} \subseteq \{a, b, \{a, b\}\}$
 - d) $\{a,b\} \in \{a,b,\{a,b\}\}$

- 1) Determine whether each of the following is true or false.
 - a) $\emptyset \in \emptyset$ False.
 - b) $\emptyset \subset \emptyset$ True.
 - c) $\{a, b\} \subseteq \{a, b, \{a, b\}\}$
 - d) $\{a,b\} \in \{a,b,\{a,b\}\}$

- 1) Determine whether each of the following is true or false.
 - a) $\emptyset \in \emptyset$ False.
 - b) $\emptyset \subset \emptyset$ True.
 - c) $\{a,b\} \subseteq \{a,b,\{a,b\}\}$ True.
 - d) $\{a,b\} \in \{a,b,\{a,b\}\}$

- 1) Determine whether each of the following is true or false.
 - a) $\emptyset \in \emptyset$ False.
 - b) $\emptyset \subset \emptyset$ True.
 - c) $\{a,b\} \subseteq \{a,b,\{a,b\}\}$ True.
 - d) $\{a,b\} \in \{a,b,\{a,b\}\}$ True.

• Definition:

Regular expressions

Regular expressions are a *finite* representation of languages.

Inductive definition of regular expressions for languages over an alphabet Σ . A regular expression is a string over alphabet $\Sigma_1 = \Sigma \cup \{(,),\emptyset,\cup,^*\}$.

- 1. \emptyset and each $\sigma \in \Sigma$ are regular expressions.
- 2. If α and β are regular expressions, then

$$(\alpha\beta), (\alpha\cup\beta), \alpha^*$$

are regular expressions.

3. Nothing else is a regular expression.

2) Show that if a and b are distinct symbols, then

$${a,b}^* = {a}^*({b}{a}^*)^*$$

Hint:

• Two sets A and B are equal (A = B) if $A \subseteq B$ and $B \subseteq A$.

2) Show that if a and b are distinct symbols, then

$${a,b}^* = {a}^*({b}{a}^*)^*$$

Proof:

It is clear that $\{a\}^*(\{b\}\{a\}^*)^* \subseteq \{a,b\}^*$. Need to show that $\{a,b\}^* \subseteq \{a\}^*(\{b\}\{a\}^*)^*$.

2) Show that if a and b are distinct symbols, then

$${a,b}^* = {a}^*({b}{a}^*)^*$$

Proof:

It is clear that ${a}^*({b}{a}^*)^* \subseteq {a,b}^*$.

Need to show that $\{a,b\}^* \subseteq \{a\}^*(\{b\}\{a\}^*)^*$.

For any string $w \in \{a,b\}^*$. If we mark all the occurrence of b's in w, then w can be rewritten as: $w \in a^*ba^*ba^*ba^*ba^*ba^*$

3) True or False. Justification.

```
a) abcd \in (a(cd)^*b)^*
```

b)
$$\{a^nb^n: n \ge 0\}\{b^nc^n: n \ge 0\} = \{a^nb^{2n}c^n: n \ge 0\}$$

3) True or False. Justification.

a)
$$abcd \in (a(cd)^*b)^*$$

False.

Because any nonempty string in $(a(cd)^*b)^*$ must ends with b.

3) True or False. Justification.

b)
$${a^nb^n : n \ge 0}{b^nc^n : n \ge 0} = {a^nb^{2n}c^n : n \ge 0}$$

False.

Counter example: abbbcc

- 4) Let $\Sigma = \{a, b\}$. Write regular expressions for the following sets.
 - a) All strings in Σ^* with no more than 3 a's.
 - b) All strings in Σ^* with a number of a's divisible by 3.
 - c) All strings in Σ^* that does not have aab as a substring.

- 4) Let $\Sigma = \{a, b\}$. Write regular expressions for the following sets.
 - a) All strings in Σ^* with no more than 3 a's.

$$b^*(a \cup \emptyset)b^*(a \cup \emptyset)b^*(a \cup \emptyset)b^*$$

- b) All strings in Σ^* with a number of a's divisible by 3.
- c) All strings in Σ^* that does not have aab as a substring.

- 4) Let $\Sigma = \{a, b\}$. Write regular expressions for the following sets.
 - a) All strings in Σ^* with no more than 3 a's.

$$b^*(a \cup \emptyset)b^*(a \cup \emptyset)b^*(a \cup \emptyset)b^*$$

b) All strings in Σ^* with a number of a's divisible by 3.

$$b^*(b^*ab^*ab^*ab^*)^*$$

c) All strings in Σ^* that does not have *aab* as a substring.

- 4) Let $\Sigma = \{a, b\}$. Write regular expressions for the following sets.
 - a) All strings in Σ^* with no more than 3 a's.

$$b^*(a \cup \emptyset)b^*(a \cup \emptyset)b^*(a \cup \emptyset)b^*$$

b) All strings in Σ^* with a number of a's divisible by 3.

$$b^*(b^*ab^*ab^*ab^*)^*$$

c) All strings in Σ^* that does not have *aab* as a substring.

$$(ab \cup b)^*a^*$$

5) Prove that if *L* is regular, then so is

$$\operatorname{Pref}(L) = \{w : wu \in L \text{ for some string u}\}.$$

Hint: Follow the definition of regular expression.

- Proof(sketch):
 - i. We first prove that if for any $L(a), a \in \Sigma \cup \emptyset$, then Pref(L(a)) is regular.
 - ii. Then we prove that if L1 and L2 are regular and have their *Pref* also regular, so do their *union*, concatenation and *Kleene Star*.