



COMP 3721

Tutorial 4

- NFA \rightarrow Regular Expression
- NFA = DFA = Regular Expressions
- Pumping Theorem


NFA \rightarrow Regular Expression

- 1) Write regular expressions for the languages accepted by the following NFA.




NFA \rightarrow Regular Expression

- 1) Write regular expressions for the languages accepted by the following NFA.




a^*




NFA \rightarrow Regular Expression

- 1) Write regular expressions for the languages accepted by the following NFA.



a^*



$(a(ba \cup b))^*$

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

a) $Subseq(L) = \{w_1 \dots w_k : x = x_0 w_1 \dots w_k x_k \in L \text{ for some } x_0, \dots, x_k\}$.

b) $L^R = \{w : w^R \in L\}$.

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

a) $Subseq(L) = \{w_1 \dots w_k : x = x_0 w_1 \dots w_k x_k \in L \text{ for some } x_0, \dots, x_k\}$.

Proof:

Since L is regular, it is accepted by some NFA $M = \{K, \Sigma, \Delta, s, F\}$.

Next we will construct a NFA M' that accepts $Subseq(L)$.

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

a) $Subseq(L) = \{w_1 \dots w_k : x = x_0 w_1 \dots w_k x_k \in L \text{ for some } x_0, \dots, x_k\}$.

Proof:

Since L is regular, it is accepted by some NFA $M = \{K, \Sigma, \Delta, s, F\}$.

Next we will construct a NFA M' that accepts $Subseq(L)$.

We look at any pair (p, q) of states in M , if there is a path from p to q , we draw an arrow from p to q labeled with e .

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

a) $Subseq(L) = \{w_1 \dots w_k : x = x_0 w_1 \dots w_k x_k \in L \text{ for some } x_0, \dots, x_k\}$.

Proof:

Since L is regular, it is accepted by some NFA $M = \{K, \Sigma, \Delta, s, F\}$.

Next we will construct a NFA M' that accepts $Subseq(L)$.

We look at any pair (p, q) of states in M , if there is a path from p to q , we draw an arrow from p to q labeled with e .

More precisely, $M' = (K, \Sigma, \Delta', s, F)$ where

$\Delta' = \Delta \cup \{(p, e, q) : \text{there is a path from } p \text{ to } q \text{ in } M\}$.

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

a) $Subseq(L) = \{w_1 \dots w_k : x = x_0 w_1 \dots w_k x_k \in L \text{ for some } x_0, \dots, x_k\}$.

Proof:

Since L is regular, it is accepted by some NFA $M = \{K, \Sigma, \Delta, s, F\}$.

Next we will construct a NFA M' that accepts $Subseq(L)$.

We look at any pair (p, q) of states in M , if there is a path from p to q , we draw an arrow from p to q labeled with e .

More precisely, $M' = (K, \Sigma, \Delta', s, F)$ where

$\Delta' = \Delta \cup \{(p, e, q) : \text{there is a path from } p \text{ to } q \text{ in } M\}$.

We claim that M' accepts $Subseq(L)$ and hence $Subseq(L)$ is regular. Details are omitted

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

b) $L^R = \{w : w^R \in L\}.$

Proof:

Since L is regular, it is accepted by some NFA $M = \{K, \Sigma, \Delta, s, f\}$. Without loss of generality, we assume that M has single final state.

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

b) $L^R = \{w : w^R \in L\}.$

Proof:

Since L is regular, it is accepted by some NFA $M = \{K, \Sigma, \Delta, s, f\}$. Without loss of generality, we assume that M has single final state.

Next we will construct a NFA M' that accepts L^R by reversing all the transitions in M .

NFA = DFA = Regular Expressions

2) Prove that if L is regular, then the following languages are also regular

b) $L^R = \{w : w^R \in L\}.$

Proof:

Since L is regular, it is accepted by some NFA $M = \{K, \Sigma, \Delta, s, f\}$. Without loss of generality, we assume that M has single final state.

Next we will construct a NFA M' that accepts L^R by reversing all the transitions in M .

More precisely, we let $M' = (K, \Sigma, \Delta', s', f')$ where $s' = f$, $f' = s$, and $\Delta' = \{(p, a, q) : (q, a, p) \in \Delta\}.$

We claim that M' accepts L^R and hence L^R is regular.

Details are omitted

Pumping Theorem

We use the pumping theorem to prove that a language is NOT regular. To prove that L is not regular, we show the following, which contradicts the P.T.:

$$\forall n \geq 1,$$

$$\exists w \in L, |w| \geq n,$$

$$\forall x, y, z \text{ such that } w = xyz \text{ and } y \neq e, |xy| \leq n,$$

$$\text{but } \exists i \geq 0, xy^iz \notin L.$$

Pumping Theorem

- Assume L is regular.
- Then, by Pumping Theorem, there exists an integer n such that *all* strings of length $\geq n$ in L can be pumped.
- Find *one* string w in L that has length $\geq n$ and is not pumpable no matter how you split it into 3 parts, i.e.,:
 - Choose *one* string $w \in L$ with $|w| \geq n$.
 - Consider *all* valid ways of splitting w into x, y, z .
 - For each way of splitting, demonstrate *one* value i such that $xy^iz \notin L$
- The existence of one string w that is not pumpable contradicts the Pumping Theorem.
- Hence L is not regular.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$

b) $L = \{ww : w \in \{a, b\}^*\}$

c) $L = \{(bab)^i (babbab)^i : i \geq 1\}.$

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

Proof: Assume L is regular. Then pumping theorem applies for some integer n .

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

Proof: Assume L is regular. Then pumping theorem applies for some integer n .

Consider the string $w = a^{n+1}b^n \in L$.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

Proof: Assume L is regular. Then pumping theorem applies for some integer n .

Consider the string $w = a^{n+1}b^n \in L$.

Let x , y , and z be specified as in pumping theorem.

Then $x = a^i$, $y = a^j$, $z = a^k b^n$ for $i, k \geq 0$ and $j > 0$.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

Proof: Assume L is regular. Then pumping theorem applies for some integer n .

Consider the string $w = a^{n+1}b^n \in L$.

Let x , y , and z be specified as in pumping theorem.

Then $x = a^i$, $y = a^j$, $z = a^k b^n$ for $i, k \geq 0$ and $j > 0$.

By the theorem, xy^0z is in L .

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

Proof: Assume L is regular. Then pumping theorem applies for some integer n .

Consider the string $w = a^{n+1}b^n \in L$.

Let x , y , and z be specified as in pumping theorem.

Then $x = a^i$, $y = a^j$, $z = a^k b^n$ for $i, k \geq 0$ and $j > 0$.

By the theorem, xy^0z is in L .

But this is impossible since there are at most n a 's and exactly n b 's in xy^0z .

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

Proof: Assume L is regular. Then pumping theorem applies for some integer n .

Consider the string $w = a^{n+1}b^n \in L$.

Let x , y , and z be specified as in pumping theorem.

Then $x = a^i$, $y = a^j$, $z = a^k b^n$ for $i, k \geq 0$ and $j > 0$.

By the theorem, xy^0z is in L .

But this is impossible since there are at most n a 's and exactly n b 's in xy^0z .

Hence L is not regular.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

b) $L = \{ww : w \in \{a, b\}^*\}$

c) $L = \{(bab)^i (babbab)^i : i \geq 1\}$.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

b) $L = \{ww : w \in \{a, b\}^*\}$ Not regular.

c) $L = \{(bab)^i (babbab)^i : i \geq 1\}$.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

b) $L = \{ww : w \in \{a, b\}^*\}$ Not regular.

Proof: Assume L is regular. Then pumping theorem applies for some integer n .

Consider the string $w = a^{n+1}ba^{n+1}b \in L$.

Let x , y , and z be specified as in pumping theorem.

Then $x = a^i$, $y = a^j$, $z = a^kba^{n+1}b$ for $i, k \geq 0$ and $j > 0$.

By the theorem, xy^0z is in L .

But this is impossible since there are at most n a 's and exactly n b 's in xy^0z .

Hence L is not regular.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

b) $L = \{ww : w \in \{a, b\}^*\}$ Not regular.

c) $L = \{(bab)^i (babbab)^i : i \geq 1\}$.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.

b) $L = \{ww : w \in \{a, b\}^*\}$ Not regular.

c) $L = \{(bab)^i (babbab)^i : i \geq 1\}$. Regular.

Pumping Theorem

3) Are the following languages on $\{a, b\}$ regular or not? Prove your answer.

- a) $L = \{a^i b^j : i > j \geq 1\}$ Not regular.
- b) $L = \{ww : w \in \{a, b\}^*\}$ Not regular.
- c) $L = \{(bab)^i (babbab)^i : i \geq 1\}$. Regular.

Proof.

In fact,

$$L = (babbabbab)^+ = (babbabbab)(babbabbab)^*$$

Properties of R. L.

- 4) Are the following statements true or false?
 - a) Every subset of a regular language is regular.

Properties of R. L.

- 4) Are the following statements true or false?
- a) Every subset of a regular language is regular.
False.

Properties of R. L.

4) Are the following statements true or false?

a) Every subset of a regular language is regular.

False.

Counter example:

$\{a^n b^n : n \geq 0\}$ is a subset of $(a \cup b)^*$, but is not regular.

Properties of R. L.

- 4) Are the following statements true or false?
- a) Every subset of a regular language is regular. False.
 - b) If L is regular, then so is $L' = \{xy : x \in L \text{ and } y \notin L\}$.

Properties of R. L.

- 4) Are the following statements true or false?
- a) Every subset of a regular language is regular. False.
 - b) If L is regular, then so is $L' = \{xy : x \in L \text{ and } y \notin L\}$.
True.

Properties of R. L.

- 4) Are the following statements true or false?
- a) Every subset of a regular language is regular. False.
 - b) If L is regular, then so is $L' = \{xy : x \in L \text{ and } y \notin L\}$.
True.

Regular language is closed under *Concatenation* and *Complementation*.

$$L' = L\bar{L}$$

Properties of R. L.

- 4) Are the following statements true or false?
- a) Every subset of a regular language is regular. False.
 - b) If L is regular, then so is $L' = \{xy : x \in L \text{ and } y \notin L\}$. True.
 - c) $\{w : w = w^R\}$ is regular.

Properties of R. L.

4) Are the following statements true or false?

- a) Every subset of a regular language is regular. False.
- b) If L is regular, then so is $L' = \{xy : x \in L \text{ and } y \notin L\}$. True.
- c) $\{w : w = w^R\}$ is regular.
False.

Can be proved by pumping theorem.

Properties of R. L.

- 4) Are the following statements true or false?
- a) Every subset of a regular language is regular. False.
 - b) If L is regular, then so is $L' = \{xy : x \in L \text{ and } y \notin L\}$. True.
 - c) $\{w : w = w^R\}$ is regular. False