COMPASS_TEMPEST_SGW_2024: November w/ original standard curve

Stephanie J. Wilson

2025-02-11

##Set Up

Read in first data file and assess standard curves

```
##     Machine    User Run_Date Sample_Type Sample1 Sample_Year Sample_Month
## 1 Varian GC Blakley 20260417        0.05 STD_CO2          NA           NA
## 2 Varian GC Blakley 20260417     STD_CO2 STD_CO2          NA           NA
## 3 Varian GC Blakley 20260417     STD_CO2 STD_CO2          NA           NA
## 4 Varian GC Blakley 20260417     STD_CO2 STD_CO2          NA           NA
## 5 Varian GC Blakley 20260417     STD_CO2 STD_CO2          NA           NA
## 6 Varian GC Blakley 20260417     STD_CO2 STD_CO2          NA           NA
##   Sample_Day Sample_Time Time_Zone G_W Sample_Plot Sample_Grid Sample_Depth
## 1         NA          NA        NA  NA          NA          NA           NA
## 2         NA          NA        NA  NA          NA          NA           NA
## 3         NA          NA        NA  NA          NA          NA           NA
## 4         NA          NA        NA  NA          NA          NA           NA
## 5         NA          NA        NA  NA          NA          NA           NA
## 6         NA          NA        NA  NA          NA          NA           NA
##   Sample_ID Dilution_Factor    STD_Conc CO2_Area CH4_Area Field_Notes Lab_Notes
## 1      0ppm               1 0.000000000        0       NA          NA        NA
## 2    1mL_LT               1 0.008333333       NA       NA          NA        NA
## 3    2mL_LT               1 0.016666667       NA       NA          NA        NA
## 4    6mL_LT               1 0.050000000    63988       NA          NA        NA
## 5   12mL_LT               1 0.100000000   130481       NA          NA        NA
## 6   24mL_LT               1 0.200000000   261002       NA          NA        NA
##   Sample_Order
## 1           NA
## 2           NA
## 3           NA
## 4           NA
## 5           NA
## 6           NA
```

```
## `geom_smooth()` using formula = 'y ~ x'

## Warning: Removed 1 row containing non-finite outside the scale range
## (`stat_smooth()`).

## Warning: Removed 1 row containing non-finite outside the scale range
## (`stat_poly_eq()`).
```

```
## Warning: Removed 1 row containing missing values or values outside the scale range
## (`geom_point()`).
```




```
##
## Call:
## lm(formula = stds_ch4_low$CH4_Area ~ stds_ch4_low$STD_Conc)
##
## Residuals:
##     Min      1Q  Median      3Q     Max
## -572.07 -340.33   83.54  242.01  575.16
##
## Coefficients:
##                       Estimate Std. Error t value Pr(>|t|)
## (Intercept)            688.089    180.870   3.804  0.00668 **
## stds_ch4_low$STD_Conc  752.098      4.685 160.530 9.61e-14 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 436.7 on 7 degrees of freedom
##   (1 observation deleted due to missingness)
## Multiple R-squared:  0.9997, Adjusted R-squared:  0.9997
## F-statistic: 2.577e+04 on 1 and 7 DF,  p-value: 9.605e-14
```

```
## `geom_smooth()` using formula = 'y ~ x'
```

```
## Warning: Removed 1 row containing non-finite outside the scale range
## (`stat_smooth()`).

## Warning: Removed 1 row containing non-finite outside the scale range
## (`stat_poly_eq()`).

## Warning: Removed 1 row containing missing values or values outside the scale range
## (`geom_point()`).
```




```
##
## Call:
## lm(formula = stds_ch4$CH4_Area ~ stds_ch4$STD_Conc)
##
## Residuals:
##    Min     1Q Median     3Q    Max
## -10484  -1944  -1510  -1101  20922
##
## Coefficients:
##                   Estimate Std. Error t value Pr(>|t|)
## (Intercept)       2345.350   2101.477   1.116    0.286
## stds_ch4$STD_Conc  752.105      1.378 545.932   <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 7050 on 12 degrees of freedom
```

```
##   (1 observation deleted due to missingness)
## Multiple R-squared:      1,  Adjusted R-squared:      1
## F-statistic: 2.98e+05 on 1 and 12 DF,  p-value: < 2.2e-16
```

```
## `geom_smooth()` using formula = 'y ~ x'
```

```
## Warning: Removed 4 rows containing non-finite outside the scale range
## (`stat_smooth()`).
```

```
## Warning: Removed 4 rows containing non-finite outside the scale range
## (`stat_poly_eq()`).
```

```
## Warning: Removed 4 rows containing missing values or values outside the scale range
## (`geom_point()`).
```




```
##
## Call:
## lm(formula = stds_co2$CO2_Area ~ stds_co2$STD_Conc)
##
## Residuals:
##    Min     1Q Median     3Q    Max
## -52641 -12644  -6920  12718  62934
##
## Coefficients:
```

```
##                    Estimate Std. Error t value Pr(>|t|)
## (Intercept)           10996      13670   0.804    0.444
## stds_co2$STD_Conc   1316151       6996 188.126 7.13e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 32560 on 8 degrees of freedom
##   (4 observations deleted due to missingness)
## Multiple R-squared:  0.9998, Adjusted R-squared:  0.9997
## F-statistic: 3.539e+04 on 1 and 8 DF,  p-value: 7.133e-16
```

```
##   X          Curve        R2    Slope  Intercept Run_Date
## 1 1  Slope_CH4_low 0.9996407 633.0314  -320.6965   202404
## 2 2 Slope_CH4_high 0.9993443 601.5512 14817.1191   202404
## 3 3      Slope_CO2 0.9999907 168.3200 10075.2183   202404
## 4 4  Slope_CH4_low 0.9996407 633.0314  -320.6965   202404
## 5 5 Slope_CH4_high 0.9993443 601.5512 14817.1191   202404
## 6 6      Slope_CO2 0.9999907 168.3200 10075.2183   202404
```



Now calculate the CH4 & CO2 concentrations in ppm

```
#head(raw)
```

```r
#pull out methane standards
Samples <-  raw %>%
  filter(!str_detect(Sample_Type, "STD_CH4")) %>%
  filter(!str_detect(Sample_Type, "STD_CO2")) %>%
  filter(!str_detect(Sample_Type, "Blank")) %>%
  filter(!str_detect(Sample_Type, "Chk_STD")) %>%
   filter(!str_detect(Sample_Type, "CHKSTD")) %>%
  filter(!str_detect(Sample_Type, "NA"))
#head(Samples)

#Now flag any areas that are above the 100ppm area for CH4
Samples$CH4_Curve <- ifelse(Samples$CH4_Area >71000, "High", "Low")
#head(Samples)

#Calculate CH4 concentrations in ppm
Samples$CH4_Conc_ppm <- ifelse(Samples$CH4_Area >71000, (Samples$CH4_Area-Slope_CH4_high$Intercept)/Slo

#Calculate CO2 concentrations in ppm
Samples$CO2_Conc_ppm <- ((Samples$CO2_Area-Slope_CO2$Intercept)/Slope_CO2$Slope)

#head(Samples)

##########make flags for any dilutions needed
#highest CH4 standard = 10000
#highest CO2 standard = 50000

Samples$CH4_Flag <- ifelse(Samples$CH4_Conc_ppm >10000, "Needs Dilution", "Within Range")
Samples$CO2_Flag <- ifelse(Samples$CO2_Conc_ppm >50000, "Needs Dilution", "Within Range")
#head(Samples)
```

Check the Check Standards

```
## Warning: Using ‘size‘ aesthetic for lines was deprecated in ggplot2 3.4.0.
## i Please use ‘linewidth‘ instead.
## This warning is displayed once per session.
## Call ‘lifecycle::last_lifecycle_warnings()‘ to see where this warning was
## generated.
```



Dilution correct samples

```
## Warning: Removed 1 row containing missing values or values outside the scale range
## (`geom_bar()`).
```


CH4: Green = Within Range
CH4 (ppm)
0
10
20
0mL_LT
0ppm
12mL_LT
24mL_LT
6mL_LT


CO2: Green = Within Range
CO2 (ppm)
0.0
0.1
0.2
0.3
0mL_LT
0ppm
12mL_LT
24mL_LT
6mL_LT

If samples are water calculate gas in water



Write out processed data & slopes - NOT THIS ONE

##Combine sample data with expected concentrations

Plot Expected vs Actual Conc





Plot Samples in Run Order