

Salinity affects microbial activity and soil organic matter content in tidal wetlands

EMBER M. MORRISSEY, JAIMIE L. GILLESPIE, JOSEPH C. MORINA and RIMA B. FRANKLIN

Department of Biology, Virginia Commonwealth University, 1000 W Cary Street, Richmond, VA 23284, USA

Abstract

Climate change-associated sea level rise is expected to cause saltwater intrusion into many historically freshwater ecosystems. Of particular concern are tidal freshwater wetlands, which perform several important ecological functions including carbon sequestration. To predict the impact of saltwater intrusion in these environments, we must first gain a better understanding of how salinity regulates decomposition in natural systems. This study sampled eight tidal wetlands ranging from freshwater to oligohaline (0–2 ppt) in four rivers near the Chesapeake Bay (Virginia). To help isolate salinity effects, sites were selected to be highly similar in terms of plant community composition and tidal influence. Overall, salinity was found to be strongly negatively correlated with soil organic matter content (OM%) and C : N, but unrelated to the other studied environmental parameters (pH, redox, and above- and below-ground plant biomass). Partial correlation analysis, controlling for these environmental covariates, supported direct effects of salinity on the activity of carbon-degrading extracellular enzymes (β -1, 4-glucosidase, 1, 4- β -cellobiosidase, β -D-xylosidase, and phenol oxidase) as well as alkaline phosphatase, using a per unit OM basis. As enzyme activity is the putative rate-limiting step in decomposition, enhanced activity due to salinity increases could dramatically affect soil OM accumulation. Salinity was also found to be positively related to bacterial abundance (qPCR of the 16S *rRNA* gene) and tightly linked with community composition (T-RFLP). Furthermore, strong relationships were found between bacterial abundance and/or composition with the activity of specific enzymes (1, 4- β -cellobiosidase, arylsulfatase, alkaline phosphatase, and phenol oxidase) suggesting salinity's impact on decomposition could be due, at least in part, to its effect on the bacterial community. Together, these results indicate that salinity increases microbial decomposition rates in low salinity wetlands, and suggests that these ecosystems may experience decreased soil OM accumulation, accretion, and carbon sequestration rates even with modest levels of saltwater intrusion.

Keywords: carbon cycling, decomposition, extracellular enzyme activity, marsh, microbial community structure, saltwater intrusion, sea level rise

Received 13 August 2013 and accepted 26 September 2013

Introduction

Climate change is predicted to alter the global hydrological cycle in many ways. For example, rising sea levels (Nakada & Inoue, 2005; Wigley, 2005; Church & White, 2006), reduced precipitation in watersheds (Smith *et al.*, 2005) with resulting declines in stream flow (Milley *et al.*, 2005), and global increases in water consumption (Gleick, 2003) may result in widespread saltwater intrusion into freshwater coastal ecosystems. Of particular concern is the impact of increased salinity on tidal freshwater wetlands, where it has been shown to drive changes in microbial metabolism (Weston *et al.*, 2011; Neubauer *et al.*, 2013), nutrient cycling (Weston *et al.*, 2006; Marton *et al.*, 2012), plant community composition (Sharpe & Baldwin, 2012), and primary production (Baldwin & Mendelssohn, 1998). Taken together, these changes may significantly alter

the carbon (C) biogeochemistry and organic matter (OM) storage capacity of freshwater wetlands (Craft, 2007; Loomis & Craft, 2010). Wetlands store an estimated 45–70% of all terrestrial C (Mitra *et al.*, 2005), making them important targets for conservation and major players in the global C cycle (Mcleod *et al.*, 2011). One of the reasons for the high C sequestration rate of wetlands is that decomposition slows in water-saturated anaerobic soils (Reddy & DeLaune, 2008). Microbial decomposition of soil organic C and plant detritus begins with extracellular enzyme-mediated hydrolysis of complex substrates into monomers and oligomers that can be directly used for metabolism (Shi, 2011). This enzymatic hydrolysis has been proposed by many researchers to regulate decomposition rates (Sinsabaugh *et al.*, 1991; Sinsabaugh & Moorhead, 1994; Schimel & Weintraub, 2003; Freeman *et al.*, 2004; Allison & Vitousek, 2005) and has been tied to rates of soil respiration in multiple ecosystems including wetlands (Sinsabaugh & Findlay, 1995; Freeman *et al.*, 1998; Margesin *et al.*, 2000).

Correspondence: Rima B. Franklin, tel. + 804 828 6753, fax + 804 828 0503, e-mail: rbfranklin@vcu.edu

Elevated salinity has been reported to both increase (Weston *et al.*, 2006; Craft, 2007), and decrease decomposition rates in wetland ecosystems (Rejmánková & Houdková, 2006; Roache *et al.*, 2006; Neubauer, 2012). These inconsistencies highlight the need for a more mechanistic understanding of how salinity affects decomposition, which can best be achieved by simultaneously studying extracellular enzymes, the proximal agents of decomposition, as well as the microorganisms responsible for enzyme production. Several prior studies have documented a salinity effect on extracellular enzyme activity (EEA) in both aquatic (Nausch *et al.*, 1998; Cunha *et al.*, 2000; Mulholland *et al.*, 2003; Neubauer *et al.*, 2013) and soil ecosystems (Rietz & Haynes, 2003; Rejmánková & Sirova, 2007). Likewise, an effect of salinity on microbial community structure has been found by scientists considering both environmental gradients (Casamayor *et al.*, 2002; Blum *et al.*, 2004; Crump *et al.*, 2004; Asghar *et al.*, 2012) and experimental manipulations (Langenheder *et al.*, 2003; Mandeel, 2006). However, very few studies have simultaneously studied the effects of salinity on microbial communities and the enzymes they produce (Rietz & Haynes, 2003; Pinckney *et al.*, 2011). This sort of information is essential if we are to understand naturally occurring patterns in soil C across estuarine systems and develop a

predictive understanding of how saltwater intrusion will influence decomposition and C biogeochemistry in wetlands.

This study provides insight into the potential consequences of saltwater intrusion into freshwater ecosystems by investigating changes in microbial decomposition processes and soil OM content along a natural salinity gradient. Specifically, this work examines variation in bacterial community structure and EEA in tidal wetlands ranging from fresh to oligohaline in the Chesapeake Bay watershed (Virginia). In an attempt to isolate the effect of salinity, sampling locations were close enough in proximity that weather, land use, tidal influence, and underlying lithology were likely similar, and site selection targeted nearly identical plant communities.

Materials and methods

Site description and sampling

Samples were collected during a two week period in June 2010 from eight tidal wetlands in Virginia (Fig. 1), varying in salinity from completely fresh (ca. 0 ppt) to oligohaline (ca. 2 ppt). At each wetland, an area dominated by *Peltandra virginica* (min 75% above-ground biomass) was located and a




Fig. 1 Map of sampling locations along four tidal rivers proximal to the Chesapeake Bay (Virginia). The lowest salinity sites (white symbols, ppt < 0.1) are Walkerton Landing (WL) and James River National Wildlife Refuge (JR). Intermediate salinity sites (gray symbols, 0.1 < ppt < 0.4) are Yarmouth Creek (YC), Blackstump Creek (BC), and Morris Creek (MC). The most saline sites (black symbols, ppt > 0.4) are Gleason Marsh (GM), Sweet Hall Marsh (SH), and College Creek (CC).

10 × 10 m² plot was established. Within each plot, five sampling stations (1 m² each) were randomly selected with the caveat that the minimum separation distance between stations was 3 m. All living plant material found within each 1 m² area was harvested to determine above-ground biomass (AGB, clipped to the soil surface). In addition, a 60 cm³ plastic syringe was modified to form a miniature soil corer and used to collect soil for quantifying below-ground biomass (BGB). Lastly, samples of surface soil (0–10 cm) were placed in airtight plastic bags and returned to the laboratory on ice, at which point each sample was homogenized and subdivided for analysis of soil properties, enzyme activity (ca. 50 g stored at 4 °C), and bacterial community structure (ca. 5 g stored at –20 °C).

Soil properties

In the laboratory, soil was immediately analyzed for pH and redox potential (Hanna Combo pH and ORP probe, Smithfield, RI, USA), OM concentration (% as loss on ignition, 425 °C for 12 h), and C and N content (Perkin Elmer Series II CHNS/O Analyzer 2400; Waltham, MA, USA). In addition, porewater was collected from 5-ml soil samples by centrifugation (3000 g for 15 min), filtered using a 0.45 µm pore size mixed cellulose ester syringe filter, and stored at –20 °C until it could be analyzed for the concentration of chloride via ion chromatography (Dionex ICS-1000, Sunnyvale CA, USA). Salinity (ppt) was then calculated as described by Bianchi (2006).

Plant biomass

The plant material harvested from each station was dried (70 °C for 7 days) to estimate AGB as kg of dry plant material per m². BGB was determined by submerging each relevant soil core in 35 ml of tap water (24 h) prior to straining through a No. 45 metal mesh (U.S.A. Standard Testing Sieve, A.S.T.M. E-11 specification, Fisher Scientific, Waltham MA, USA). Roots were collected by hand, dried (95 °C for 96 h), and BGB was calculated as mg of dry roots per cm³ of soil.

Enzyme activity

Rates of EEA were determined within 1 week of sampling using the fluorimetric and colorimetric microplate assays described in Neubauer *et al.* (2013) and a Synergy 2 plate reader (Biotek, Winooski, VT, USA). Briefly, enzyme activity associated with breakdown of cellulose [β -1,4-glucosidase (BG) and 1,4- β -cellobiosidase (CBH)], hemicellulose [β -D-xylosidase (BX)], and lignin [phenol oxidase (POX)], as well as the release of phosphorus [alkaline phosphatase (AP)] and sulfur [arylsulfatase (AS)] from organic molecules, was measured. All substrates and reagents were obtained from Sigma-Aldrich Co. Ltd (St. Louis, MO, USA). Methylumbelliferone (MUB)-linked assays relied on the following substrates: 4-MUB β -D-glucopyranoside (BG), 4-MUB β -D-cellobioside (CBH), 4-MUB- β -D-xylopyranoside (BX), 4-MUB-phosphate (AP), and 4-MUB-sulfate (AS) and incubations of either 1

(CBH, AP) or 4 h (BG, BX, AS) at 30 °C with gentle agitation. POX activity was measured colorimetrically via the oxidation of l-DOPA after a 30 min incubation using the methods of Sin-sabaugh *et al.* (2003).

Bacterial abundance

Whole-community DNA was extracted from 0.25 g subsamples of soil using the Mo Bio PowerSoil DNA Isolation Kit (Carlsbad, CA, USA) and then stored at –20 °C. DNA purity and concentration were analyzed using a Nanodrop ND-1000 (Thermo Scientific, Wilmington, DE, USA). All DNA extracts and PCR products were verified using agarose gel (1.5%) electrophoresis and ethidium bromide staining.

Bacterial abundance was estimated using the quantitative polymerase chain reaction (qPCR) approach outlined in Fierer *et al.* (2005); specifically, we targeted the 16S rRNA gene using the Eub338 and Eub517 primers and analytical details presented in Morrissey *et al.* (2013). Assays utilized SsoAdvanced SYBR Green qPCR Supermix (BioRad, Hercules, CA, USA) and the BioRad CFX 96 Real-Time System; data were analyzed using Bio-Rad CFX Manager Version 2.1. Samples were run in triplicate and averaged; standard curves were established using genomic DNA from *Escherichia coli* (Strain #11775, ATCC, Manassas, VA, USA; average efficiency = 101%, $r^2 = 0.99$).

Bacterial community structure

Bacterial community structure was analyzed using terminal restriction fragment length polymorphism (T-RFLP) targeting the 16S rRNA gene of bacteria and the protocols described in Morrissey *et al.* (2013). Briefly, PCRs included 1.2 ng of template DNA and the domain-specific primers 27F (labeled with FAM) and 1492R at a concentration of 0.2 µM each. PCR products were digested using Hha1 (New England Biolabs, Ipswich, MA, USA) and detected via capillary electrophoresis using a MegaBACE 1000 DNA Analysis System (Amersham Biosciences, Buckinghamshire, UK). T-RFLP fragments between 70 and 400 base pairs (bp) were analyzed using Fragment Profiler software (Version 1.2; Amersham Biosciences, Buckinghamshire, UK) with a 1-bp size differential and a 40 relative fluorescent unit peak height threshold. The T-RFLP results were recorded as a binary data matrix describing the presence or absence of each terminal restriction fragment in each sample.

Statistical analyses

Shapiro–Wilks tests confirmed that soil pH, redox potential (mV), AGB (kg m^{–2}), and BGB (mg cm^{–3}) data were normally distributed. A natural log (ln) transformation was necessary for salinity (ppt), all enzyme activity measures (µmoles g-OM^{–1} h^{–1}), and bacterial abundance (copies g-OM^{–1}); transformed values were utilized in all subsequent statistical analyses. Site differences were analyzed using one-way analysis of variance (ANOVA; $n = 5$, $df = 39$, $\alpha = 0.05$) with Tukey's HSD for post hoc comparisons. Direct and

partial correlations (Pearson's r , $n = 40$) were used to evaluate relationships between bacterial abundance, enzyme activity, and environmental variables; statistical significance was established using a sequential Holm–Bonferroni adjustment to correct for multiple comparisons (Holm, 1979). All ANOVAs and Pearson's correlations were performed using the JMP statistical software (Version JMP Pro 9.0.2, Cary, NC, USA; Sall, 2005).

The binary matrix of T-RFLP data was converted to a set of Jaccard coefficients, which were used for all statistical analyses that considered bacterial community composition. Visualization of overall similarity between samples was achieved using nonmetric multidimensional scaling (NMDS, 2-dimensional solution) and site differences were analyzed using one-way nonparametric multivariate analysis of variance (NP-MANOVA; $n = 5$, $df = 39$, $\alpha = 0.05$). Mantel and partial Mantel tests (r_M) were performed to compare the changes in bacterial community composition (Jaccard similarity matrix) with the changes in environmental and enzyme variables (Euclidian distance matrices). All community composition analyses were performed in PAST statistical software package (Version 2.10; Hammer, 2001).

Results

Site differences

Sites varied significantly with respect to porewater salinity, from fresh (0.03 ppt at JR) to brackish (1.85 ppt at CC, Table 1). Likewise, soil characteristics (pH, redox, %OM, and C : N) and macrophyte biomass (AGB and BGB) exhibited significant differences between sites, though all parameters were within the natural ranges expected for these types of wetlands (Reddy & DeLaune, 2008; Barendregt *et al.*, 2009).

The activity of all enzymes varied significantly across sites (Table 2, rates in $\mu\text{moles g-OM}^{-1} \text{ h}^{-1}$). For the hydrolytic carbon-degrading enzymes (CBH, BG, and BX), rates were lowest at JR (CBH = 0.13, BX = 0.03) and/or WL (BG = 0.09, BX = 0.03) and highest (ca. 5-times greater) at SH (mean: CBH = 0.75, BG = 0.36, and BX = 0.12). This pattern was also observed for POX [lowest at JR (mean: 31) and highest at SH (mean: 1071); ca. 35-fold increase] and AP [lowest at JR (mean: 0.22) and highest at SH and GM (mean: 1.39); ca. sixfold increase]. Activity of AS also varied ~sixfold from the site with the lowest activity (BC=0.08) to the site with the highest activity (YC = 0.45).

Bacterial abundance differed significantly across sites (Table 1), ranging from the lowest at BC (8.3×10^9 copies g-OM^{-1}) to the highest at GM (35.3×10^9 copies g-OM^{-1} , ca. 4-times greater). The community structure of bacteria also differed across sites (NP-MANOVA $F = 7.32$, $P < 0.01$); all pairwise comparisons between sites yielded $P < 0.03$.

Table 1 Site locations and mean (\pm SE) of environmental parameters for each site. Values with the same superscript letter are not significantly different from each other (ANOVA and Tukey's HSD post hoc test with $\alpha = 0.05$; salinity and bacterial abundance were natural log (ln) transformed values prior to testing)

Site	GPS (N)	GPS (W)	Porewater salinity (ppt)	Soil properties				Plant biomass		Bacterial abundance ($\times 10^9$ copies g-OM^{-1})
				pH	Redox (mV)	OM (%)	C : N	AGB* (kg m^{-2})	BGB† (mg cm^{-3})	
JR	37°16'27"	77°09'18"	0.03 \pm 0.01 ^a	6.1 \pm 0.1 ^{ab}	−11 \pm 5 ^b	37.4 \pm 2.9 ^a	12.7 \pm 0.6 ^{ab}	0.23 \pm 0.04 ^b	4.8 \pm 1.1 ^b	13.5 \pm 2.6 ^{ab}
WL	37°43'60"	77°00'96"	0.04 \pm 0.01 ^a	5.6 \pm 0.2 ^b	309 \pm 26 ^a	35.8 \pm 1.7 ^a	12.6 \pm 0.5 ^{ab}	0.19 \pm 0.04 ^b	4.4 \pm 1.8 ^b	12.4 \pm 2.6 ^{ab}
YC	37°19'64"	76°52'26"	0.15 \pm 0.03 ^b	6.4 \pm 0.1 ^a	−96 \pm 9 ^b	10.4 \pm 2.0 ^b	11.4 \pm 0.2 ^{ab}	0.16 \pm 0.02 ^a	23.8 \pm 3.4 ^b	23.7 \pm 2.8 ^{bc}
MC	37°16'78"	76°53'38"	0.16 \pm 0.01 ^b	6.2 \pm 0.2 ^{ab}	−65 \pm 28 ^b	18.2 \pm 0.6 ^b	11.5 \pm 0.3 ^{ab}	0.12 \pm 0.03 ^b	6.34 \pm 0.3 ^b	18.9 \pm 2.8 ^{abc}
BC	37°18'76"	76°51'89"	0.17 \pm 0.02 ^b	6.2 \pm 0.1 ^{ab}	23 \pm 19 ^b	34.2 \pm 5.0 ^a	12.9 \pm 0.6 ^a	0.11 \pm 0.03 ^a	30.0 \pm 3.4 ^b	8.3 \pm 2.1 ^a
GM	37°38'14"	76°51'39"	0.54 \pm 0.04 ^c	5.6 \pm 0.1 ^a	255 \pm 58 ^a	15.3 \pm 0.4 ^b	10.8 \pm 0.4 ^b	0.53 \pm 0.12 ^b	1.0 \pm 0.2 ^a	35.3 \pm 8.9 ^c
SH	37°33'02"	76°53'31"	0.88 \pm 0.08 ^{cd}	6.2 \pm 0.2 ^{ab}	−64 \pm 31 ^b	14.2 \pm 0.6 ^b	11.3 \pm 0.3 ^{ab}	0.14 \pm 0.05 ^b	6.1 \pm 2.2 ^b	28.2 \pm 4.3 ^{bc}
CC	37°15'08"	76°42'60"	1.85 \pm 0.11 ^d	6.4 \pm 0.2 ^a	−73 \pm 22 ^b	16.4 \pm 0.7 ^b	10.8 \pm 0.7 ^b	0.20 \pm 0.04 ^b	9.1 \pm 2.7 ^b	28.2 \pm 4.6 ^{bc}

*Above-ground biomass.

†Below-ground biomass.

Table 2 Mean (\pm SE) enzyme activity rates for each site. Values with the same superscript letter are not significantly different from each other (ANOVA and Tukey's HSD post hoc test on natural log (ln) transformed values, $\alpha = 0.05$)

Site	Enzyme Activity ($\mu\text{moles g-OM}^{-1} \text{ h}^{-1}$)					
	CBH	BG	BX	POX	AP	AS
JR	0.13 ± 0.01^a	0.11 ± 0.02^{ab}	0.03 ± 0.01^a	31 ± 9^a	0.22 ± 0.03^a	0.10 ± 0.02^{ab}
WL	0.15 ± 0.01^a	0.09 ± 0.01^a	0.03 ± 0.00^a	100 ± 20^b	0.31 ± 0.03^a	0.09 ± 0.01^{ab}
YC	0.46 ± 0.01^{bc}	0.28 ± 0.04^c	0.10 ± 0.01^b	365 ± 26^c	0.87 ± 0.04^b	0.45 ± 0.00^c
MC	0.41 ± 0.03^b	0.18 ± 0.03^{abc}	0.09 ± 0.01^b	222 ± 39^{bc}	0.84 ± 0.07^b	0.24 ± 0.04^{bc}
BC	0.20 ± 0.05^a	0.25 ± 0.06^{bc}	0.07 ± 0.01^b	295 ± 101^b	0.36 ± 0.09^a	0.08 ± 0.04^a
GM	0.61 ± 0.03^{bc}	0.26 ± 0.03^c	0.09 ± 0.01^b	501 ± 74^{cd}	1.39 ± 0.05^b	0.34 ± 0.02^c
SH	0.75 ± 0.08^c	0.36 ± 0.05^c	0.12 ± 0.02^b	1071 ± 238^d	1.39 ± 0.13^b	0.40 ± 0.04^c
CC	0.50 ± 0.06^{bc}	0.23 ± 0.02^{bc}	0.09 ± 0.01^b	250 ± 39^b	1.10 ± 0.12^b	0.36 ± 0.04^c

Correlation analysis

Salinity. Salinity was negatively correlated with soil OM content ($r = -0.65$, $P < 0.01$) and C:N ($r = -0.50$, $P < 0.01$, Fig. 2), but unrelated to the other environmental variables (all $|r| < 0.29$, $P > 0.07$). Salinity exhibited strong positive correlations with EEA (Fig. 3; all $r > 0.58$, all $P < 0.01$) and bacterial abundance (Fig. 4; $r = 0.56$, $P < 0.01$). Similarly, salinity was the strongest environmental correlate with bacteria community structure ($r_M = 0.32$, $P < 0.01$). This relationship can be visualized on the lower panel of Fig. 4 where increasing salinity is associated with a positive shift on the NMDS Axis 1, and a negative shift on Axis 2. This is reflected by a salinity vector that represents the strength and direction of the correlation of salinity with each NMDS axis.

Soil OM. In addition to the relationship with salinity, soil OM was strongly correlated with C:N ratio ($r = 0.67$, $P < 0.01$), but unrelated to the other environmental parameters (all $|r| < 0.26$, $P > 0.09$). Furthermore, OM was negatively correlated with enzyme activity rates (Pearson's r : CBH = -0.91 , BG = -0.65 , BX = -0.80 , POX = -0.73 , AP = -0.90 , AS = -0.91 ; all $P < 0.01$) as well as bacterial abundance ($r = -0.64$, $P < 0.01$). Changes in soil OM were not significantly related to changes in bacterial community composition following the Holm's Bonferroni correction ($r_M = 0.13$, $P = 0.04$).

Other environmental variables. The only other significant relationships among environmental variables were between redox and pH ($r = -0.73$, $P < 0.01$) and between AGB and both pH ($r = -0.47$, $P < 0.01$) and redox ($r = 0.52$, $P < 0.01$). In addition to salinity and OM (discussed above), EEA was correlated with C:N (Pearson's r : CBH = -0.65 , BG = -0.42 , BX = -0.46 , POX = -0.48 , AP = -0.71 , AS = -0.74 ; all $P < 0.01$),

but unrelated to all other measured environmental variables. The only other significant correlation for bacterial abundance was with AGB ($r = 0.46$, $P < 0.01$). Bacteria community structure was unrelated to all other environmental variables (all $r_M < 0.12$ and $P > 0.05$). Full correlation results are presented in Table S1.

Partial correlation analysis

To determine the extent to which salinity was a direct driver of EEA and the bacterial community, a partial correlation analysis was performed to control for the main environmental covariates (%OM and C:N; Table 3). All enzymes remained significantly correlated with salinity after controlling for C:N ($r > 0.35$, $P < 0.04$); similarly, relationships remained significant after controlling for OM ($r > 0.32$, $P < 0.05$) except for AS rates. Bacterial abundance remained significantly correlated with salinity after controlling for C:N, but not OM. The association between salinity and community structure was robust and remained highly significant after controlling for covariates with salinity (both $r_M > 0.29$ with $P < 0.01$).

Relationships between the bacterial community and EEA were also examined. Bacterial abundance was significantly correlated with all enzymes except BG, and three of these relationships remained significant after controlling for covariation with salinity via partial correlation (Table 4). Furthermore, the composition of the bacterial community was significantly related to CBH, POX, and AP, even following a partial Mantel test to account for any shared relationship with salinity.

Discussion

The aim of this study was to examine changes in bacterial community structure and function (EEA) along a salinity gradient from fresh to oligohaline to better understand how salinity regulates soil OM




Fig. 2 Variation in (a) organic matter and (b) C : N with salinity [natural log (ln) scale]. Correlation results are shown in the lower right corner (Pearson's r , both $P < 0.01$). Data points are colored by salinity (white < 0.1 ppt, gray $= 0.1 < \text{ppt} < 0.4$, and black > 0.4 ppt).

decomposition and storage in wetlands. The role of salinity as a driver of ecosystem processes in tidal freshwater wetlands is particularly important in light of the potential for climate change-associated saltwater intrusion. Generally, there have been few studies that focus on this range of salinity (Poffenbarger *et al.*, 2011; Neubauer, 2012; Sharpe & Baldwin, 2013), even though the transition from fresh to oligohaline is what many historically freshwater systems are likely to experience as a consequence of sea level rise in the coming decades (Woodroffe & Murray-Wallace, 2012).

The negative relationship between soil OM and salinity exhibited in the current work is consistent with Craft (2007), who found soil organic C (%) to be negatively related to salinity in tidal wetlands across the conterminous United States. Specifically, for the 63 studies surveyed in Craft's work, soil bulk density and organic C content were consistently lower in saltier marshes than their freshwater counterparts regardless of geographical region. We consider three nonexclusive possible explanations for this pattern: (i) enhanced tidal influences that increase OM export and/or sediment deposition in saltier marshes; (ii) decreased plant production and/or altered community composition, which changes the quality and quantity of OM deposited on the marsh surface; and (iii) increased decomposition rates at more saline sites.

Tidal range has been shown to be a strong regulator of aqueous export of total organic C and suspended solids from marsh ecosystems (Childers *et al.*, 2000) and could potentially affect mineral sediment deposition (Chmura & Hung, 2004). Consequently, a greater tidal range at the more saline wetlands could contribute to the lower levels of soil OM at these sites. Although not directly measured in this study, we expect the differences in tidal range across the sites we sampled were not sufficient to explain the OM patterns. For instance, tidal range predictions by NOAA on the James River (20 year average from <http://tidesandcurrents.noaa.gov>) indicate that the mean tidal range near our most saline site is only a few inches greater than for the most upriver freshwater site (2.26 vs. 2.15 ft). In addition, soil texture (determined by the hydrometer method using one aggregate sample per site; results not presented) did not correlate with salinity (considering % sand, silt, and clay, all $P > 0.29$), which suggests no consistent changes in sediment deposition along the salinity gradient of our sites.

The second explanation of the salinity-OM relationship we observed – that is, the effect of saltwater on plant production and/or community composition – was also not well supported. We saw no correlation between above- or below-ground plant biomass and either salinity or soil OM content, and plant production differed little across sites (Table 1). This is likely due to the relatively narrow salinity range we considered, as prior work has shown decreased plant productivity (McKee & Mendelssohn, 1989; Więski *et al.*, 2010) as well as differences in below-ground biomass allocation (Neubauer *et al.*, 2005) along broader salinity gradients. Aside from biomass production, salinity could influence plant community composition and diversity (Latham *et al.*, 1994; Więski *et al.*, 2010; Tuxen *et al.*, 2011). Because various plant species differ in polymer composition (Kögel-Knabner, 2002), this could affect the




Fig. 3 Variation in extracellular enzyme activity (rates in $\mu\text{moles g-OM}^{-1} \text{h}^{-1}$) with salinity; both parameters are presented using a natural log (ln) scale. Correlations results are shown in the lower right corner of each panel (Pearson's r , all $P < 0.01$). Data points are colored by salinity (white < 0.1 ppt, gray $= 0.1 < \text{ppt} < 0.4$, and black > 0.4 ppt). Enzyme abbreviations are as follows: (a) CBH = 1, 4- β -cellobiosidase, (b) BG = β -1, 4-glucosidase, (c) BX = β -D-xylosidase, (d) AS = arylsulfatase, (e) POX = phenol oxidase, and (f) AP = alkaline phosphatase.

quality of the litter deposited on the marsh surface and potentially alter rates of OM accumulation. While this possibility cannot be completely ruled out in our study, a large variation in litter quality is not expected due to the fact that site selection targeted plots that were dominated by a single perennial species (*Peltandra virginica*).

Given the similar tidal regimes and plant communities at each of our sites, the third of our possible explanations for the salinity-OM relationship – increased decomposition under more saline conditions – is best supported by our results. In particular, we documented strong relationships between salinity and enzyme activity (Fig. 3), soil OM content (Fig. 2), and microbial community structure (Fig. 4) that typify enhanced decomposition. This could result from a combination of factors whereby salinity: (i) increases the bioavailability of organic substrates; (ii) facilitates enzyme activity;

and (iii) stimulates changes in microbial community structure. Furthermore, we propose that salinity's influence in this study derives largely from the effect of increased ionic strength on the molecular stability and sorption of both enzymes and their organic substrates. With regard to OM, increased ionic strength can make organic particles more accessible by disrupting soil microaggregates (Rengasamy & Sumner, 1998) and increasing the number of macropore spaces in the soil profile (Brady & Weil, 2004) potentially affecting soil aeration (Blackwell *et al.*, 1990; Kühne *et al.*, 2012). Ionic strength could also affect solute concentrations of organic compounds by decreasing their sorption to soil particles (Reemtsma *et al.*, 1999; Mavi & Marschner, 2012). Both these processes would increase microbial access to organic substrates and, as past studies have found enzyme activity rates to be positively related to




Fig. 4 Variation in (a) bacterial abundance and (b) community composition with salinity. Bacterial abundance and salinity axes use a natural log (ln) scale. Community composition is displayed using an NMDS ordination diagram (stress = 0.20) wherein the vector presents the relationship of salinity with each axis. Correlation results for salinity with abundance (Pearson's r) and composition (Mantel test, r_M) are shown in the lower right corner. Data points are colored by salinity (white < 0.1 ppt, gray = 0.1 < ppt < 0.4, and black > 0.4 ppt).

target molecule availability (Shackle *et al.*, 2000; Allison & Vitousek, 2005; Morrissey *et al.*, 2013), this could explain the higher levels of enzyme activity we observed at more saline sites (Fig. 3). Similarly, the abundance of cations in seawater can affect inorganic N exchange and P sorption (Portnoy & Giblin, 1997;

Table 3 Partial correlation analysis comparing salinity to enzyme activity (Pearson's r), bacterial abundance (Pearson's r), and bacterial community composition (Mantel test, r_M) while controlling for soil OM and C : N

	Partial correlation controlling for			
	OM		C : N	
	r	P	r	P
Enzyme				
CBH	0.52	<0.01*	0.65	<0.01*
BG	0.33	0.04*	0.51	<0.01*
BX	0.44	<0.01*	0.64	<0.01*
POX	0.38	0.02*	0.57	<0.01*
AP	0.55	<0.01*	0.68	<0.01*
AS	-0.05	0.74	0.36	0.03*
Bacteria				
Abundance	0.26	0.11	0.48	<0.01*
Composition	0.30	<0.01*	0.32	<0.01*

* $P < 0.05$.

Weston *et al.*, 2006, 2011), allowing higher salinity to liberate otherwise limiting nutrients (Weston *et al.*, 2010). It is important to point out that the electrostatic interactions described above are interactively affected by other environmental parameters (e.g., pH) and soil physicochemical properties like surface area. For example, increased pH can limit the binding capacity of clay compounds, leading to decreased humic acid sorption (Vermeer *et al.*, 1998; Abate & Masini, 2003). All these effects are also somewhat dependent on the chemical nature of the C compounds involved. In particular, the behavior of humic substances, which constitutes 50–80% of the natural OM (Shaker *et al.*, 2012), will be governed by the relative abundance of functional groups like carboxyls, hydroxyls, and phenolics.

Besides ionic strength effects on organic substrate availability, it is also necessary to consider how changes in salt concentration may directly impact extracellular enzymes. With regard to sorption, several studies have documented a decrease in mobility and catalytic activity when enzymes are adsorbed to clay surfaces (Quiquampoix, 1987; Fusi *et al.*, 1989; Tietjen & Wetzel, 2003), and such immobilization reduces the probability of contact between enzyme and substrate (Lammiro *et al.*, 2010). However, scientists' understanding of how these proteins interact with mineral surfaces is limited, especially when considering confounding factors such as solution pH and enzyme molecular weight. As it appears that the majority of extracellular enzymes are adsorbed to soil particles (Kandeler, 1990; Gianfreda & Bollag, 1994; Violante *et al.*, 1995; Lipson & Nasholm, 2001; Nannipieri *et al.*, 2003), this is an essential area for

Table 4 Direct and partial (controlling for salinity) correlation analysis comparing enzyme activity to bacterial abundance (Pearson's r) and community composition (Mantel test, r_M)

Enzyme	Bacterial abundance				Bacteria Composition			
	Direct		Partial		Direct		Partial	
	r	P	r	P	r_M	P	r_M	P
CBH	0.62	<0.01*	0.34	0.03*	0.26	<0.01*	0.14	0.02*
BG	0.31	0.05	-0.05	0.75	0.09	0.09	-0.01	0.52
BX	0.44	<0.01*	0.06	0.73	0.11	0.06	-0.04	0.73
POX	0.39	0.01*	<0.01	0.98	0.42	<0.01*	0.35	<0.01*
AP	0.63	<0.01*	0.36	0.02*	0.27	<0.01*	0.14	0.02*
AS	0.60	<0.01*	0.41	0.01*	0.10	0.08	0.03	0.25

* $P < 0.05$.

future research if we are to develop a predictive understanding of how C mineralization is affected by saltwater intrusion.

Changes in salinity also have the potential to impact extracellular enzyme activity by influencing molecular stability and protein confirmation states. Most laboratory studies show a detrimental effect of salt on enzyme activity, but these assays use purified enzymes in solution and only a select group of substrates (Das *et al.*, 1997; Fang *et al.*, 2010). Work in soils similarly shows high salt concentrations inhibit enzyme activity, but these studies target hypersaline conditions well beyond the range relevant to our study (e.g., Tripathi *et al.*, 2007; Yun *et al.*, 2010; Pan *et al.*, 2013). The limited reports from wetland soils are inconsistent. Even within a single study, results vary by enzyme, and there is evidence that activity can simultaneously be enhanced, suppressed, and unaffected by salinity (Wu *et al.*, 2008; Jackson & Vallaire, 2009; Saviozzi *et al.*, 2011; Chambers *et al.*, 2013; Neubauer *et al.*, 2013). There are several noteworthy differences between the studies cited above and our own, in which we found salinity significantly enhanced enzyme activity for a diversity of substrates (e.g., breakdown of both labile and recalcitrant compounds, acquisition of C, N, and P). First, the research efforts differed dramatically in their potential for plant effects. Our experimental design minimized the influence of salinity-induced differences in plant biomass and community composition, which was a significant covariant in much of the prior work. Second, our study compared sites with relatively modest salinity (0–2 ppt); this contrasts, for example, the work of Chambers *et al.* (2013) that compared 'fresh' (0.5 ppt), 'brackish' (13 ppt), and 'saline' (26 ppt) conditions. Future studies need to consider a broad range of exposure levels as ecosystem responses to salinity are highly nonlinear. Third, our study represented field conditions integrating prolonged exposure to varying salinity levels.

Studies like Jackson & Vallaire (2009) assessed the response to short-term (55 days) increases in laboratory microcosms. In this context, our findings suggest that long-term shifts in C cycling may not be consistent with short-term disturbance-type responses, as hypothesized by Neubauer *et al.* (2013), and highlight a need for *in situ* work combined with manipulative studies.

Combined, these lines of reasoning help explain the strong correlation we observed between salinity and enzyme activity and provide several possible mechanisms by which salt water intrusion could directly influence decomposition rates. Salinity effects could also be indirect, and mediated through changes in microbial community structure. In this study, we observed a consistent shift in bacterial abundance and community composition with increased salinity (Fig. 4) and found these parameters to be correlated with the activity of several extracellular enzymes (Table 4). This could indicate that community structure (i.e., abundance and composition) regulates, at least in part, organic polymer breakdown and thus could constrain decomposition rates. Several prior studies have similarly documented a link between extracellular enzyme activity and changes in composition of the soil microbial community (e.g., Gallo *et al.*, 2004; Costa *et al.*, 2007; Morrissey *et al.*, 2013). The shifts in bacterial community structure we observed may be in part due to the physiological effects of increased saltwater, and numerous recent studies have demonstrated a link between salinity and microbial community composition (Bouvier & del Giorgio, 2002; Langenheder *et al.*, 2003; Langenheder & Ragnarsson, 2007; Berga *et al.*, 2012). Some researchers have even reported that salinity is the most important factor determining the distribution patterns of microorganisms across the globe (Lozupone & Knight, 2007; Auguet *et al.*, 2010). Besides ionic strength, saltwater intrusion brings with it specific ions that may affect microbial functional groups. For

instance, elevated sulfate can stimulate the growth of sulfate-reducing bacteria, which then outcompete the methanogens typically found in freshwater wetlands (Weston *et al.*, 2006; Chambers *et al.* 2011; Weston *et al.*, 2011).

In conclusion, salinity was found to be strongly related to bacterial community structure and decomposition activity in tidal wetlands ranging from fresh to oligohaline (Figs 3 and 4). The results of this study suggest that via abiotic and/or microbial processes, salinity is stimulating extracellular enzyme activity. This increased enzyme activity could reflect increased decomposition rates, contributing to the negative relationship between salinity and soil OM content in these ecosystems (Fig. 2). If saltwater intrusion does increase soil OM decomposition; previously freshwater wetlands may face reduced soil OM accumulation leading to lower accretion rates (Callaway *et al.*, 1997; Craft, 2007). Consequently, these ecosystems may have more difficulty keeping pace with sea level rise than previously anticipated, potentially leading to ecosystem loss and large releases of stored C (DeLaune & White, 2011). More research on the mechanisms underlying salinity's regulation of enzyme activity and decomposition is sorely needed if we are to quantitatively predict salinity-induced changes in C cycling in tidal wetlands.

Acknowledgements

This work was funded in part by VCU Howard Hughes Medical Institute Summer Scholars Program awards to Morina and Gillespie. Thanks to Amy Jenkins-Lederer, Caitlin Muse, Aaron Porter, Jamie Bourne, John Furry, Ed Crawford, Ariana Johns, and Susan Chong for field and laboratory assistance. We would like to acknowledge Cyrus Brame and Merry Maxwell of the US Fish and Wildlife Service, Bob Greenlee of the Virginia Department of Game and Inland Fisheries, and Jim Perry of the Virginia Institute for Marine Sciences for helping us with site access. Special thanks to Christine Prasse for assistance in enzyme methodologies and David Berrier whose constructive comments improved this manuscript.

References

Abate G, Masini JC (2003) Influence of pH and ionic strength on removal processes of a sedimentary humic acid in a suspension of vermiculite. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, **226**, 25–34.

Allison SD, Vitousek PM (2005) Responses of extracellular enzymes to simple and complex nutrient inputs. *Soil Biology and Biochemistry*, **37**, 937–944.

Asghar HN, Setia R, Marschner P (2012) Community composition and activity of microbes from saline soils and non-saline soils respond similarly to changes in salinity. *Soil Biology and Biochemistry*, **47**, 175–178.

Auguet JC, Barberan A, Casamayor EO (2010) Global ecological patterns in uncultured Archaea. *The ISME Journal*, **4**, 182–190.

Baldwin AH, Mendelsohn IA (1998) Effects of salinity and water level on coastal marshes: an experimental test of disturbance as a catalyst for vegetation change. *Aquatic Botany*, **61**, 255–268.

Barendregt A, Whigham DF, Baldwin AH (2009) *Tidal Freshwater Wetlands*. Margraf Publishers, Weikersheim.

Berga M, Székely AJ, Langenheder S (2012) Effects of disturbance intensity and frequency on bacterial community composition and function. *PLoS ONE*, **7**, e36959.

Bianchi TS (2006) *Biochemistry of Estuaries*. Oxford University Press, North Carolina.

Blackwell PS, Ringrose-Voase AJ, Jayawardane NS, Olsson KA, McKenzie DC, Mason WK (1990) The use of air-filled porosity and intrinsic permeability to air to characterize structure of macropore space and saturated hydraulic conductivity of clay soils. *European Journal of Soil Science*, **41**, 215–228.

Blum LK, Roberts MS, Garland JL, Mills AL (2004) Distribution of microbial communities associated with the dominant high marsh plants and sediments of the United States east coast. *Microbial Ecology*, **48**, 375–388.

Bouvier TC, del Giorgio PA (2002) Compositional changes in free-living bacterial communities along a salinity gradient in two temperate estuaries. *Limnology and Oceanography*, **47**, 453–470.

Brady NC, Weil RR (2004) *Elements of the Nature and Properties of Soils*, 2nd edn, Pearson Prentice Hall, New Jersey.

Callaway JC, DeLaune RD, Patrick WH Jr (1997) Sediment accretion rates from four coastal wetlands along the Gulf of Mexico. *Journal of Coastal Research*, **13**, 181–191.

Casamayor EO, Massana R, Benlloch S *et al.* (2002) Changes in archaeal, bacterial and eukaryal assemblages along a salinity gradient by comparison of genetic fingerprinting methods in a multipond solar saltern. *Environmental Microbiology*, **4**, 338–348.

Chambers LG, Reddy KR, Osborne TZ (2011) Short-term response of carbon cycling to salinity pulses in a freshwater wetland. *Soil Science Society of America Journal*, **75**, 2000–2007.

Chambers LG, Osborne TZ, Reddy KR (2013) Effect of salinity-altering pulsing events on soil organic carbon loss along an intertidal wetland gradient: a laboratory experiment. *Biogeochemistry*, **115**, 363–383.

Childers DL, Day JW, McKellar HN (2000) Twenty more years of marsh and estuarine flux studies: revisiting Nixon (1980). In: *Concepts and Controversies in Tidal Marsh Ecology* (eds Weinstein MP, Kreeger DQ) pp. 391–424, John Wiley & Sons, New York.

Chmura GL, Hung GA (2004) Controls on salt marsh accretion: a test in salt marshes of Eastern Canada. *Estuaries*, **27**, 70–81.

Church JA, White NJ (2006) A 20th century acceleration in global sea-level rise. *Geophysical Research Letters*, **33**, L01602.

Costa AL, Paixão SM, Caçador I, Carolino M (2007) CLPP and EEA profiles of microbial communities in salt marsh sediments. *Journal of Soils and Sediments*, **7**, 418–425.

Craft C (2007) Freshwater input structures soil properties, vertical accretion, and nutrient accumulation of Georgia and US tidal marshes. *Limnology and Oceanography*, **52**, 1220–1230.

Crump BC, Hopkinson CS, Sogin ML, Hobbie JE (2004) Microbial biogeography along an estuarine salinity gradient: combined influences of bacterial growth and residence time. *Applied and Environmental Microbiology*, **70**, 1494–1505.

Cunha MA, Almeida MA, Alcântara F (2000) Patterns of ectoenzymatic and heterotrophic bacterial activities along a salinity gradient in a shallow tidal estuary. *Marine Ecology Progress Series*, **204**, 1–12.

Das JR, Bhat SG, Gowda LR (1997) Purification and characterization of a polyphenol oxidase from the Kew cultivar of Indian pineapple fruit. *Journal of Agricultural and Food Chemistry*, **45**, 2031–2035.

DeLaune RD, White JR (2011) Will coastal wetlands continue to sequester carbon in response to an increase in global sea level?: a case study of the rapidly subsiding Mississippi river deltaic plain. *Climate Change*, **110**, 297–314.

Fang Z, Fang W, Liu J *et al.* (2010) Cloning and characterization of a beta-glucosidase from marine microbial metagenome with excellent glucose tolerance. *Journal of Microbiology and Biotechnology*, **20**, 1351–1358.

Fierer N, Jackson JA, Vilgalys R, Jackson RB (2005) Assessment of soil microbial community structure by use of taxon-specific quantitative PCR assays. *Applied and Environmental Microbiology*, **71**, 4117–4120.

Freeman C, Nevison GB, Hughes S, Reynolds B, Hudson J (1998) Enzymic involvement in the biogeochemical responses of a Welsh peatland to a rainfall enhancement manipulation. *Biology and Fertility of Soils*, **27**, 173–178.

Freeman C, Ostle NJ, Fenner N, Kang H (2004) A regulatory role for phenol oxidase during decomposition in peatlands. *Soil Biology and Biochemistry*, **36**, 1663–1667.

Fusi P, Ristori GG, Calamai L, Stotzy G (1989) Adsorption and binding of protein on “clean” (homoionic) and “dirty” (coated with Fe oxyhydroxides) montmorillonite, illite and kaolinite. *Soil Biology and Biochemistry*, **21**, 911–920.

Gallo M, Amonette R, Lauber C, Sinsabaugh RL, Zak DR (2004) Microbial community structure and oxidative enzyme activity in nitrogen-amended north temperate forest soils. *Microbial Ecology*, **48**, 218–229.

Gianfreda L, Bollag JM (1994) Effect of soils on the behavior of immobilized enzymes. *Soil Science Society of America Journal*, **58**, 1672–1681.

Gleick PH (2003) Water use. *Annual Review of Environment and Resources*, **28**, 275–314.

- Hammer Ø (2001) PAST: Paleontological statistics software package for education and data analysis. *Palaentologica Electronica*, **4**, 1–9.
- Holm S (1979) A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, **6**, 65–70.
- Jackson CR, Vallaire SC (2009) Effects of salinity and nutrients on microbial assemblages in Louisiana wetland sediments. *Wetlands*, **29**, 277–287.
- Kandeler E (1990) Characterization of free and adsorbed phosphatases in soils. *Biology and Fertility of Soils*, **9**, 199–202.
- Kögel-Knabner I (2002) The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter. *Soil Biology and Biochemistry*, **34**, 139–162.
- Kühne A, Schack-Kirchner H, Hildebrand EE (2012) Gas diffusivity in soils compared to ideal isotropic porous media. *Journal of Plant Nutrition and Soil Science*, **175**, 34–45.
- Lammirato C, Miltner A, Wick LY, Kästner M (2010) Hydrolysis of cellobiose by β -glucosidase in the presence of soil minerals—Interactions at solid–liquid interfaces and effects on enzyme activity levels. *Soil Biology and Biochemistry*, **42**, 2203–2210.
- Langenheder S, Ragnarsson H (2007) The role of environmental and spatial factors for the composition of aquatic bacterial communities. *Ecology*, **88**, 2154–2161.
- Langenheder S, Kisand V, Wikner J, Tranvik LJ (2003) Salinity as a structuring factor for the composition and performance of bacterioplankton degrading riverine DOC. *FEMS Microbiology Ecology*, **45**, 189–202.
- Latham PJ, Pearlstone LG, Kitchens WM (1994) Species association changes across a gradient of freshwater, oligohaline, and mesohaline tidal marshes along the lower Savannah River. *Wetlands*, **14**, 174–183.
- Lipson D, Nasholm T (2001) The unexpected versatility of plants: organic nitrogen use and availability in terrestrial ecosystems. *Oecologia*, **128**, 305–316.
- Loomis MJ, Craft CB (2010) Carbon sequestration and nutrient (nitrogen, phosphorus) accumulation in river-dominated tidal marshes, Georgia, USA. *Soil Science Society of America Journal*, **74**, 1028–1036.
- Lozupone CA, Knight R (2007) Global patterns in bacterial diversity. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 11436–11440.
- Mandee QA (2006) Biodiversity of the genus *Fusarium* in saline soil habitats. *Journal of Basic Microbiology*, **46**, 480–494.
- Margesin R, Zimmerbauer A, Schinner F (2000) Monitoring of bioremediation by soil biological activities. *Chemosphere*, **40**, 339–346.
- Marton JM, Herbert ER, Craft CB (2012) Effects of salinity on denitrification and greenhouse gas production from laboratory-incubated tidal forest soils. *Wetlands*, **32**, 347–357.
- Mavi MS, Marschner P (2012) Drying and wetting in saline and saline-sodic soils - effects on microbial activity, biomass and dissolved organic carbon. *Plant and Soil*, **355**, 51–62.
- McKee KL, Mendelsohn IA (1989) Response of a freshwater marsh plant community to increased salinity and increased water level. *Aquatic Botany*, **34**, 301–316.
- McLeod E, Chmura GL, Bouillon S *et al.* (2011) A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Frontiers in Ecology and the Environment*, **9**, 552–560.
- Milley PCD, Dunne KA, Vecchia AV (2005) Global patterns of trends in streamflow and water availability in a changing climate. *Nature*, **437**, 347–350.
- Mitra S, Wassmann R, Vlek PL (2005) An appraisal of global wetland area and its organic carbon stock. *Current Science*, **88**, 25–35.
- Morrissey EM, Berrier DJ, Neubauer SC, Franklin RB (2013) Using microbial communities and extracellular enzymes to link soil organic matter characteristics to greenhouse gas production in a tidal freshwater wetland. *Biogeochemistry*. Available at: <http://link.springer.com/article/10.1007/s10533-013-9894-5> (accessed 28 October 2013). doi: 10.1007/s10533-013-9894-5.
- Mulholland MR, Lee C, Gilbert PM (2003) Extracellular enzyme activity and uptake of carbon and nitrogen along an estuarine salinity and nutrient gradient. *Marine Ecology Progress Series*, **258**, 3–17.
- Nakada M, Inoue H (2005) Rates and causes of recent global sea-level rise inferred from long tide gauge data records. *Quaternary Science Reviews*, **24**, 1217–1222.
- Nannipieri P, Ascher J, Ceccherini M, Landi L, Pietramellara G, Renella G (2003) Microbial diversity and soil functions. *European Journal of Soil Science*, **54**, 655–670.
- Nausch M, Pollehne F, Kerstan E (1998) Extracellular enzyme activities in relation to hydrodynamics in the Pomeranian Bight (Southern Baltic Sea). *Microbial Ecology*, **36**, 251–258.
- Neubauer SC (2012) Ecosystem responses of a tidal freshwater marsh experiencing saltwater intrusion and altered hydrology. *Estuaries and Coasts*, **36**, 491–607.
- Neubauer SC, Givler K, Valentine S, Megonigal JP (2005) Seasonal patterns and plant-mediated controls of subsurface wetland biogeochemistry. *Ecology*, **86**, 3334–3344.
- Neubauer SC, Franklin RB, Berrier DJ (2013) Saltwater intrusion into tidal freshwater marshes alters the biogeochemical processing of organic carbon. *Biogeochemical Discussion*, **10**, 10685–10720.
- Pan C, Liu C, Zhao H, Wang Y (2013) Changes of soil physico-chemical properties and enzyme activities in relation to grassland salinization. *European Journal of Soil Biology*, **55**, 13–19.
- Pinckney JL, Long RA, Paerl HW (2011) Structural and functional responses of microbial mats to reductions in nutrient and salinity stressors in a Bahamian hypersaline lagoon. *Aquatic Microbial Ecology*, **62**, 289–298.
- Poffenbarger HJ, Needelman BA, Megonigal JP (2011) Salinity influence on methane emissions from tidal marshes. *Wetlands*, **31**, 831–842.
- Portnoy JW, Giblin AE (1997) Biogeochemical effects of seawater restoration to diked salt marshes. *Ecological Applications*, **7**, 1054–1063.
- Quiquampoix H (1987) A stepwise approach to the understanding of extracellular enzyme activity in soil I. Effect of electrostatic interactions on the conformation of a β -D-glucosidase adsorbed on different mineral surfaces. *Biochimie*, **69**, 753–763.
- Reddy KR, DeLaune RD (2008) Coastal wetlands: Mississippi River deltaic plain coastal marshes, Louisiana. *Biogeochemistry of Wetlands: Science and Applications*, pp. 669–702. CRC Press, Boca Raton.
- Reemtsma T, Bredow A, Gehring M (1999) The nature and kinetics of organic matter release from soil by salt solutions. *European Journal of Soil Science*, **50**, 53–64.
- Rietz DN, Haynes RJ (2003) Effects of irrigation-induced salinity and sodicity on soil microbial activity. *Soil Biology and Biochemistry*, **35**, 845–854.
- Rejmánková E, Houdková K (2006) Wetland plant decomposition under different nutrient conditions: what is more important, litter quality or site quality? *Biogeochemistry*, **80**, 245–262.
- Rejmánková E, Sirova D (2007) Wetland macrophyte decomposition under different nutrient conditions: relationships between decomposition rate, enzyme activities and microbial biomass. *Soil Biology and Biochemistry*, **39**, 526–538.
- Rengasamy P, Sumner ME (1998) Processes involved in sodic behavior. In: *Sodic Soils: Distribution, Properties, Management, and Environmental Consequences* (eds Sumner ME, Naidu R), pp. 35–50. Oxford University Press, Oxford.
- Roache MC, Bailey PC, Boon PI (2006) Effects of salinity on the decay of the freshwater macrophyte, *Triglochin procerum*. *Aquatic Botany*, **84**, 45–52.
- Sall J (2005) *JMP Start Statistics: A Guide to Statistics and Data Analysis using JMP and JMP IN Software*. SAS Institute Inc, Cary, NC.
- Saviozzi A, Cardelli R, Di Puccio R (2011) Impact of salinity on soil biological activities: a laboratory experiment. *Communications in Soil Science and Plant Analysis*, **42**, 358–367.
- Schimel JP, Weintraub MN (2003) The implication of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biology and Biochemistry*, **35**, 549–563.
- Shackle VJ, Freeman C, Reynolds B (2000) Carbon supply and the regulation of enzyme activity in constructed wetlands. *Soil Biology and Biochemistry*, **32**, 1935–1940.
- Shaker AM, Komy ZR, Heggy SEM, El-Sayed MEA (2012) Kinetic study for adsorption humic acid on soil minerals. *Journal of Physical Chemistry*, **116**, 10889–10896.
- Sharpe PJ, Baldwin AH (2012) Tidal marsh plant community response to sea-level rise: a mesocosm study. *Aquatic Botany*, **101**, 34–40.
- Sharpe PJ, Baldwin AH (2013) Wetland plant species richness across estuarine gradients: the role of environmental factors and the mid-domain effect. *Aquatic Botany*, **107**, 23–32.
- Shi W (2011) Agriculture and ecological significance of soil enzymes: soil carbon sequestration and nutrient cycling. In: *Soil Enzymology* (eds Varma A, Shukla G), pp. 43–60. Springer, New York.
- Sinsabaugh RL, Findlay S (1995) Microbial production, enzyme activity, and carbon turnover in surface sediments of the Hudson River estuary. *Microbial Ecology*, **30**, 127–141.
- Sinsabaugh RL, Moorhead DL (1994) Resource allocation to extracellular enzyme production: a model for nitrogen and phosphorus control of litter decomposition. *Soil Biology and Biochemistry*, **26**, 1305–1311.
- Sinsabaugh RL, Antibus RK, Linkins AE (1991) An enzymatic approach to the analysis of microbial activity during plant litter decomposition. *Agriculture Ecosystems and Environment*, **34**, 43–54.
- Sinsabaugh RL, Saiya-Cork K, Long T, Osgood MP, Neher DA, Zak DR, Norby RJ (2003) Soil microbial activity in a *Liquidambar* plantation unresponsive to CO₂-driven increases in primary production. *Applied Soil Ecology*, **24**, 263–271.
- Smith SJ, Thomson AM, Rosenberg NJ, Izaurralde RC, Brown RA, Wigley TML (2005) Climate change impacts for the conterminous USA: an integrated assessment - Part I. Scenarios and context. *Climatic Change*, **69**, 7–25.
- Tietjen T, Wetzel RG (2003) Extracellular enzyme clay mineral complexes: enzyme adsorption, alteration of enzyme activity, and protection from photodegradation. *Aquatic Ecology*, **37**, 331–339.

- Tripathi S, Chakraborty A, Chakrabarti K, Bandyopadhyay BK (2007) Enzyme activities and microbial biomass in coastal soils of India. *Soil Biology and Biochemistry*, **39**, 2840–2848.
- Tuxen K, Schile L, Stralberg D *et al.* (2011) Mapping changes in tidal wetland vegetation composition and pattern across a salinity gradient using high spatial resolution imagery. *Wetlands Ecology and Management*, **19**, 141–157.
- Vermeer AWP, van Riemsdijk WH, Koopal LK (1998) Adsorption of humic acid to mineral particles. 1. Specific and electrostatic interactions. *Langmuir*, **14**, 2810–2819.
- Violante A, De Cristofaro A, Rao MA, Gianfreda L (1995) Physicochemical properties of protein-smectite and protein-Al (OH)-smectite complexes. *Clay Minerals*, **30**, 325–336.
- Weston NB, Dixon RE, Joye SB (2006) Ramifications of increased salinity in tidal freshwater sediments: geochemistry and microbial pathways of organic matter mineralization. *Journal of Geophysical Research: Biogeosciences*, **111**, 2005–2012.
- Weston NB, Giblin AE, Banta GT, Hopkinson CS, Tucker J (2010) The effects of varying salinity on ammonium exchange in estuarine sediments of the Parker River, Massachusetts. *Estuaries and Coasts*, **33**, 985–1003.
- Weston NB, Vile MA, Neubauer SC, Velinsky DJ (2011) Accelerated microbial organic matter mineralization following salt-water intrusion into tidal freshwater marsh soils. *Biogeochemistry*, **102**, 135–151.
- Wieski K, Guo H, Craft CB, Pennings SC (2010) Ecosystem functions of tidal fresh, brackish, and salt marshes on the Georgia Coast. *Estuaries and Coasts*, **33**, 161–169.
- Wigley TML (2005) The climate change commitment. *Science*, **307**, 1766–1769.
- Woodroffe CD, Murray-Wallace CV (2012) Sea-level rise and coastal change: the past as a guide to the future. *Quaternary Science Reviews*, **54**, 4–11.
- Wu Y, Tam NFY, Wong MH (2008) Effects of salinity on treatment of municipal wastewater by constructed mangrove wetland microcosms. *Marine Pollution Bulletin*, **57**, 727–734.
- Yun EY, Ro HM, Lee GT, Choi WJ (2010) Salinity effects on chlorpyrifos degradation and phosphorus fractionation in reclaimed coastal tideland soils. *Geosciences Journal*, **14**, 371–378.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Correlations coefficients (Pearson's r , top right half of the matrix) and P -values (lower left half) between all evaluated parameters. Enzyme ($\mu\text{moles g-OM}^{-1} \text{ h}^{-1}$) and bacterial abundance (16S rRNA gene copies g-OM^{-1}) data were natural log transformed (\ln) prior to analysis. Bacterial community composition was represented using a Jaccard similarity matrix derived from 16S rRNA T-RFLP; correlations presented below are from associated Mantel tests (r_M). Abbreviations are as follows: CBH, 1, 4- β -cellobiosidase; BG, β -1, 4-glucosidase; BX, β -D-xylosidase; AS, arylsulfatase; POX, phenol oxidase; AP, alkaline phosphatase; BGB, below-ground plant biomass; and AGB, above-ground plant biomass.