

Model-Based Clustering of Time Series in Group-Specific Functional Subspaces

Dani Chu

Simon Fraser University

STAT 843, Feb 11th 2019

Introduction

About the Paper

- ▶ Paper by Charles Bouveyron & Julien Jacques
- ▶ Published in 2011
- ▶ Published in the international, journal Advances in Data Analysis and Classification (ADAC)
- ▶ Cited 22
- ▶ [Link to paper](#)
- ▶ Bouveyron, C. and Jacques, J. (2011). [Model-based clustering of time series in group-specific functional subspaces.](#)
Advances in Data Analysis and Classification, 5(4):281–300

National Football League (NFL) Example

Figure: Example of using model based clustering for multivariate functional data to identify similar shaped routes in the NFL. (Open in Adobe Acrobat)

Table of Contents

1. Introduction
2. Clustering
3. Functional Latent Mixture Model
4. Maximum Likelihood (EM)
5. Examples
6. R Demo

Clustering

Clustering

- ▶ Identify groups of homogeneous data without prior knowledge of these groups
- ▶ Examples: k-means, hierarchical classification, model-based clustering
- ▶ Clustering time series data is difficult since the data lives in an infinite dimensional space

Time Series Clustering

- ▶ Notion of pdfs do not exist for functional data
- ▶ Transform the data first to a finite dimensional problem
- ▶ Clustering time series data is difficult since the data lives in an infinite dimensional space
 - ▶ Discretizing the time interval
 - ▶ Decomposing into basis of functions
 - ▶ FPCA

Time Series Clustering Issues

- ▶ Descretization of the observed curves leads to high-dimensional data sets (sometimes $n < p$).
- ▶ In these cases, model-based clustering suffer from numerical problems.
- ▶ Can assume that high-dimensional data live in group-specific subspaces

Functional Latent Mixture Model

Plan

- ▶ Identify K homogeneous clusters among the data
- ▶ Introduce a family of mixture models designed for multivariate functional data

Reconstructing the Functional Form

- ▶ Let $\{x_1, \dots, x_n\}$ be independent realizations of a L_2 -continuous stochastic process $X = \{X(t)\}$, $t \in [0, T]$
- ▶ We observe discrete observations $x_{i,j} = x_i(t_{i,j})$ at a finite set of ordered times $\{t_{i,j} : j = 1, \dots, m_i\}$
- ▶ Must reconstruct the functional form of the data from the discrete observations
- ▶ Assume curves belong to a finite dimensional space spanned by a basis of functions

Basis Expansion

- ▶ Consider a basis $\{\psi_1, \dots, \psi_p\}$
- ▶ $\gamma = (\gamma_1(X), \dots, \gamma_p(X))$ a random vector in \mathbb{R}^p
- ▶ p is assumed to be fixed and known
- ▶ Assume that the stochastic process X admits the basis expansion

$$X(t) = \sum_{j=1}^p \gamma_j(X) \psi_j(t)$$

Estimation of Basis Expansion

- ▶ Each observed curve is $x_i(t) = \sum_{j=1}^p \gamma_{i,j} \psi_j(t)$
- ▶ Estimated by least square smoothing.
- ▶ A latent mixture model is proposed for modelling the coefficient vectors $\{\gamma_1, \dots, \gamma_n\} \in \mathbb{R}$ of the observed curves
- ▶ $\gamma_i = (\gamma_{i,1}, \dots, \gamma_{i,p})$, for $i = 1, \dots, n$

Assumptions

- ▶ Consider a set of n_k observed curves described by their coefficient vectors $\{\gamma_1, \dots, \gamma_{n_k}\} \in \mathbb{R}$
- ▶ Assume the gammas are independent realizations of a random vector $\Gamma \in \mathbb{R}^p$
- ▶ Actual stochastic process associated with the k th cluster can be described in a low-dimensional functional latent subspace $\mathbb{E}_k[0, T]$ of $L_2[0, T]$ with dimension $d_k \leq p$
- ▶ Let $\mathbb{E}_k[0, T]$ be spanned by the first d_k elements of a group-specific basis of functions $\{\phi_{kj}\}_{j=1, \dots, d_k}$ in $L_2[0, T]$

Assumptions

- ▶ The group-specific basis is obtained from $\{\psi\}_{j=1,\dots,p}$ by a linear transformation $\phi_{kj} = \sum_{l=1}^p q_{k,jl} \psi_l$
- ▶ With an orthogonal $p \times p$ matrix $Q_k = (q_{k,jl})$
- ▶ Split into two parts $[U_k, V_k]$
- ▶ U_k of size $p \times d_k$ with $U_k^t U_k = I_{d_k}$
- ▶ V_k of size $p \times (p - d_k)$ with $V_k^t V_k = I_{p-d_k}$
- ▶ $U_k^t V_k = 0$

Latent Expansion Coefficients

- ▶ $\lambda_1, \dots, \lambda_{n_k}$ is the latent expansion coefficients of the curves in the group-specific basis $\{\phi_{kj}\}_{j=1, \dots, d_k}$
- ▶ Assumed to be independent realizations of a latent random vector $\Lambda \in \mathbb{R}^{d_k}$
- ▶ The relationship between $\{\phi_{kj}\}_{j=1, \dots, d_k}$ and $\{\psi_j\}_{j=1, \dots, p}$ suggest s that the random vectors Γ and Λ are linked through the following linear transformation for the k th group

$$\Gamma = U_k \Lambda + \epsilon$$

- ▶ where $\epsilon \in \mathbb{R}^p$ is an an independent and random noise term

Distributional Assumptions

- ▶ Λ assumed to be distributed according to a multivariate Gaussian density, $\Lambda \sim \mathcal{N}(m_k, S_k)$
- ▶ For the k th group we have m_k the mean and $S_k = \text{diag}(a_{k1}, \dots, a_{kd_k})$ the covariance matrix
- ▶ ϵ distributed according to a multivariate Gaussian density $\epsilon \sim \mathcal{N}(0, \Xi_k)$
- ▶ We then get that for the k th cluster $\Gamma \sim \mathcal{N}(\mu_k, \Sigma_k)$
 - ▶ $\mu_k = U_k m_k$
 - ▶ $\Sigma_k = U_k S_k U_k^t + \Xi_k$

Variances

- ▶ Assume that the noise covariance matrix Ξ_k is such that $\Delta_k = \text{cov}(Q_k^t \Gamma) = Q_k^t \Sigma_k Q_k$ has the following form
 - ▶ A diagonal matrix
 - ▶ $a_{k1} \dots a_{kd_k}$ along the first d_k entries
 - ▶ $b_k \dots b_k$ along the remaining $(p - d_k)$ entries.
 - ▶ $a_{kj} > b_j$ for $j = 1, \dots, d_k$
- ▶ Practically the variance of the actual data of the k th group is modeled by $a_{k1} \dots a_{kd_k}$
- ▶ b_k models the variance of the noise
- ▶ The dimension d_k is the intrinsic dimension of the latent subspace of the k th group.

The Problem Set Up

- ▶ Consider a set of n observed curves $\{x_1, \dots, x_n\}$, where $x_i = \{x_i(t)\}_{t \in [0, T]}$ ($1 \leq i \leq n$)
- ▶ We want to cluster the curves into K homogeneous groups
- ▶ Assume there exists a latent random variable $Z = (Z_1, \dots, Z_K) \in \{0, 1\}^K$ indicating the group membership of X
- ▶ Z_k is equal to 1 if X belongs to the k th group and 0 otherwise
- ▶ Want to predict the value of $z_i = (z_{i1}, \dots, z_{iK})$ of Z for each observed curve x_i

Summary of Model

- ▶ Each x_i is a sample path of X admitting a basis expansion summarised by the coefficient vector γ_i whose distribution is now a mixture of Gaussians with density

$$p(\gamma) = \sum_{k=1}^K \pi_k \phi(\gamma; \mu_k, \Sigma_k)$$

- ▶ where ϕ is the standard Gaussian density function, $\mu_k = U_k m_k$, $\Sigma_k = Q_k \Delta_k Q - k^t$ and $\pi_k = P(Z_k = 1)$ is the prior probability of the k th group
- ▶ This mixture model will be referred to as FLM_[$a_{kj} b_k Q_k d_k$] model

Constraints on the Full Model

- ▶ Can constrain the model parameters within or between groups
- ▶ Constrain the first d_k diagonal elements of Δ_k to be equal within each class ($a_{k1} = \dots = a_{kd_k}$)
 - ▶ Assumes each matrix Δ_k only has two different eigenvalues a_k and b_k .
- ▶ Can fix the parameters b_k to be common across classes.
 - ▶ Assume that the behavior of the error components outside the class specific subspaces is common
 - ▶ modelling the noise outside the latent subspace by b

Submodels

For

- ▶ $\rho = Kp + K - 1$ the number of parameters needed to estimate the means and proportions
- ▶ $\tau = \sum_{k=1}^K d_k [p - (d_k + 1)/2]$ the number of parameters needed to estimate the orientation matrices Q_k and $D = \sum_{k=1}^K d_k$

Here are a selection of submodels and the number of parameters required to estimate the model

FLM Model	Number of Parameters
$[a_{kj}b_kQ_kd_k]$	$\rho + \tau + 2K + D$
$[a_{kj}bQ_kd_k]$	$\rho + \tau + K + D + 1$
$[a_kb_kQ_kd_k]$	$\rho + \tau + 3K$
$[ab_kQ_kd_k]$	$\rho + \tau + 2K + 1$
$[a_kbQ_kd_k]$	$\rho + \tau + 2K + 1$
$[abQ_kd_k]$	$\rho + \tau + K + 2$

Maximum Likelihood (EM)

funHDDC Algorithm

- ▶ In model-based clustering, the estimation of model parameters is traditionally done by maximizing the likelihood through the EM algorithm
- ▶ Iterative algorithm consists in maximizing the complete likelihood rather than directly maximizing the likelihood which is an intractable problem with incomplete data

Log-likelihood

Given the coefficient vectors $\gamma_1, \dots, \gamma_n$ of the observed curves x_1, \dots, x_n the complete log-likelihood of the data under the FLM model has the following form

$$l_c(\theta; \gamma_1, \dots, \gamma_n, z_1, \dots, z_n) = -\frac{1}{2} \sum_{k=1}^K \eta_k \left[\sum_{j=1}^{d_k} \left(\log(a_{kj}) + \frac{q_{kj}^t C_k q_{kj}}{a_{kj}} \right) + \left(\log(b_k) + \frac{q_{kj}^t C_k q_{kj}}{b_k} \right) - 2 \log(\pi_k) \right] + \xi$$

where $\theta = (\pi_k, \mu_k, a_{kj}, b_k, q_{kj})$ for $1 \leq j \leq d_k$ and $1 \leq k \leq K$, q_{kj} is the j th column of Q_k , $C_k = \frac{1}{\eta_k} \sum_{i=1}^n z_{ik} (\gamma_i - \mu_k)^t (\gamma_i - \mu_k)$, $\eta_k = \sum_{i=1}^n z_{ik}$ and ξ is a term not depending on the parameter θ .

E step I

- ▶ At iteration q , compute the expectation of the complete log-likelihood conditionally on the current value of the parameter $\theta^{(q-1)}$
- ▶ So just need to compute $t_{ik}^{(q)} = e[Z_{ik}|\gamma_i, \theta^{(q-1)}]$
- ▶ For the FLM_[$a_k b_k Q_k d_k$] model, the posterior probability $t_{ik}^{(q)}$ can be computed for iteration q .

$$t_{ik}^{(q)} = 1 / \sum_{l=1}^K \exp \left(H_k^{(q-1)}(\gamma_i) - H_l^{(q-1)}(\gamma_i) \right)$$

E step II

With $H_k^{(q-1)}(\gamma)$ defined for $\gamma \in \mathbb{R}^p$ as:

$$H_k^{(q-1)}(\gamma) = \|\mu_k^{(q-1)} - P_k(\gamma)\|_{D_k}^2 + \frac{1}{b_k^{(q-1)} \|\gamma - P_k(\gamma)\|^2} + \sum_{j=1}^{d_k} \log \left(a_{kj}^{(q-1)} \right) + (p - d_k) \log \left(b_k^{(q-1)} \right) - 2 \log \left(\pi_k^{((q-1))} \right)$$

where $\|\cdot\|_{D_k}^2$ is a norm on the latent space \mathbb{E}_k defined by $\|y\|_{D_k}^2 = y^t \mathcal{D}_k y$, $\mathcal{D}_k = \tilde{Q} \Delta_k^{-1} \tilde{Q}^t$ and \tilde{Q} is a $p \times p$ matrix containing the d_k vectors of U_k completed by zeros such as $\tilde{Q} = [U_k, 0_{p-d}]$, P_k is the projection operator on the latent space \mathbb{E}_k defined by $P_k(\gamma) = U_k U_k^t (\gamma - \mu_k) + \mu_k$

E step III

- ▶ $H_k(\gamma)$ is mainly based on two distances
 1. The distance between the projection of γ on \mathbb{E}_k and the current mean of the k th group
 2. The distance between the observation and the subspace \mathbb{E}_k
- ▶ The classification function favors the assignment of a new observation to the class for which it is close to the subspace and for which its projection on the class subspace is close to the mean of the class
- ▶ The variance terms a_k and b_k balance the importance of both distances
- ▶ If the data is noisy (b_k is large) it is natural to balance the distance $\|\gamma - P_k(\gamma)\|^2$ by $1/b_k$

M step I (Mixture Proportions and Means)

- ▶ The next step is to estimate the model parameters by maximizing the expectation of the complete likelihood conditionally on the posterior probabilities $t_{ik}^{(q)}$ computed in the E step.
- ▶ Mixture proportions and means are updated as usual by:

$$\pi_k^{(q)} = \frac{n_k^{(q)}}{n}, \quad \mu_k^{(q)} = \frac{1}{n_k^{(q)}} \sum_{i=1}^n t_{ik}^{(q)} \gamma_i$$

- ▶ where $n_k^{(q)} = \sum_{i=1}^n t_{ik}^{(q)}$

M step II (a_{kj}, b_k, q_{kj})

- ▶ Introduce $C_k^{(q)} = \frac{1}{n_k^{(q)}} \sum_{i=1}^n t_{ik}^{(q)} \left(\gamma_i - \mu_k^{(q)} \right)^t \left(\gamma_i - \mu_k^{(q)} \right)$, the sample covariance matrix of group k
- ▶ Introduce $W = (w_{jk})_{q \leq j, k \leq p} = \int_0^T \psi_j(t) dt$, the matrix of inner products between the basis functions
- ▶ the first d_k columns of Q_k are updated by the eigenvectors associated with the largest eigenvalues of $W^{\frac{1}{2}} C_k^{(q)} W^{\frac{1}{2}}$
- ▶ Variance parameters $a_{kj}, j = 1, \dots, d_k$ are updated by the d_k largest eigenvalues $W^{\frac{1}{2}} C_k^{(q)} W^{\frac{1}{2}}$
- ▶ the variance parameters b_k are updated by
$$b_k^{(q)} = \text{trace} \left(W^{\frac{1}{2}} C_k^{(q)} W^{\frac{1}{2}} \right) - \sum_{j=1}^{d_k} \hat{a}_{kj}^{(q)}$$

Summary of EM Algorithm

- ▶ fun HDDC models and clusters time series objects through their projections in group-specific functional principal subspaces.
- ▶ These group-specific functional principal subspaces are obtained by performing FPCA conditionally on the posterior probabilities t_{ik}
- ▶ No discriminative information is lost since the b_k term models the variance outside the subspaces

Hyper-parameters

- ▶ We do not estimate d_k or K and they cannot be found from maximizing the likelihood since they control the model complexity
- ▶ Can use BIC criterion to select both hyperparameters.

Classifying Observations

- ▶ The last step is to add a classification for each observation
- ▶ We do this using the maximum a posteriori (MAP) rule
- ▶ Assigning an observation $\gamma_i \in \mathbb{R}^P$ to the group for which γ_i has the highest posterior probability $P(Z_{ik} = 1 | \gamma_i)$
- ▶ Assign the observation γ_i to the group with the highest $t_{ik}^{(q_f)}$ where q_f is the last iteration of the algorithm before its convergence.

Convergence

- ▶ Since it is an EM-based algorithm we are guaranteed convergence to a local maximum
- ▶ Can execute the algorithm several times from random initializations to try and find a global maximum

Examples

Example 1 National Basketball Association (NBA)

- ▶ In their paper “Possession Sketches: Mapping NBA Strategies”, [Miller and Bornn, 2017] use this idea to cluster similar movement patterns of NBA players to develop a dictionary of actions that players make throughout a possession
- ▶ They then use this dictionary as a vocabulary to do topic analysis on NBA possession.
- ▶ They are able to then group together possession with similar offensive structure




Figure: Example of Movement Clusters in NBA Possessions

Example 2 Canadian Temperatures

- ▶ The original paper uses an example with the Canadian temperature data provided in the *fda* package
- ▶ Use a basis of 20 natural cubic splines
- ▶ Use BIC to select 4 clusters




Figure: Colors indicating the cluster membership plotted with respect to geographical positions

R Demo

References



Bouveyron, C. and Jacques, J. (2011).

Model-based clustering of time series in group-specific functional subspaces.

Advances in Data Analysis and Classification, 5(4):281–300.



Miller, A. C. and Bornn, L. (2017).

Possession sketches : Mapping nba strategies.

In *Proceedings of the 2017 MIT Sloan Sports Analytics Conference*.