

Notes on running MOM5 ocean/sea-ice simulations forced with JRA55-do

Riccardo Farneti¹

¹rfarneti@ictp.it

Version of February 14, 2023

Abstract. Notes on how to preprocess and prepare JRA55-do surface atmospheric data sets for MOM5 ocean-sea-ice simulations.

1 JRA55-do dataset

5 JRA55-do (T sujino et al., 2018) is a surface dataset for driving ocean-sea ice models and used in phase 2 of OMIP (OMIP-2; Griffies et al., 2016). JRA55-do corrects the atmospheric reanalysis product JRA-55 using satellite and other atmospheric reanalysis products. The merits of JRA55-do are the high horizontal resolution (~55 km) and temporal interval (3 h). JRA55-do can suitably replace the current CORE/OMIP-1 dataset (T sujino et al., 2020).

The JRA55-do datasets (Table 1) are downloaded from input4MIPs and any update from:

10 <https://climate.mri-jma.go.jp/pub/ocean/JRA55-do/>.

Presently, v1.5.0 covers the period 1958 to 2022.

2 Preprocessing of JRA55-do dataset

All forcing files are yearly, from 1958 to 2022. In order to produce a single file for each surface forcing variable we need to:

1. *Padding*: [optional] padding is necessary when using single files as it includes on each file the last and first time step of the previous and next year, respectively. This is done using the routines in

15 `/pad_JRA_main`.

2. *Concatenation*: next we concatenate all years into a single file with the NCO command `ncrcat`. I have found that the CDO commands `mergetime` and `cat` would modify the metadata for variables and axes causing the FATAL error:

“friver / couldnt recognize axis atts in time_interp_external”.

20 3. *Edit the attributes*: the model might crash because `mpp_io_read.inc` returns a FATAL error:

“VAR ATT too long”.

In this case modify the attributes with `nco`:

```
ncatted -h -0 -a history,,d,,name_of_file.nc
```

4. The required surface forcing files are:

- 25 - JRA_psl.1958_2022.nc
- JRA_tas.1958_2022.nc
- JRA_huss.1958_2022.nc
- JRA_uas.1958_2022.nc
- JRA_vas.1958_2022.nc
- 30 - JRA_rlds.1958_2022.nc
- JRA_rsds.1958_2022.nc
- JRA_prsa.1958_2022.nc
- JRA_prsn.1958_2022.nc

2.1 Interpolation of files

35 Some of the forcing fields, and the SSS restoring field, need to be interpolated onto the model grid.

1. SSS: Sea surface salinity is restored to the JRA field:

```
sos_input4MIPs_atmosphericState_OMIP_MRI - JRA55 - do - 1 - 5 - 0_gr_195501 - 2012 - clima.nc
```

(a) SSS is interpolated using the routines in

```
/OM4_025_salt_restore including interp_and_fill.py.
```

40 Necessary input are ocean_hgrid.nc and ocean_mask.nc.

(b) The model might crash because mpp_io_read.inc returns a FATAL error: “VAR ATT too long”.

In this case modify the attributes with nco:

```
ncatted -h -0 -a history,,d,,salt_sfc_restore.nc
```

```
ncatted -h -0 -a comment,,d,,salt_sfc_restore.nc
```

45 (c) The sea surface salinity restoring file, with variable salt is:

```
      - salt_sfc_restore.nc
```

2. *friver and licalvf*: Both liquid and solid runoff are regridded on the model grid.

(a) (optional) First, both variables are padded.

(b) Then, *friver and licalvf* are regridded using the routines in

50 /OM4_025_runoff including regrid_runoff.py.

Necessary input are ocean_hgrid.nc and ocean_mask.nc.

(c) kshedstrom@alaska_edu suggested to modify both variables by remove NaNs (with xarray: `.fillna(0)`) and deleting the `_FillValue` attribute as FMS would not accept it.

I have found this would cause the model to crash with a FATAL error:

55

“friver / couldnt recognize axis atts in time_interp_external”.

(d) Finally, both variables are concatenated into a single file with the NCO command `ncrcat`:

- JRA_friver.1958_2022.nc
- JRA_licalvf.1958_2022.nc

Table 1. JRA-55 based surface dataset for driving ocean/sea-ice models (JRA55-do)

| variable | long name | units | frequency | nominal resolution |
|----------|---|---------------------------------------|-----------|--------------------|
| uas | eastward near-surface (10m) wind speed | [m s ⁻¹] | 3hr | 50 km |
| vas | northward near-surface (10m) wind speed | [m s ⁻¹] | 3hr | 50 km |
| tas | Near-Surface (10 m) Air Temperature | [K] | 3hr | 50 km |
| huss | Near-Surface (10 m) Specific Humidity | | 3hr | 50 km |
| psl | Sea Level Pressure | [Pa] | 3hr | 50 km |
| rsds | Surface Downwelling Shortwave Radiation | [W m ⁻²] | 3hr | 50 km |
| rlds | Surface Downwelling Longwave Radiation | [W m ⁻²] | 3hr | 50 km |
| prra | Rainfall Flux | [Kg m ⁻² s ⁻¹] | 3hr | 50 km |
| prsn | Snowfall Flux | [Kg m ⁻² s ⁻¹] | 3hr | 50 km |
| friver | Water Flux into Sea Water from Rivers (liquid water runoff) | [Kg m ⁻² s ⁻¹] | day | 25 km |
| licalvf | Land ice calving flux (solid ice runoff) | [Kg m ⁻² s ⁻¹] | day | 25 km |

References

- 60 Griffies, S., Biastoch, A., Boning, C., Bryan, F., Danabasoglu, G., Chassignet, E., England, M., Gerdes, R., Haak, H., and Hallberg, R.: Coordinated Ocean-ice Reference Experiments (COREs), *Ocean Modelling*, 26, 1–46, 2009.
- Griffies, S. M., Danabasoglu, G., Durack, P. J., Adcroft, A. J., Balaji, V., Böning, C. W., Chassignet, E. P., Curchitser, E., Deshayes, J., Drange, H., Fox-Kemper, B., Gleckler, P. J., Gregory, J. M., Haak, H., Hallberg, R. W., Heimbach, P., Hewitt, H. T., Holland, D. M., Ilyina, T., Jungclaus, J. H., Komuro, Y., Krasting, J. P., Large, W. G., Marsland, S. J., Masina, S., McDougall, T. J., Nurser, A. J. G., Orr, 65 J. C., Pirani, A., Qiao, F., Stouffer, R. J., Taylor, K. E., Treguier, A. M., Tsujino, H., Uotila, P., Valdivieso, M., Wang, Q., Winton, M., and Yeager, S. G.: OMIP contribution to CMIP6: experimental and diagnostic protocol for the physical component of the Ocean Model Intercomparison Project, *Geoscientific Model Development*, 9, 3231–3296, 2016.
- Tsujino, H., Urakawa, S., Nakano, H., Small, R. J., Kim, W. M., Yeager, S. G., Danabasoglu, G., Suzuki, T., Bamber, J. L., Bentsen, M., Böning, C. W., Bozec, A., Chassignet, E. P., Curchitser, E., Boeira Dias, F., Durack, P. J., Griffies, S. M., Harada, Y., Ilicak, M., Josey, 70 S. A., Kobayashi, C., Kobayashi, S., Komuro, Y., Large, W. G., Le Sommer, J., Marsland, S. J., Masina, S., Scheinert, M., Tomita, H., Valdivieso, M., and Yamazaki, D.: JRA-55 based surface dataset for driving ocean–sea-ice models (JRA55-do), *Ocean Modelling*, 130, 79–139, 2018.
- Tsujino, H., Urakawa, L. S., Griffies, S. M., Danabasoglu, G., Adcroft, A. J., Amaral, A. E., Arsouze, T., Bentsen, M., Bernardello, R., Böning, C. W., Bozec, A., Chassignet, E. P., Danilov, S., Dussin, R., Exarchou, E., Fogli, P. G., Fox-Kemper, B., Guo, C., Ilicak, M., Iovino, 75 D., Kim, W. M., Koldunov, N., Lapin, V., Li, Y., Lin, P., Lindsay, K., Liu, H., Long, M. C., Komuro, Y., Marsland, S. J., Masina, S., Nummelin, A., Rieck, J. K., Ruprich-Robert, Y., Scheinert, M., Sicardi, V., Sidorenko, D., Suzuki, T., Tatebe, H., Wang, Q., Yeager, S. G., and Yu, Z.: Evaluation of global ocean–sea-ice model simulations based on the experimental protocols of the Ocean Model Intercomparison Project phase 2 (OMIP-2), *Geoscientific Model Development*, 13, 3643–3708, 2020.