

regional-mom6: Automatic generation of regional configurations for the Modular Ocean Model v6 in Python

Ashley J. Barnes^{1,2}, Navid C. Constantinou^{1,2,3,4}, Angus H. Gibson¹, Andrew E. Kiss^{1,2}, Chris Chapman^{5*}, John Reilly^{6*}, and Dhruv Bhagtani^{1,2*}

¹ Australian National University, Australia ² ARC Centre of Excellence for Climate Extremes, Australia ³ University of Melbourne, Australia ⁴ ARC Centre of Excellence for the Weather of the 21st Century, Australia ⁵ CSIRO Environment, Hobart, Tasmania, Australia ⁶ University of Tasmania, Australia * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

Summary

regional-mom6 is a Python package that provides an easy and versatile way to set up regional configurations of the Modular Ocean Model version 6 (MOM6).

Regional ocean modeling

In the ocean, fast and small-scale motions (from ~100m to ~100km varying at time scales of hours to days) play an important role in shaping the large-scale ocean circulation and climate (length scales ~10,000km varying at decadal time scales) (de Lavergne et al., 2022; Gula et al., 2022; Melet et al., 2022). Despite the increase in computational power and the use of graphical processing units that can bring breakthrough performance and speedup (Silvestri et al., 2023), there are always processes, boundary, or forcing features that are smaller than the model's grid spacing and, thus, remain unresolved in global ocean models. Regional ocean models can be run at higher resolutions while limiting the required computational resources.

A regional ocean model simulates the ocean only in a prescribed region, which is a subset of the global ocean. To do that, we need to apply open boundary conditions at the region's boundaries, that is, we need to impose conditions that mimic the oceanic flow that we are not simulating (Orlanski, 1976). For example, Figure 1 shows the surface currents from a regional ocean simulation of the Tasman sea that was configured using the regional-mom6 package. The boundaries of the domain depicted in Figure 1 are forced with the ocean flow from a global ocean reanalysis product. Higher-resolution regional ocean models improve the representation of smaller-scale motions, such as tidal beams, mixing, mesoscale and sub-mesoscale circulation, as well as the oceanic response to smaller-scale bathymetric or coastal features (such as headlands, islands, sea-mounts, or submarine canyons) and surface forcing (such as atmospheric fronts and convective storms). Regional modelling further allows for the “downscaling” of coarse-resolution global ocean or climate models, permitting the representation of the variation in local conditions that might otherwise be contained within only a few (or even a single!) model grid cells in a global model.

Modular Ocean Model version 6

MOM6 is a widely-used open-source, general circulation ocean–sea ice model, written in Fortran (Adcroft et al., 2019). MOM6 contains several improvements over its predecessor

MOM5 (Griffies, 2014), including the implementation of the Arbitrary-Lagrangian-Eulerian vertical coordinates (Bleck, 2002; Griffies et al., 2020), more efficient tracer advection schemes, and state-of-the-art parameterizations of sub-grid scale physics. Pertinent for our discussion, MOM6 provides support for open boundary conditions and thus is becoming popular for regional ocean modeling studies (see, e.g., Ross et al. (2023), Ross et al. (2024)) in addition to global configurations. However, setting up a regional configuration for MOM6 can be challenging, time consuming, and often involves using several programming languages, a few different tools, and also manually editing/tweaking some input files. The regional-mom6 package overcomes these difficulties, automatically generating a regional MOM6 configuration of the user's choice with relatively simple domain geometry, that is, rectangular domains.




Figure 1: A snapshot of the surface ocean currents from a regional ocean simulation of the Tasman sea using MOM6. The simulation is forced by the GLORYS and ERA5 reanalysis datasets and configured with a horizontal resolution of 1/80th degree and 100 vertical levels (see Barnes (2024) for the source code).

regional-mom6

The regional-mom6 package takes as input various datasets that contain the ocean initial condition, the boundary forcing (ocean and atmosphere) for the regional domain, and the seafloor topography. The input datasets can be on the Arakawa A, B, or C grids (Arakawa & Lamb, 1977); the package performs the appropriate interpolation using xESMF (Zhuang et al., 2023) under the hood, to put the everything on the C grid required by MOM6. This base grid for the regional configuration can be constructed in two ways, either by the user defining a desired resolution and choosing between pre-configured options, or by the user providing pre-existing horizontal and/or vertical MOM6 grids. The user can use MOM6's Arbitrary-Lagrangian-Eulerian vertical coordinates, regardless of the native vertical coordinates of the boundary forcing input. The package automates the re-gridding of all the required forcing input, takes care of all the metadata encoding, generates the regional grid, and ensures that the final input files are in the format expected by MOM6. Additionally, the tricky case of a regional configuration that includes the 'seam' in the longitude of the raw input data (e.g., a 10°-wide regional configuration centred at Fiji (178°E) and forced by input with native longitude coordinate in the range 180°W–180°E) is handled automatically, removing the need for any preprocessing of the input data. This automation allows users to set up a

67 regional MOM6 configuration using only Python and from the convenience of a single Jupyter
68 notebook. Herzfeld et al. (2011) provide rules of thumb to guide the user in setting regional
69 grid parameters such as the resolution.

70 regional-mom6 is installable via conda, it is continuously tested, and comes with extensive
71 documentation including tutorials and examples for setting up regional MOM6 configurations
72 using publicly-available forcing and bathymetry datasets (namely, the GLORYS dataset for ocean
73 boundary forcing (Copernicus Marine Services, 2024), the ERA5 reanalysis for atmospheric
74 forcing (Copernicus Climate Change Service, 2024), and the GEBCO dataset for seafloor
75 topography (GEBCO Bathymetric Compilation Group 2023, 2023)).

76 With the entire process for setting up a regional configuration streamlined to run within a
77 Jupyter notebook, the package dramatically reduces the barrier-to-entry for first-time users, or
78 those without a strong background in Fortran, experience in compiling and running scripts in
79 terminals, and manipulating netCDF files. Besides making regional modelling with MOM6
80 more accessible, our package can automate the generation of multiple experiments (e.g., a
81 series of perturbation experiments), saving time and effort, and improving reproducibility.

82 We designed regional-mom6 with automation of regional configurations in mind. However,
83 the package's code design and modularity make more complex configurations possible since
84 users can use their own custom-made grids with more complex boundaries and construct the
85 boundary forcing terms one by one.

86 Statement of need

87 The learning curve for setting up a regional ocean model can be steep, and it is not obvious
88 for a new user what inputs are required, nor the appropriate format. In the case of MOM6,
89 there are several tools scattered in Github repositories, for example those collected in Earth
90 System Modeling Group grid tools (Simkins et al., 2021). Also, there exist several regional
91 configuration examples but they are hardcoded for particular domains, specific input files, and
92 work only on specific high-performance computing machines (e.g., Ross et al. (2023)).

93 Until now there has been no one-stop-shop for users to learn how to get a regional MOM6
94 configuration up and running. Users are required to use several tools in several programming
95 languages and then modify – sometimes by hand – some of the input metadata to bring
96 everything into the format that MOM6 expects. Many parts of this process are not docu-
97 mented, requiring users to dig into the MOM6 Fortran source code. Recently, the Climate,
98 Ecosystems and Fisheries Initiative gathered these scattered tools into a single repository (Teng
99 et al., 2023) but at the moment they are tailored to a specific high-performance computing
100 machines and not installable as a Python package. Other ocean models have packages to
101 aid in regional configuration setup, for example Pyroms (Hedstrom & contributors, 2023) for
102 the Regional Oceanic Modelling System (ROMS; Shchepetkin & McWilliams (2005)) and
103 MITgcm_python (Naughten & Jones, 2023) for the Massachusetts Institute of Technology
104 General Circulation Model (MITgcm; Marshall et al. (1997)). With MOM6's growing user
105 base for regional applications, there is a need for a platform that walks users through regional
106 domain configuration from start to finish and, ideally, automates the process on the way. Other
107 than reducing the barrier-to-entry, automating the regional configuration process renders the
108 workflow much more reproducible; see discussion by Polton et al. (2023). regional-mom6
109 precisely meets these needs.

110 By having a shared set of tools that the community can work with and contribute to, this
111 package also facilitates collaboration and knowledge-sharing between different research groups.
112 Using a shared framework for setting up regional models, it is easier to compare and contrast
113 examples of different experiments and allows for users to gain intuition for generating their
114 chosen domain.

115 regional-mom6 package can also be used for educational purposes, for example as part of

course curricula. With the technically-challenging aspects of setting up a regional configuration now being automated by the regional-mom6 package, students can set up and run simple MOM6 regional configurations and also change parameters like the model's resolution or the forcing, run again, and see how these parameters affect the ocean flow.

Acknowledgements

We thank the vibrant community of the Consortium for Ocean–Sea Ice Modeling in Australia (cosima.org.au) and also Josué Martínez-Moreno and Callum Shakespeare for useful discussions during the development of this package. We acknowledge support from the Australian Research Council under DECRA Fellowship DE210100749 (N.C.C.) and grant LP200100406 (A.E.K.). We would also like to acknowledge the code and notes by James Simkins, Andrew Ross, and Rob Cermak, which helped us to troubleshoot and improve the algorithms in our package.

References

- Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C. O., Dunne, J. P., Griffies, S. M., Hallberg, R., Harrison, M. J., Held, I. M., Jansen, M. F., John, J. G., Krasting, J. P., Langenhorst, A. R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan, A., ... Zhang, R. (2019). The GFDL global ocean and sea ice model OM4.0: Model description and simulation features. *Journal of Advances in Modeling Earth Systems*, 11(10), 3167–3211. <https://doi.org/10.1029/2019MS001726>
- Arakawa, A., & Lamb, V. R. (1977). Computational design of the basic dynamical processes of the UCLA general circulation model. *Methods in Computational Physics: Advances in Research and Applications*, 17(Supplement C), 173–265. <https://doi.org/10.1016/B978-0-12-460817-7.50009-4>
- Barnes, A. J. (2024). Tasman-tides. In *GitHub repository*. GitHub. <https://github.com/ashjbarnes/tasman-tides>
- Bleck, R. (2002). An oceanic general circulation model framed in hybrid isopycnic-Cartesian coordinates. *Ocean Modelling*, 4(1), 55–88. [https://doi.org/10.1016/S1463-5003\(01\)00012-9](https://doi.org/10.1016/S1463-5003(01)00012-9)
- Copernicus Climate Change Service. (2024). *ECMWF Reanalysis v5*. European Centre for Medium-Range Weather Forecasts. <https://doi.org/10.48670/moi-00021>
- Copernicus Marine Services. (2024). *Global ocean physics reanalysis*. Mercator Ocean International. <https://doi.org/10.48670/moi-00021>
- de Lavergne, C., Groeskamp, S., Zika, J., & Johnson, H. L. (2022). Chapter 3 – the role of mixing in the large-scale ocean circulation. In M. Meredith & A. Naveira Garabato (Eds.), *Ocean mixing* (pp. 35–63). Elsevier. <https://doi.org/10.1016/B978-0-12-821512-8.00010-4>
- GEBCO Bathymetric Compilation Group 2023. (2023). *The GEBCO_2023 Grid - a continuous terrain model of the global oceans and land*. NERC EDS British Oceanographic Data Centre NOC. <https://doi.org/10.5285/f98b053b-0cbc-6c23-e053-6c86abc0af7b>
- Griffies, S. M. (2014). Elements of the modular ocean model (MOM). *GFDL Ocean Group Tech. Rep.*, 7, 47. https://mom-ocean.github.io/assets/pdfs/MOM5_manual.pdf
- Griffies, S. M., Adcroft, A., & Hallberg, R. W. (2020). A primer on the vertical Lagrangian-remap method in ocean models based on finite volume generalized vertical coordinates. *Journal of Advances in Modeling Earth Systems*, 12(10), e2019MS001954. <https://doi.org/10.1029/2019MS001954>

- 160 Gula, J., Taylor, J., Shcherbina, A., & Mahadevan, A. (2022). Chapter 8 – submesoscale
161 processes and mixing. In M. Meredith & A. Naveira Garabato (Eds.), *Ocean mixing* (pp.
162 181–214). Elsevier. <https://doi.org/10.1016/B978-0-12-821512-8.00015-3>
- 163 Hedstrom, K., & contributors. (2023). Pyroms. In *GitHub repository*. GitHub. <https://github.com/ESMG/pyroms>
- 165 Herzfeld, M., Schmidt, M., Griffies, S. M., & Liang, Z. (2011). Realistic test cases for limited
166 area ocean modelling. *Ocean Modelling*, 37(1), 1–34. <https://doi.org/10.1016/j.ocemod.2010.12.008>
- 168 Marshall, J., Adcroft, A., Hill, C., Perelman, L., & Heisey, C. (1997). A finite-volume,
169 incompressible Navier Stokes model for studies of the ocean on parallel computers. *Journal of*
170 *Geophysical Research: Oceans*, 102(C3), 5753–5766. <https://doi.org/10.1029/96JC02775>
- 171 Melet, A. V., Hallberg, R., & Marshall, D. P. (2022). Chapter 2 – the role of ocean mixing
172 in the climate system. In M. Meredith & A. Naveira Garabato (Eds.), *Ocean mixing* (pp.
173 5–34). Elsevier. <https://doi.org/10.1016/B978-0-12-821512-8.00009-8>
- 174 Naughten, K., & Jones, D. (2023). MITgcm_python. In *GitHub repository*. GitHub. https://github.com/knaughten/mitgcm_python
- 176 Orlandi, I. (1976). A simple boundary condition for unbounded hyperbolic flows. *Journal of*
177 *Computational Physics*, 21(3), 251–269. [https://doi.org/10.1016/0021-9991\(76\)90023-1](https://doi.org/10.1016/0021-9991(76)90023-1)
- 178 Polton, J., Harle, J., Holt, J., Katavouta, A., Partridge, D., Jardine, J., Wakelin, S., Rulent, J.,
179 Wise, A., Hutchinson, K., Byrne, D., Bruciaferri, D., O'Dea, E., De Dominicis, M., Mathiot,
180 P., Coward, A., Yool, A., Palmieri, J., Lessin, G., ... Rousset, C. (2023). Reproducible and
181 relocatable regional ocean modelling: Fundamentals and practices. *Geoscientific Model*
182 *Development*, 16(5), 1481–1510. <https://doi.org/10.5194/gmd-16-1481-2023>
- 183 Ross, A. C., Stock, C. A., Adcroft, A., Curchitser, E., Hallberg, R., Harrison, M. J., Hedstrom,
184 K., Zadeh, N., Alexander, M., Chen, W., Drenkard, E. J., Pontavice, H. du, Dussin, R.,
185 Gomez, F., John, J. G., Kang, D., Lavoie, D., Resplandy, L., Roobaert, A., ... Simkins, J.
186 (2023). A high-resolution physical–biogeochemical model for marine resource applications in
187 the northwest Atlantic (MOM6-COBALT-NWA12 v1.0). *Geoscientific Model Development*,
188 16(23), 6943–6985. <https://doi.org/10.5194/gmd-16-6943-2023>
- 189 Ross, A. C., Stock, C. A., Koul, V., Delworth, T. L., Lu, F., Wittenberg, A., & Alexander,
190 M. A. (2024). Dynamically downscaled seasonal ocean forecasts for North American East
191 Coast ecosystems. *EGUsphere*, 2024, 1–40. <https://doi.org/10.5194/egusphere-2024-394>
- 192 Shchepetkin, A. F., & McWilliams, J. C. (2005). The regional oceanic modeling system
193 (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model.
194 *Ocean Modelling*, 9(4), 347–404. <https://doi.org/10.1016/j.ocemod.2004.08.002>
- 195 Silvestri, S., Wagner, G. L., Hill, C., Ardakani, M. R., Blaschke, J., Campin, J.-M., Churavy,
196 V., Constantinou, N. C., Edelman, A., Marshall, J., Ramadhan, A., Souza, A., & Ferrari,
197 R. (2023). *Oceananigans.jl: A model that achieves breakthrough resolution, memory, and*
198 *energy efficiency in global ocean simulations*. <https://arxiv.org/abs/2309.06662>
- 199 Simkins, J., Cermak, R., Hedstrom, K., & Gibson, A. (2021). Earth System Modeling Group
200 (ESMG) gridtools. In *GitHub repository*. GitHub. <https://github.com/ESMG/gridtools>
- 201 Teng, Y.-C., Ross, A., & Morrison, T. (2023). CEFI-regional-MOM6: Essential tools, XML
202 files, and source codes for collaborators of the Climate, Ecosystems, and Fisheries Initiative
203 (CEFI) to conduct simulations. In *GitHub repository*. GitHub. <https://github.com/NOAA-GFDL/CEFI-regional-MOM6>
- 205 Zhuang, J., Dussin, R., Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich,
206 B., Schupfner, M., Filipe, Levang, S., Gauthier, C., Jüling, A., Almansi, M., Scott,
207 R., RondeauG, Rasp, S., Smith, T. J., Stachelek, J., Plough, M., & Li, X. (2023).

208 xESMF: Universal regridding for geospatial data. In *GitHub repository*. Zenodo. [https:](https://github.com/pangeo-data/xESMF)
209 [//github.com/pangeo-data/xESMF](https://github.com/pangeo-data/xESMF)

DRAFT