

# regional\_mom6: Automatic generation of regional configurations for the Modular Ocean Model 6 in Python

Ashley J. Barnes<sup>1,2</sup>, Navid C. Constantinou<sup>1,2</sup>, Angus H. Gibson<sup>1</sup>, Chris Chapman<sup>3</sup>, Dhruv Bhagtani<sup>1,2</sup>, John Reilly<sup>4</sup>, and Andrew E. Kiss<sup>1,2</sup>

<sup>1</sup> Australian National University, Australia <sup>2</sup> ARC Centre of Excellence in Climate Extremes, Australia <sup>3</sup> CSIRO Oceans and Atmosphere, Hobart, Tasmania, Australia <sup>4</sup> University of Tasmania, Australia

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

## Summary

Modular Ocean Model 6 (MOM6) is a widely-used general circulation open-source ocean-sea ice-ice shelf model developed mainly at the Geophysical Fluid Dynamics Laboratory (GFDL) (Adcroft et al., 2019). Among other improvements over its predecessor MOM5 (citation), MOM6 allows open boundary conditions and thus it is becoming popular also for regional ocean modeling studies (see, e.g., Ross et al. (2023), Ross et al. (2024)). However, setting up a regional configuration for MOM6 can be challenging and time consuming and often involves using several programming languages, a few different tools, and also editing/fiddling some input files by hand. The regional\_mom6 python package automates the procedure of generating a regional MOM6 configuration.


The regional\_mom6 package takes as input various datasets that containing the ocean initial condition, the boundary forcing (ocean and atmosphere) for the regional domain, and the bathymetry. The inputs datasets can be on the Arakawa A, B, or C grids (Arakawa & Lamb, 1977); the package performs the appropriate interpolation using xESMF (Zhuang et al., 2023) under the hood, to put the everything on the C grid required by MOM6. Thus, the package automates the re-gridding of all the required forcing input, takes care of all the metadata encoding, generates the regional grid, and deals with few other miscellaneous steps required. This allows users to setup a regional MOM6 configurations using only Python and from a single Jupyter notebook.

regional\_mom6 is continuously tested and comes with an extensive documentation that also includes documented tutorials/examples for setting up regional MOM6 configurations using publicly-available forcing and bathymetry datasets (namely, the GLORYS dataset for ocean boundary forcing (Copernicus Marine Services, 2024), the ERA5 for the atmospheric forcing (Copernicus Climate Change Service, 2024), and the GEBCO dataset for bathymetry (GEBCO Bathymetric Compilation Group 2023, 2023)).

Having the entire process for setting up a regional configuration running in a Jupyter notebook dramatically reduces the barrier of entry for first-time users, or those without a strong background in FORTRAN, experience in compiling and running scripts in terminals, and manipulating netCDF files. Besides making regional modelling with MOM6 more accessible, our package can automate the generation of multiple experiments (e.g., series of perturbation experiments), saving time, effort, and improving reproducibility.

We designed regional\_mom6 with automation of regional configurations in mind. However, the package's code design and modularity allows more complex configurations possible

42 since users can use their own custom-made grids with more complex boundaries and construct  
43 the boundary forcing terms one by one.



**Figure 1:** A snapshot of surface currents from a MOM6 regional simulation of the Tasman sea. The simulation is forced by GLORYS and ERA5 reanalyses datasets. The source code at Barnes (2024).

44 [Figure 1](#) show the surface currents from a region ocean simulation of the Tasman sea configured  
45 with `regional_mom6`. In this case, the forcing datasets are GLORYS and ERA5, with an 1/80th  
46 degree horizontal resolution and 100 vertical levels.

## 47 Statement of need

48 The learning curve for setting up a regional ocean model can be quite steep. In the case of  
49 MOM6, there are several tools scattered in Github repositories, like, for example, those collected  
50 in ESMG's grid tools ([Cermak et al., 2021](#)). Also, there exist several regional configuration  
51 example but that are hardcoded for particular domains, specific input files, and work only on  
52 certain HPC machines.

53 There is no one-stop-shop for users to learn how to get a regional MOM6 configuration up and  
54 running. Users are required to use several tools and in several programming languages and then  
55 modify –sometimes by hand– some of the input metadata to bring everything in the format that  
56 MOM6 expects. Often parts of this process are not documented bringing users down to their  
57 knees since they end up having to figure it out a lot of things by themselves, e.g., by deciphering  
58 FORTRAN code. Other ocean models have packages to aid in regional configuration setup.  
59 Characteristically, Pyroms ([Hedstrom & contributors, 2023](#)) for the Regional Oceanic Modelling  
60 System (ROMS; Shchepetkin & McWilliams (2005)) and MITgcm\_python ([Naughten & Jones, 2023](#))  
61 for the Massachusetts Institute of Technology General Circulation Model (MITgcm;  
62 Marshall et al. (1997)). With MOM6's growing user base for regional applications, there is a  
63 need for a platform that walks users through regional domain configuration from from start to  
64 finish and, ideally, automates the process on the way. `regional_mom6` comes to fill in precisely  
65 this need.

66 By having a shared set of tools that the community can work with and contribute to, this  
67 package also facilitates collaboration and knowledge-sharing between different research groups.

68 Using a shared framework for setting up regional models, it is easier to compare and contrast  
69 examples of different experiments and allows for user to gain intuition for generating their  
70 chosen domain.

71 Another potential advantage of a package that allows users to automatically obtain regional  
72 configurations of MOM6 is in education. With the technically-challenging aspects of setting  
73 up a regional configuration now being automated by `regional_mom6` package, students can  
74 setup and run simple MOM6 regional configurations and also change parameters like resolution  
75 or forcing, run again, and see how these parameters affect the ocean flow.

## 76 Acknowledgements

77 We thank the Consortium for Ocean–Sea Ice Modeling in Australia ([cosima.org.au](https://cosima.org.au)) for useful  
78 discussions during the development of this package. N.C.C. acknowledges funding from the  
79 Australian Research Council under DECRA Fellowship DE210100749. We would also like to  
80 acknowledge the code and notes by James Simkins, Andrew Ross, and Rob Cermak, which  
81 helped us to troubleshoot and improve the algorithms in our package.

## 82 References

- 83 Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C. O., Dunne, J.  
84 P., Griffies, S. M., Hallberg, R., Harrison, M. J., Held, I. M., Jansen, M. F., John, J. G.,  
85 Krasting, J. P., Langenhorst, A. R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan,  
86 A., ... Zhang, R. (2019). The GFDL global ocean and sea ice model OM4.0: Model  
87 description and simulation features. *Journal of Advances in Modeling Earth Systems*,  
88 11(10), 3167–3211. <https://doi.org/10.1029/2019MS001726>
- 89 Arakawa, A., & Lamb, V. R. (1977). Computational design of the basic dynamical processes  
90 of the UCLA general circulation model. *Methods in Computational Physics: Advances*  
91 *in Research and Applications*, 17(Supplement C), 173–265. [https://doi.org/10.1016/](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)  
92 [B978-0-12-460817-7.50009-4](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)
- 93 Barnes, A. J. (2024). Tasman-tides. In *GitHub repository*. [https://github.com/ashjbarnes/](https://github.com/ashjbarnes/tasman-tides)  
94 [tasman-tides](https://github.com/ashjbarnes/tasman-tides)
- 95 Cermak, R., Simkins, J., Hedstrom, K., & Gibson, A. (2021). ESMG gridtools. In *GitHub*  
96 *repository*. Github. <https://github.com/ESMG/gridtools>
- 97 Copernicus Climate Change Service. (2024). *ECMWF Reanalysis v5*. European Centre for  
98 Medium-Range Weather Forecasts. <https://doi.org/10.48670/moi-00021>
- 99 Copernicus Marine Services. (2024). *Global ocean physics reanalysis*. Mercator Ocean  
100 International. <https://doi.org/10.48670/moi-00021>
- 101 GEBCO Bathymetric Compilation Group 2023. (2023). *The GEBCO\_2023 Grid - a continuous*  
102 *terrain model of the global oceans and land*. NERC EDS British Oceanographic Data  
103 Centre NOC. <https://doi.org/10.5285/f98b053b-0cbc-6c23-e053-6c86abc0af7b>
- 104 Hedstrom, K., & contributors. (2023). Pyroms. In *GitHub repository*. Github. [https://](https://github.com/ESMG/pyroms)  
105 [github.com/ESMG/pyroms](https://github.com/ESMG/pyroms)
- 106 Marshall, J., Adcroft, A., Hill, C., Perelman, L., & Heisey, C. (1997). A finite-volume,  
107 incompressible Navier Stokes model for studies of the ocean on parallel computers. *Journal of*  
108 *Geophysical Research: Oceans*, 102(C3), 5753–5766. <https://doi.org/10.1029/96JC02775>
- 109 Naughten, K., & Jones, D. (2023). MITgcm\_python. In *GitHub repository*. Github. [https://](https://github.com/knaughten/mitgcm_python)  
110 [github.com/knaughten/mitgcm\\_python](https://github.com/knaughten/mitgcm_python)

- 111 Ross, A. C., Stock, C. A., Adcroft, A., Curchitser, E., Hallberg, R., Harrison, M. J., Hedstrom,  
112 K., Zadeh, N., Alexander, M., Chen, W., Drenkard, E. J., Pontavice, H. du, Dussin, R.,  
113 Gomez, F., John, J. G., Kang, D., Lavoie, D., Resplandy, L., Roobaert, A., ... Simkins, J.  
114 (2023). A high-resolution physical–biogeochemical model for marine resource applications in  
115 the northwest Atlantic (MOM6-COBALT-NWA12 v1.0). *Geoscientific Model Development*,  
116 *16*(23), 6943–6985. <https://doi.org/10.5194/gmd-16-6943-2023>
- 117 Ross, A. C., Stock, C. A., Koul, V., Delworth, T. L., Lu, F., Wittenberg, A., & Alexander,  
118 M. A. (2024). Dynamically downscaled seasonal ocean forecasts for North American East  
119 Coast ecosystems. *EGUsphere*, 2024, 1–40. <https://doi.org/10.5194/egusphere-2024-394>
- 120 Shchepetkin, A. F., & McWilliams, J. C. (2005). The regional oceanic modeling system  
121 (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model.  
122 *Ocean Modelling*, *9*(4), 347–404. <https://doi.org/10.1016/j.ocemod.2004.08.002>
- 123 Zhuang, J., Dussin, R., Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich,  
124 B., Schupfner, M., Filipe, Levang, S., Gauthier, C., Jüling, A., Almansi, M., Scott,  
125 R., RondeauG, Rasp, S., Smith, T. J., Stachelek, J., Plough, M., & Li, X. (2023).  
126 xESMF: Universal regridding for geospatial data. In *GitHub repository*. Zenodo. <https://github.com/pangeo-data/xESMF>

DRAFT