

regional_mom6: Automatic generation of regional configurations for the Modular Ocean Model 6 in Python

Ashley J. Barnes^{1,2}, Navid C. Constantinou^{1,2}, Angus H. Gibson¹, Chris Chapman³, Dhruv Bhagtani^{1,2}, John Reilly⁴, and Andrew E. Kiss^{1,2}

¹ Australian National University, Australia ² ARC Centre of Excellence in Climate Extremes, Australia ³ CSIRO Oceans and Atmosphere, Hobart, Tasmania, Australia ⁴ University of Tasmania, Australia

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

Summary

Modular Ocean Model 6 (MOM6) is a widely-used general circulation open-source ocean-sea ice-ice shelf model developed mainly at the Geophysical Fluid Dynamics Laboratory (GFDL) (Adcroft et al., 2019). Among other improvements over its predecessor MOM5 (citation), MOM6 allows open boundary conditions and thus it is becoming popular also for regional ocean modeling studies (see, e.g., Ross et al. (2023), Ross et al. (2024)). However, setting up a regional configuration for MOM6 can be challenging and time consuming and often involves using several programming languages, a few different tools, and also editing/fiddling some input files by hand. The regional_mom6 python package automates the procedure of generating a regional MOM6 configuration.


The regional_mom6 package takes as input various datasets that containing the ocean initial condition, the boundary forcing (ocean and atmosphere) for the regional domain, and the bathymetry. The inputs datasets can be on the Arakawa A, B, or C grids (Arakawa & Lamb, 1977); the package performs the appropriate interpolation using xESMF (Zhuang et al., 2023) under the hood, to put the everything on the C grid required by MOM6. Thus, the package automates the re-gridding of all the required forcing input, takes care of all the metadata encoding, generates the regional grid, and deals with few other miscellaneous steps required. This allows users to setup a regional MOM6 configurations using only Python and from a single Jupyter notebook.

regional_mom6 is continuously tested and comes with an extensive documentation that also includes documented tutorials/examples for setting up regional MOM6 configurations using publicly-available forcing and bathymetry datasets (namely, the GLORYS dataset for ocean boundary forcing (Copernicus Marine Services, 2024), the ERA5 for the atmospheric forcing (Copernicus Climate Change Service, 2024), and the GEBCO dataset for bathymetry (GEBCO Bathymetric Compilation Group 2023, 2023)).

Having the entire process for setting up a regional configuration running in a Jupyter notebook dramatically reduces the barrier of entry for first-time users, or those without a strong background in FORTRAN, experience in compiling and running scripts in terminals, and manipulating netCDF files. Besides making regional modelling with MOM6 more accessible, our package can automate the generation of multiple experiments (e.g., series of perturbation experiments), saving time, effort, and improving reproducibility.

We designed regional_mom6 with automation of regional configurations in mind. However, the package's code design and modularity allows more complex configurations possible

42 since users can use their own custom-made grids with more complex boundaries and construct
43 the boundary forcing terms one by one.



45 Statement of need

46 The learning curve for setting up a regional ocean model can be quite steep. In the case of
47 MOM6, there are several tools scattered in Github repositories, like, for example, those collected
48 in ESMG's grid tools (Cermak et al., 2021). Also, there exist several regional configuration
49 example but that are hardcoded for particular domains, specific input files, and work only on
50 certain HPC machines.

51 There is no one-stop-shop for users to learn how to get a regional MOM6 configuration up and
52 running. Users are required to use several tools and in several programming languages and then
53 modify—sometimes by hand—some of the input metadata to bring everything in the format that
54 MOM6 expects. Often parts of this process are not documented bringing users down to their
55 knees since they end up having to figure it out a lot of things by themselves, e.g., by deciphering
56 FORTRAN code. Other ocean models have packages to aid in regional configuration setup.
57 Characteristically, Pyroms (Hedstrom & contributors, 2023) for the Regional Oceanic Modelling
58 System (ROMS; Shchepetkin & McWilliams (2005)) and MITgcm_python (Naughten & Jones,
59 2023) for the Massachusetts Institute of Technology General Circulation Model (MITgcm;
60 Marshall et al. (1997)). With MOM6's growing user base for regional applications, there is a
61 need for a platform that walks users through regional domain configuration from from start to
62 finish and, ideally, automates the process on the way. regional_mom6 comes to fill in precisely
63 this need.

64 By having a shared set of tools that the community can work with and contribute to, this
65 package also facilitates collaboration and knowledge-sharing between different research groups.
66 Using a shared framework for setting up regional models, it is easier to compare and contrast
67 examples of different experiments and allows for user to gain intuition for generating their

68 chosen domain.

69 Another potential advantage of a package that allows users to automatically obtain regional
70 configurations of MOM6 is in education. With the technically-challenging aspects of setting
71 up a regional configuration now being automated by `regional_mom6` package, students can
72 setup and run simple MOM6 regional configurations and also change parameters like resolution
73 or forcing, run again, and see how these parameters affect the ocean flow.

74 Acknowledgements

75 We thank the Consortium for Ocean–Sea Ice Modeling in Australia (cosima.org.au) for useful
76 discussions during the development of this package. N.C.C. acknowledges funding from the
77 Australian Research Council under DECRA Fellowship DE210100749. We would also like to
78 acknowledge the code and notes by James Simkins, Andrew Ross, and Rob Cermak, which
79 helped us to troubleshoot and improve the algorithms in our package.

80 References

- 81 Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C. O., Dunne, J.
82 P., Griffies, S. M., Hallberg, R., Harrison, M. J., Held, I. M., Jansen, M. F., John, J. G.,
83 Krasting, J. P., Langenhorst, A. R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan,
84 A., ... Zhang, R. (2019). The GFDL global ocean and sea ice model OM4.0: Model
85 description and simulation features. *Journal of Advances in Modeling Earth Systems*,
86 11(10), 3167–3211. <https://doi.org/10.1029/2019MS001726>
- 87 Arakawa, A., & Lamb, V. R. (1977). Computational design of the basic dynamical processes
88 of the UCLA general circulation model. *Methods in Computational Physics: Advances*
89 *in Research and Applications*, 17(Supplement C), 173–265. [https://doi.org/10.1016/](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)
90 [B978-0-12-460817-7.50009-4](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)
- 91 Cermak, R., Simkins, J., Hedstrom, K., & Gibson, A. (2021). ESMG gridtools. In *GitHub*
92 *repository*. Github. <https://github.com/ESMG/gridtools>
- 93 Copernicus Climate Change Service. (2024). *ECMWF Reanalysis v5*. European Centre for
94 Medium-Range Weather Forecasts. <https://doi.org/10.48670/moi-00021>
- 95 Copernicus Marine Services. (2024). *Global ocean physics reanalysis*. Mercator Ocean
96 International. <https://doi.org/10.48670/moi-00021>
- 97 GEBCO Bathymetric Compilation Group 2023. (2023). *The GEBCO_2023 Grid - a continuous*
98 *terrain model of the global oceans and land*. NERC EDS British Oceanographic Data
99 Centre NOC. <https://doi.org/10.5285/f98b053b-0cbc-6c23-e053-6c86abc0af7b>
- 100 Hedstrom, K., & contributors. (2023). Pyroms. In *GitHub repository*. Github. [https://](https://github.com/ESMG/pyroms)
101 github.com/ESMG/pyroms
- 102 Marshall, J., Adcroft, A., Hill, C., Perelman, L., & Heisey, C. (1997). A finite-volume,
103 incompressible Navier Stokes model for studies of the ocean on parallel computers. *Journal of*
104 *Geophysical Research: Oceans*, 102(C3), 5753–5766. <https://doi.org/10.1029/96JC02775>
- 105 Naughten, K., & Jones, D. (2023). MITgcm_python. In *GitHub repository*. Github. [https://](https://github.com/knaughten/mitgcm_python)
106 github.com/knaughten/mitgcm_python
- 107 Ross, A. C., Stock, C. A., Adcroft, A., Curchitser, E., Hallberg, R., Harrison, M. J., Hedstrom,
108 K., Zadeh, N., Alexander, M., Chen, W., Drenkard, E. J., Pontavice, H. du, Dussin, R.,
109 Gomez, F., John, J. G., Kang, D., Lavoie, D., Resplandy, L., Roobaert, A., ... Simkins, J.
110 (2023). A high-resolution physical–biogeochemical model for marine resource applications in

- 111 the northwest Atlantic (MOM6-COBALT-NWA12 v1.0). *Geoscientific Model Development*,
112 16(23), 6943–6985. <https://doi.org/10.5194/gmd-16-6943-2023>
- 113 Ross, A. C., Stock, C. A., Koul, V., Delworth, T. L., Lu, F., Wittenberg, A., & Alexander,
114 M. A. (2024). Dynamically downscaled seasonal ocean forecasts for North American East
115 Coast ecosystems. *EGUsphere*, 2024, 1–40. <https://doi.org/10.5194/egusphere-2024-394>
- 116 Shchepetkin, A. F., & McWilliams, J. C. (2005). The regional oceanic modeling system
117 (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model.
118 *Ocean Modelling*, 9(4), 347–404. <https://doi.org/10.1016/j.ocemod.2004.08.002>
- 119 Zhuang, J., Dussin, R., Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich,
120 B., Schupfner, M., Filipe, Levang, S., Gauthier, C., Jüling, A., Almansi, M., Scott,
121 R., RondeauG, Rasp, S., Smith, T. J., Stachelek, J., Plough, M., & Li, X. (2023).
122 xESMF: Universal regridding for geospatial data. In *GitHub repository*. Zenodo. <https://github.com/pangeo-data/xESMF>
123

DRAFT