

# regional-mom6: Automatic generation of regional configurations for the Modular Ocean Model v6 in Python

Ashley J. Barnes<sup>1,2</sup>, Navid C. Constantinou<sup>1,2,3,4</sup>, Angus H. Gibson<sup>1</sup>, Andrew E. Kiss<sup>1,2</sup>, Chris Chapman<sup>5</sup>, John Reilly<sup>6</sup>, and Dhruv Bhagtani<sup>1,2</sup>

<sup>1</sup> Australian National University, Australia <sup>2</sup> ARC Centre of Excellence for Climate Extremes, Australia <sup>3</sup> University of Melbourne, Australia <sup>4</sup> ARC Centre of Excellence for the Weather of the 21st Century, Australia <sup>5</sup> CSIRO Environment, Hobart, Tasmania, Australia <sup>6</sup> University of Tasmania, Australia

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary


regional-mom6 is a Python package that provides an easy and versatile way to set up regional configurations of the Modular Ocean Model version 6 (MOM6).

In the ocean, fast and small-scale motions (from ~100m to ~100km varying at time scales of hours to days) play an important role in shaping the large-scale ocean circulation and climate (length scales ~10,000km varying at decadal time scales) (de Lavergne et al., 2022; Gula et al., 2022; Melet et al., 2022). Despite the increase in computational power and the use of graphical processing units that can bring breakthrough performance and speedup (Silvestri et al., 2023), there are always processes, boundary, or forcing features that are smaller than the model's grid spacing and, thus, remain unresolved in operational ocean models. Regional ocean models can be run at higher resolutions while limiting the required computational resources.

A regional ocean model simulates the ocean only in a prescribed region, which is a subset of the global ocean. To do that, we need to apply open boundary conditions at the region's boundaries, that is, we need to impose conditions that mimic the oceanic flow that we are not simulating (Orlanski, 1976). For example, Figure 1 shows the surface currents from a regional ocean simulation of the Tasman sea that was configured using the regional-mom6 package. The boundaries of the domain depicted in Figure 1 are forced with the ocean flow from a global ocean reanalysis product. Higher-resolution regional ocean models improve the representation of smaller-scale motions, such as tidal beams, mixing, mesoscale and sub-mesoscale circulation, as well as the oceanic response to smaller-scale bathymetric or coastal features (such as headlands, islands, sea-mounts, or submarine canyons) and surface forcing (such as atmospheric fronts and convective storms). Regional modelling further allows for the “downscaling” of coarse-resolution global ocean or climate models, permitting the representation of the variation in local conditions that might otherwise be contained within only a few (or even a single!) model grid cells in a global model.

MOM6 is a widely-used open-source, general circulation ocean–sea ice model, written in Fortran (Adcroft et al., 2019). MOM6 contains several improvements over its predecessor MOM5 (Griffies, 2014), including the implementation of the Arbitrary-Lagrangian-Eulerian vertical coordinates (Bleck, 2002; Griffies et al., 2020), more efficient tracer advection schemes, and state-of-the art parameterizations of sub-grid scale physics. Pertinent for our discussion, MOM6 provides support for open boundary conditions and thus is becoming popular for regional ocean modeling studies (see, e.g., Ross et al. (2023), Ross et al. (2024)) in addition to global configurations. However, setting up a regional configuration for MOM6 can be challenging,

time consuming, and often involves using several programming languages, a few different tools, and also manually editing/tweaking some input files. The regional-mom6 package overcomes these difficulties, automatically generating a regional MOM6 configuration of the user's choice with relatively simple domain geometry, that is, rectangular domains.



**Figure 1:** A snapshot of the surface ocean currents from a regional ocean simulation of the Tasman sea using MOM6. The simulation is forced by the GLORYS and ERA5 reanalysis datasets and configured with a horizontal resolution of 1/80th degree and 100 vertical levels (see Barnes (2024) for the source code).

The regional-mom6 package takes as input various datasets that contain the ocean initial condition, the boundary forcing (ocean and atmosphere) for the regional domain, and the seafloor topography. The input datasets can be on the Arakawa A, B, or C grids (Arakawa & Lamb, 1977); the package performs the appropriate interpolation using xESMF (Zhuang et al., 2023) under the hood, to put the everything on the C grid required by MOM6. This base grid for the regional configuration can be constructed in two ways, either by the user defining a desired resolution and choosing between pre-configured options, or by the user providing pre-existing horizontal and/or vertical MOM6 grids. The user can use MOM6's Arbitrary-Lagrangian-Eulerian vertical coordinates (Griffies et al., 2020), regardless of the native vertical coordinates of the boundary forcing input. The package automates the re-gridding of all the required forcing input, takes care of all the metadata encoding, generates the regional grid, and ensures that the final input files are in the format expected by MOM6. Additionally, the tricky case of a regional configuration that includes the 'seam' in the longitude of the raw input data (e.g. a 10°-wide regional configuration centred at Fiji (178°E) and forced by input with native longitude coordinate in the range 180°W–180°E) is handled automatically, removing the need for any preprocessing of the input data. This automation allows users to set up a regional MOM6 configuration using only Python and from the convenience of a single Jupyter notebook. Herzfeld et al. (2011) provide rules of thumb to guide the user in setting regional grid parameters such as the resolution.

regional-mom6 is installable via conda, it is continuously tested, and comes with extensive documentation including tutorials and examples for setting up regional MOM6 configurations using publicly-available forcing and bathymetry datasets (namely, the GLORYS dataset for ocean boundary forcing (Copernicus Marine Services, 2024), the ERA5 reanalysis for atmospheric

70 forcing ([Copernicus Climate Change Service, 2024](#)), and the GEBCO dataset for seafloor  
71 topography ([GEBCO Bathymetric Compilation Group 2023, 2023](#))).

72 With the entire process for setting up a regional configuration streamlined to run within a  
73 Jupyter notebook, the package dramatically reduces the barrier-to-entry for first-time users, or  
74 those without a strong background in Fortran, experience in compiling and running scripts in  
75 terminals, and manipulating netCDF files. Besides making regional modelling with MOM6  
76 more accessible, our package can automate the generation of multiple experiments (e.g., a  
77 series of perturbation experiments), saving time and effort, and improving reproducibility.

78 We designed regional-mom6 with automation of regional configurations in mind. However,  
79 the package's code design and modularity make more complex configurations possible since  
80 users can use their own custom-made grids with more complex boundaries and construct the  
81 boundary forcing terms one by one.

## 82 Statement of need

83 The learning curve for setting up a regional ocean model can be steep, and it is not obvious for  
84 a new user what inputs are required, nor the appropriate format. In the case of MOM6, there  
85 are several tools scattered in Github repositories, for example those collected in Earth System  
86 Modeling Group grid tools ([Simkins et al., 2021](#)). Also, there exist several regional configuration  
87 examples (e.g., **[ADD 1-2 CITATIONS OF REPOS HERE?]**) but they are hardcoded for  
88 particular domains, specific input files, and work only on specific high-performance computing  
89 machines.

90 Until now there has been no one-stop-shop for users to learn how to get a regional MOM6  
91 configuration up and running. Users are required to use several tools in several programming  
92 languages and then modify – sometimes by hand – some of the input metadata to bring  
93 everything into the format that MOM6 expects. Many parts of this process are not documented,  
94 requiring users to dig into the MOM6 Fortran source code. Other ocean models have packages  
95 to aid in regional configuration setup, for example Pyroms ([Hedstrom & contributors, 2023](#))  
96 for the Regional Oceanic Modelling System (ROMS; Shchepetkin & McWilliams (2005)) and  
97 MITgcm\_python ([Naughten & Jones, 2023](#)) for the Massachusetts Institute of Technology  
98 General Circulation Model (MITgcm; Marshall et al. (1997)). With MOM6's growing user  
99 base for regional applications, there is a need for a platform that walks users through regional  
100 domain configuration from start to finish and, ideally, automates the process on the way. Other  
101 than reducing the barrier-to-entry, automating the regional configuration renders the workflow  
102 much more reproducible; see discussion by Polton et al. (2023). regional-mom6 precisely  
103 meets these needs.

104 By having a shared set of tools that the community can work with and contribute to, this  
105 package also facilitates collaboration and knowledge-sharing between different research groups.  
106 Using a shared framework for setting up regional models, it is easier to compare and contrast  
107 examples of different experiments and allows for users to gain intuition for generating their  
108 chosen domain.

109 regional-mom6 package can also be used for educational purposes, for example as part of  
110 course curricula. With the technically-challenging aspects of setting up a regional configuration  
111 now being automated by the regional-mom6 package, students can set up and run simple  
112 MOM6 regional configurations and also change parameters like the model's resolution or the  
113 forcing, run again, and see how these parameters affect the ocean flow.

## 114 Acknowledgements

115 We thank the Consortium for Ocean–Sea Ice Modeling in Australia ([cosima.org.au](#)), Josué  
116 Martínez-Moreno, and Callum Shakespeare for useful discussions during the development of

117 this package. We acknowledge support from the Australian Research Council under DECRA  
 118 Fellowship DE210100749 (N.C.C.) and grant LP200100406 (A.E.K.). We would also like to  
 119 acknowledge the code and notes by James Simkins, Andrew Ross, and Rob Cermak, which  
 120 helped us to troubleshoot and improve the algorithms in our package.

## 121 References

- 122 Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C. O., Dunne, J.  
 123 P., Griffies, S. M., Hallberg, R., Harrison, M. J., Held, I. M., Jansen, M. F., John, J. G.,  
 124 Krasting, J. P., Langenhorst, A. R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan,  
 125 A., ... Zhang, R. (2019). The GFDL global ocean and sea ice model OM4.0: Model  
 126 description and simulation features. *Journal of Advances in Modeling Earth Systems*,  
 127 11(10), 3167–3211. <https://doi.org/10.1029/2019MS001726>
- 128 Arakawa, A., & Lamb, V. R. (1977). Computational design of the basic dynamical processes  
 129 of the UCLA general circulation model. *Methods in Computational Physics: Advances*  
 130 *in Research and Applications*, 17(Supplement C), 173–265. [https://doi.org/10.1016/](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)  
 131 [B978-0-12-460817-7.50009-4](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)
- 132 Barnes, A. J. (2024). Tasman-tides. In *GitHub repository*. GitHub. [https://github.com/](https://github.com/ashjbarnes/tasman-tides)  
 133 [ashjbarnes/tasman-tides](https://github.com/ashjbarnes/tasman-tides)
- 134 Bleck, R. (2002). An oceanic general circulation model framed in hybrid isopycnic-Cartesian  
 135 coordinates. *Ocean Modelling*, 4(1), 55–88. [https://doi.org/10.1016/S1463-5003\(01\)](https://doi.org/10.1016/S1463-5003(01)00012-9)  
 136 [00012-9](https://doi.org/10.1016/S1463-5003(01)00012-9)
- 137 Copernicus Climate Change Service. (2024). *ECMWF Reanalysis v5*. European Centre for  
 138 Medium-Range Weather Forecasts. <https://doi.org/10.48670/moi-00021>
- 139 Copernicus Marine Services. (2024). *Global ocean physics reanalysis*. Mercator Ocean  
 140 International. <https://doi.org/10.48670/moi-00021>
- 141 de Lavergne, C., Groeskamp, S., Zika, J., & Johnson, H. L. (2022). Chapter 3 – the role  
 142 of mixing in the large-scale ocean circulation. In M. Meredith & A. Naveira Garabato  
 143 (Eds.), *Ocean mixing* (pp. 35–63). Elsevier. [https://doi.org/10.1016/B978-0-12-821512-8.](https://doi.org/10.1016/B978-0-12-821512-8.00010-4)  
 144 [00010-4](https://doi.org/10.1016/B978-0-12-821512-8.00010-4)
- 145 GEBCO Bathymetric Compilation Group 2023. (2023). *The GEBCO\_2023 Grid - a continuous*  
 146 *terrain model of the global oceans and land*. NERC EDS British Oceanographic Data  
 147 Centre NOC. <https://doi.org/10.5285/f98b053b-0cbc-6c23-e053-6c86abc0af7b>
- 148 Griffies, S. M. (2014). Elements of the modular ocean model (MOM). *GFDL Ocean Group*  
 149 *Tech. Rep.*, 7, 47. [https://mom-ocean.github.io/assets/pdfs/MOM5\\_manual.pdf](https://mom-ocean.github.io/assets/pdfs/MOM5_manual.pdf)
- 150 Griffies, S. M., Adcroft, A., & Hallberg, R. W. (2020). A primer on the vertical Lagrangian-  
 151 remap method in ocean models based on finite volume generalized vertical coordinates.  
 152 *Journal of Advances in Modeling Earth Systems*, 12(10), e2019MS001954. [https://doi.](https://doi.org/10.1029/2019MS001954)  
 153 [org/10.1029/2019MS001954](https://doi.org/10.1029/2019MS001954)
- 154 Gula, J., Taylor, J., Shcherbina, A., & Mahadevan, A. (2022). Chapter 8 – submesoscale  
 155 processes and mixing. In M. Meredith & A. Naveira Garabato (Eds.), *Ocean mixing* (pp.  
 156 181–214). Elsevier. <https://doi.org/10.1016/B978-0-12-821512-8.00015-3>
- 157 Hedstrom, K., & contributors. (2023). Pyroms. In *GitHub repository*. GitHub. [https://](https://github.com/ESMG/pyroms)  
 158 [github.com/ESMG/pyroms](https://github.com/ESMG/pyroms)
- 159 Herzfeld, M., Schmidt, M., Griffies, S. M., & Liang, Z. (2011). Realistic test cases for limited  
 160 area ocean modelling. *Ocean Modelling*, 37(1), 1–34. [https://doi.org/10.1016/j.ocemod.](https://doi.org/10.1016/j.ocemod.2010.12.008)  
 161 [2010.12.008](https://doi.org/10.1016/j.ocemod.2010.12.008)



- Marshall, J., Adcroft, A., Hill, C., Perelman, L., & Heisey, C. (1997). A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers. *Journal of Geophysical Research: Oceans*, 102(C3), 5753–5766. <https://doi.org/10.1029/96JC02775>
- Melet, A. V., Hallberg, R., & Marshall, D. P. (2022). Chapter 2 – the role of ocean mixing in the climate system. In M. Meredith & A. Naveira Garabato (Eds.), *Ocean mixing* (pp. 5–34). Elsevier. <https://doi.org/10.1016/B978-0-12-821512-8.00009-8>
- Naughten, K., & Jones, D. (2023). MITgcm\_python. In *GitHub repository*. GitHub. [https://github.com/knaughten/mitgcm\\_python](https://github.com/knaughten/mitgcm_python)
- Orlanski, I. (1976). A simple boundary condition for unbounded hyperbolic flows. *Journal of Computational Physics*, 21(3), 251–269. [https://doi.org/10.1016/0021-9991\(76\)90023-1](https://doi.org/10.1016/0021-9991(76)90023-1)
- Polton, J., Harle, J., Holt, J., Katavouta, A., Partridge, D., Jardine, J., Wakelin, S., Rulent, J., Wise, A., Hutchinson, K., Byrne, D., Bruciaferri, D., O'Dea, E., De Dominicis, M., Mathiot, P., Coward, A., Yool, A., Palmieri, J., Lessin, G., ... Rousset, C. (2023). Reproducible and relocatable regional ocean modelling: Fundamentals and practices. *Geoscientific Model Development*, 16(5), 1481–1510. <https://doi.org/10.5194/gmd-16-1481-2023>
- Ross, A. C., Stock, C. A., Adcroft, A., Curchitser, E., Hallberg, R., Harrison, M. J., Hedstrom, K., Zadeh, N., Alexander, M., Chen, W., Drenkard, E. J., Pontavice, H. du, Dussin, R., Gomez, F., John, J. G., Kang, D., Lavoie, D., Resplandy, L., Roobaert, A., ... Simkins, J. (2023). A high-resolution physical–biogeochemical model for marine resource applications in the northwest Atlantic (MOM6-COBALT-NWA12 v1.0). *Geoscientific Model Development*, 16(23), 6943–6985. <https://doi.org/10.5194/gmd-16-6943-2023>
- Ross, A. C., Stock, C. A., Koul, V., Delworth, T. L., Lu, F., Wittenberg, A., & Alexander, M. A. (2024). Dynamically downscaled seasonal ocean forecasts for North American East Coast ecosystems. *EGUsphere*, 2024, 1–40. <https://doi.org/10.5194/egusphere-2024-394>
- Shchepetkin, A. F., & McWilliams, J. C. (2005). The regional oceanic modeling system (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model. *Ocean Modelling*, 9(4), 347–404. <https://doi.org/10.1016/j.ocemod.2004.08.002>
- Silvestri, S., Wagner, G. L., Hill, C., Ardakani, M. R., Blaschke, J., Campin, J.-M., Churavy, V., Constantinou, N. C., Edelman, A., Marshall, J., Ramadhan, A., Souza, A., & Ferrari, R. (2023). *Oceananigans.jl: A model that achieves breakthrough resolution, memory, and energy efficiency in global ocean simulations*. <https://arxiv.org/abs/2309.06662>
- Simkins, J., Cermak, R., Hedstrom, K., & Gibson, A. (2021). Earth System Modeling Group (ESMG) gridtools. In *GitHub repository*. GitHub. <https://github.com/ESMG/gridtools>
- Zhuang, J., Dussin, R., Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich, B., Schupfner, M., Filipe, Levang, S., Gauthier, C., Jüling, A., Almansi, M., Scott, R., RondeauG, Rasp, S., Smith, T. J., Stachelek, J., Plough, M., & Li, X. (2023). xESMF: Universal regridded for geospatial data. In *GitHub repository*. Zenodo. <https://github.com/pangeo-data/xESMF>