

# regional\_mom6: Automatic generation of regional configurations for the Modular Ocean Model 6 in Python

Ashley J. Barnes<sup>1,2</sup>, Navid C. Constantinou<sup>1,2</sup>, Angus H. Gibson<sup>1</sup>, Chris Chapman<sup>3</sup>, Dhruv Bhagtani<sup>1,2</sup>, John Reilly<sup>4</sup>, and Andrew E. Kiss<sup>1,2</sup>

<sup>1</sup> Australian National University, Australia <sup>2</sup> ARC Centre of Excellence in Climate Extremes, Australia <sup>3</sup> CSIRO Environment, Hobart, Tasmania, Australia <sup>4</sup> University of Tasmania, Australia

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

## Summary

The Modular Ocean Model version 6 (MOM6) is a widely-used open-source general circulation ocean-sea ice model developed mainly at the NOAA Geophysical Fluid Dynamics Laboratory (GFDL) (Adcroft et al., 2019). MOM6 contains several improvements over its predecessor MOM5 (Stephen M. Griffies, 2014), including the implementation of the Arbitrary-Lagrangian-Eulerian (ALE) vertical coordinates (Stephen M. Griffies et al., 2020), more efficient tracer advection schemes, and state-of-the art parameterisations of sub-grid scale physics. MOM6 provides support for open boundary conditions and thus is becoming popular for regional ocean modeling studies (see, e.g., Ross et al. (2023), Ross et al. (2024)) in addition to global configurations. However, setting up a regional configuration for MOM6 can be challenging, time consuming, and often involves using several programming languages, a few different tools, and also manually editing/tweaking some input files. The regional\_mom6 python package overcomes these difficulties, automatically generating a regional MOM6 configuration with relatively simple domain geometry.


The regional\_mom6 package takes as input various datasets that containing the ocean initial condition, the boundary forcing (ocean and atmosphere) for the regional domain, and the bathymetry. The input datasets can be on the Arakawa A, B, or C grids (Arakawa & Lamb, 1977); the package performs the appropriate interpolation using xESMF (Zhuang et al., 2023) under the hood, to put the everything on the C grid required by MOM6. Thus, the package automates the re-gridding of all the required forcing input, takes care of all the metadata encoding, generates the regional grid, and deals with a few other necessary steps. This allows users to setup a regional MOM6 configuration using only Python and from a single Jupyter notebook. The package allows the user to use MOM6's Arbitrary-Lagrangian-Eulerian vertical coordinates, regardless of the vertical coordinates of boundary forcing input. Rules-of-thumb to guide the user in setting grid parameters such as the regional domains resolution, can be found in the paper by Herzfeld et al. (2011).

regional\_mom6 is continuously tested and comes with an extensive documentation that also includes documented tutorials/examples for setting up regional MOM6 configurations using publicly-available forcing and bathymetry datasets (namely, the GLORYS dataset for ocean boundary forcing (Copernicus Marine Services, 2024), the ERA5 reanalysis for atmospheric forcing (Copernicus Climate Change Service, 2024), and the GEBCO dataset for bathymetry (GEBCO Bathymetric Compilation Group 2023, 2023)).

Having the entire process for setting up a regional configuration running in a Jupyter notebook dramatically reduces the barrier to entry for first-time users, or those without a strong back-

ground in Fortran, experience in compiling and running scripts in terminals, and manipulating netCDF files. Besides making regional modelling with MOM6 more accessible, our package can automate the generation of multiple experiments (e.g., a series of perturbation experiments), saving time and effort, and improving reproducibility.

We designed `regional_mom6` with automation of regional configurations in mind. However, the package's code design and modularity makes more complex configurations possible since users can use their own custom-made grids with more complex boundaries and construct the boundary forcing terms one by one.



**Figure 1:** A snapshot of the ocean surface currents from a MOM6 regional simulation of the Tasman sea. The simulation is forced by GLORYS and ERA5 reanalysis datasets and configured with a horizontal resolution of 1/80th degree and 100 vertical levels (see Barnes (2024) for the source code).

Figure 1 shows the surface currents from a regional ocean simulation of the Tasman sea that was configured using the `regional_mom6` package.

## Statement of need

The learning curve for setting up a regional ocean model can be quite steep, and it is not obvious for a new user what inputs are required, nor the appropriate format. In the case of MOM6, there are several tools scattered in Github repositories, for example those collected in ESMG's grid tools (Cermak et al., 2021). Also, there exist several regional configuration examples but they are hardcoded for particular domains, specific input files, and work only on specific high-performance computing machines.

Until now there has been no one-stop-shop for users to learn how to get a regional MOM6 configuration up and running. Users are required to use several tools in several programming languages and then modify—sometimes by hand—some of the input metadata to bring everything into the format that MOM6 expects. Many parts of this process are not documented, requiring users to dig into the MOM6 Fortran source code. Other ocean models have packages to aid in regional configuration setup, for example Pyroms (Hedstrom & contributors, 2023) for the Regional Oceanic Modelling System (ROMS; Shchepetkin & McWilliams (2005)) and MITgcm\_python (Naughten & Jones, 2023) for the Massachusetts Institute of Technology

68 General Circulation Model (MITgcm; Marshall et al. (1997)). With MOM6's growing user  
69 base for regional applications, there is a need for a platform that walks users through regional  
70 domain configuration from start to finish and, ideally, automates the process on the way.  
71 `regional_mom6` fills precisely this need.

72 By having a shared set of tools that the community can work with and contribute to, this  
73 package also facilitates collaboration and knowledge-sharing between different research groups.  
74 Using a shared framework for setting up regional models, it is easier to compare and contrast  
75 examples of different experiments and allows for users to gain intuition for generating their  
76 chosen domain.

77 Another potential advantage of a package that allows users to automatically obtain regional  
78 configurations of MOM6 is in education. With the technically-challenging aspects of setting  
79 up a regional configuration now being automated by the `regional_mom6` package, students  
80 can set up and run simple MOM6 regional configurations and also change parameters like  
81 resolution or forcing, run again, and see how these parameters affect the ocean flow.

## 82 Acknowledgements

83 We thank the Consortium for Ocean–Sea Ice Modeling in Australia ([cosima.org.au](https://cosima.org.au)) for useful  
84 discussions during the development of this package. We acknowledge support from the  
85 Australian Research Council under DECRA Fellowship DE210100749 (N.C.C.) and grant  
86 LP200100406 (A.E.K.). We would also like to acknowledge the code and notes by James  
87 Simkins, Andrew Ross, and Rob Cermak, which helped us to troubleshoot and improve the  
88 algorithms in our package.

## 89 References

- 90 Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C. O., Dunne, J.  
91 P., Griffies, S. M., Hallberg, R., Harrison, M. J., Held, I. M., Jansen, M. F., John, J. G.,  
92 Krasting, J. P., Langenhorst, A. R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan,  
93 A., ... Zhang, R. (2019). The GFDL global ocean and sea ice model OM4.0: Model  
94 description and simulation features. *Journal of Advances in Modeling Earth Systems*,  
95 11(10), 3167–3211. <https://doi.org/10.1029/2019MS001726>
- 96 Arakawa, A., & Lamb, V. R. (1977). Computational design of the basic dynamical processes  
97 of the UCLA general circulation model. *Methods in Computational Physics: Advances*  
98 *in Research and Applications*, 17(Supplement C), 173–265. [https://doi.org/10.1016/](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)  
99 [B978-0-12-460817-7.50009-4](https://doi.org/10.1016/B978-0-12-460817-7.50009-4)
- 100 Barnes, A. J. (2024). Tasman-tides. In *GitHub repository*. [https://github.com/ashjbarnes/](https://github.com/ashjbarnes/tasman-tides)  
101 [tasman-tides](https://github.com/ashjbarnes/tasman-tides)
- 102 Cermak, R., Simkins, J., Hedstrom, K., & Gibson, A. (2021). ESMG gridtools. In *GitHub*  
103 *repository*. Github. <https://github.com/ESMG/gridtools>
- 104 Copernicus Climate Change Service. (2024). *ECMWF Reanalysis v5*. European Centre for  
105 Medium-Range Weather Forecasts. <https://doi.org/10.48670/moi-00021>
- 106 Copernicus Marine Services. (2024). *Global ocean physics reanalysis*. Mercator Ocean  
107 International. <https://doi.org/10.48670/moi-00021>
- 108 GEBCO Bathymetric Compilation Group 2023. (2023). *The GEBCO\_2023 Grid - a continuous*  
109 *terrain model of the global oceans and land*. NERC EDS British Oceanographic Data  
110 Centre NOC. <https://doi.org/10.5285/f98b053b-0cbc-6c23-e053-6c86abc0af7b>
- 111 Griffies, Stephen M. (2014). Elements of the modular ocean model (MOM). *GFDL Ocean*  
112 *Group Tech. Rep.*, 7, 47. [https://mom-ocean.github.io/assets/pdfs/MOM5\\_manual.pdf](https://mom-ocean.github.io/assets/pdfs/MOM5_manual.pdf)

- 113 Griffies, Stephen M., Adcroft, A., & Hallberg, R. W. (2020). A primer on the vertical  
114 lagrangian-remap method in ocean models based on finite volume generalized vertical  
115 coordinates. *Journal of Advances in Modeling Earth Systems*, 12(10), e2019MS001954.  
116 <https://doi.org/10.1029/2019MS001954>
- 117 Hedstrom, K., & contributors. (2023). Pyroms. In *GitHub repository*. Github. <https://github.com/ESMG/pyroms>  
118
- 119 Herzfeld, M., Schmidt, M., Griffies, S. M., & Liang, Z. (2011). Realistic test cases for limited  
120 area ocean modelling. *Ocean Modelling*, 37(1), 1–34. [https://doi.org/10.1016/j.ocemod.](https://doi.org/10.1016/j.ocemod.2010.12.008)  
121 [2010.12.008](https://doi.org/10.1016/j.ocemod.2010.12.008)
- 122 Marshall, J., Adcroft, A., Hill, C., Perelman, L., & Heisey, C. (1997). A finite-volume,  
123 incompressible Navier Stokes model for studies of the ocean on parallel computers. *Journal of*  
124 *Geophysical Research: Oceans*, 102(C3), 5753–5766. <https://doi.org/10.1029/96JC02775>
- 125 Naughten, K., & Jones, D. (2023). MITgcm\_python. In *GitHub repository*. Github. [https://github.com/knaughten/mitgcm\\_python](https://github.com/knaughten/mitgcm_python)  
126
- 127 Ross, A. C., Stock, C. A., Adcroft, A., Curchitser, E., Hallberg, R., Harrison, M. J., Hedstrom,  
128 K., Zadeh, N., Alexander, M., Chen, W., Drenkard, E. J., Pontavice, H. du, Dussin, R.,  
129 Gomez, F., John, J. G., Kang, D., Lavoie, D., Resplandy, L., Roobaert, A., ... Simkins, J.  
130 (2023). A high-resolution physical–biogeochemical model for marine resource applications in  
131 the northwest Atlantic (MOM6-COBALT-NWA12 v1.0). *Geoscientific Model Development*,  
132 16(23), 6943–6985. <https://doi.org/10.5194/gmd-16-6943-2023>
- 133 Ross, A. C., Stock, C. A., Koul, V., Delworth, T. L., Lu, F., Wittenberg, A., & Alexander,  
134 M. A. (2024). Dynamically downscaled seasonal ocean forecasts for North American East  
135 Coast ecosystems. *EGUsphere*, 2024, 1–40. <https://doi.org/10.5194/egusphere-2024-394>
- 136 Shchepetkin, A. F., & McWilliams, J. C. (2005). The regional oceanic modeling system  
137 (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model.  
138 *Ocean Modelling*, 9(4), 347–404. <https://doi.org/10.1016/j.ocemod.2004.08.002>
- 139 Zhuang, J., Dussin, R., Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich,  
140 B., Schupfner, M., Filipe, Levang, S., Gauthier, C., Jüling, A., Almansi, M., Scott,  
141 R., RondeauG, Rasp, S., Smith, T. J., Stachelek, J., Plough, M., & Li, X. (2023).  
142 xESMF: Universal regridding for geospatial data. In *GitHub repository*. Zenodo. <https://github.com/pangeo-data/xESMF>  
143