



Faculty of Science



Rantings on Landauer's principle and low-power electronic circuits

Michael Kirkedal Thomsen

Department of Computer Science
University of Copenhagen

m.kirkedal@di.ku.dk

Based mostly on others work

COST Training School
Toruń, Aug 31 2017



Rantings on Landauer's principle and low-power electronic circuits

Rantings on Landauer's principle and low-power electronic circuits



Why?

Rantings on Landauer's principle and low-power electronic circuits

Computers are not abstract:

Information is **inevitably physical**.

[Landauer in *Feynman and Computations* (1999)]



Why?

Rantings on Landauer's principle and low-power electronic circuits

Computers are not abstract:

Information is **inevitably physical**.

[Landauer in *Feynman and Computations* (1999)]

Every **logical state** of a device *must* correspond to a **physical state** of the device.

[Feynman in *Feynman Lectures on Computation* (1985)]



Why?

Rantings on Landauer's principle and low-power electronic circuits

Computers are not abstract:

Information is **inevitably physical**.

[Landauer in *Feynman and Computations* (1999)]

Every **logical state** of a device *must* correspond to a **physical state** of the device.

[Feynman in *Feynman Lectures on Computation* (1985)]

Whenever a device changes between a logical 0 and 1, there is a **corresponding change** in the physical device (e.g. memory or a transistor in an AND gate).



Thermodynamics of computation

Rantings on Landauer's principle and low-power electronic circuits

Landauer's principle

Any computation involving **loss of information** must **dissipate** a minimum amount of energy as heat.

[Landauer, 1961]



Thermodynamics of computation

Rantings on Landauer's principle and low-power electronic circuits

Landauer's principle

Any computation involving **loss of information** must **dissipate** a minimum amount of energy as heat.

[Landauer, 1961]

A physical device must dissipate energy of at least $k \cdot T \cdot \ln(2)$ joules per lost bit. Approx. $3 \cdot 10^{-21}$ J at room temp.



Thermodynamics of computation

Rantings on Landauer's principle and low-power electronic circuits

Landauer's principle

Any computation involving **loss of information** must **dissipate** a minimum amount of energy as heat.

[Landauer, 1961]

A physical device must dissipate energy of at least $k \cdot T \cdot \ln(2)$ joules per lost bit. Approx. $3 \cdot 10^{-21}$ J at room temp.

- Gedankenexperiment has been experimentally verified.

[Bérut et al., 2012]



Thermodynamics of computation

Rantings on Landauer's principle and low-power electronic circuits

Landauer's principle

Any computation involving **loss of information** must **dissipate** a minimum amount of energy as heat.

[Landauer, 1961]

A physical device must dissipate energy of at least $k \cdot T \cdot \ln(2)$ joules per lost bit. Approx. $3 \cdot 10^{-21}$ J at room temp.

- Gedankenexperiment has been experimentally verified.

[Bérut et al., 2012]

- Can reversibly modify electrical signal with dissipation below Landauer's limit.

[Orlov et al., 2012]



Thermodynamics of computation

Reflections on Landauer's principle and low-power electronic circuits

Landauer's principle

Any computation involving **loss of information** must **dissipate** a minimum amount of energy as heat.

[Landauer, 1961]

A physical device must dissipate energy of at least $k \cdot T \cdot \ln(2)$ joules per lost bit. Approx. $3 \cdot 10^{-21}$ J at room temp.

- ▶ Gedankenexperiment has been experimentally verified.

[Bérut et al., 2012]

- ▶ Can reversibly modify electrical signal with dissipation below Landauer's limit.

[Orlov et al., 2012]

- ▶ Actual amount of energy is correlated with signal representation.

[Gershenfeld, 1996]



Landauer vs. Moore

Rantings on Landauer's principle and low-power electronic circuits

"You can spend a lot of time arguing about a sensible value but something like the following is not too unreasonable."

The Landauer switching limit at finite (GHz) clock speed:
Energy to switch 1 bit $> 100k_B T \ln(2)$

[Douglas J. Poul, url]




Landauer vs. Moore

Rantings on Landauer's principle and low-power electronic circuits

"You can spend a lot of time arguing about a sensible value but something like the following is not too unreasonable."

The Landauer switching limit at finite (GHz) clock speed:
Energy to switch 1 bit $> 100k_B T \ln(2)$

[Douglas J. Poul, url]





Transistor-based CMOS implementations I

Rantings on Landauer's principle and low-power electronic circuits

Switch-level charge-recovery logic

- Adiabatic switching
- Needs many clocks signal – hard to control
- Use compute-uncompute for energy restoration – not really reversible

[Younis and Knight, 1994]





Transistor-based CMOS implementations II

Rantings on Landauer's principle and low-power electronic circuits

Complementary pass-transistor logic

- ▶ Adiabatic switching
- ▶ Degrading signals makes control of transistors hard over deep circuits
- ▶ Fully reversible implementation (dual-line ensures conservatism)


[De Vos and Van Rentergem, 2005, Burignat and De Vos, 2012]



Future Technologies - Materials and models

Rantings on Landauer's principle and low-power electronic circuits

2D materials




The research community exploring 2D materials beyond graphene is rapidly expanding

with new entrants from different fields such as materials science, physics, chemistry and electrical engineering.

Encouraging work has already been intensively reported from different material systems such as nitrides (e.g., h-BN), dichalcogenides (e.g., MoS₂), topological insulators (e.g., Bi₂Se₃ or Bi₂Te₃) and even oxides.

- Low energy consumption
- Low switching time

IBM: Liquid transistor



Gold electrical contacts surround a semiconducting material called vanadium oxide (brownish color). When a voltage is applied to ions in


electrolyte (not seen) in contact with the material, it switches the material's state from insulating to conducting.



Future Technologies - Non-transistor models

Rantings on Landauer's principle and low-power electronic circuits


(Super) Capacitive Adiabatic Logic



Future Technologies - Non-transistor models

Rantings on Landauer's principle and low-power electronic circuits

(Super) Capacitive Adiabatic Logic




Micro-Electro-Mechanical Systems (MEMS)



Future Technologies - Non-transistor models

Rantings on Landauer's principle and low-power electronic circuits

(Super) Capacitive Adiabatic Logic





Micro-Electro-Mechanical Systems (MEMS)



Quantum Computing

- Related circuit model
- Step in synthesis of q-circuits



Thank you

?



Bibliography I



Bérut, A., Arakelyan, A., Petrosyan, A., Ciliberto, S., Dillenschneider, R., and Lutz, E. (2012).
Experimental verification of Landauer's principle linking information and thermodynamics.
Nature, 483(7388):187–189.



Burignat, S. and De Vos, A. (2012).
A review on performances of reversible ripple-carry adders.
International Journal of Electronics and Telecommunications, 58(3):205–212.



De Vos, A. and Van Rentergem, Y. (2005).
Reversible computing: from mathematical group theory to electronical circuit experiment.
In *Proceedings of the Conference on Computing Frontiers*, pages 35–44. ACM Press.



Gershenfeld, N. (1996).
Signal entropy and the thermodynamics of computation.
IBM Systems Journal, 35(3.4):577–586.



Landauer, R. (1961).
Irreversibility and heat generation in the computing process.
IBM Journal of Research and Development, 5(3):183–191.



Orlov, A. O., Lent, C. S., Thorpe, C. C., Boechler, G. P., and Snider, G. L. (2012).
Experimental test of Landauer's principle at the sub- $k_B t$ level.
Japanese Journal of Applied Physics, 51:06FE10.



Poul, D. J. (url).
The fundamental limits of computation.
<http://userweb.eng.gla.ac.uk/douglas.paul/SiGe/limits.html>.



Bibliography II



Younis, S. G. and Knight, J. T. F. (1994).

Asymptotically zero energy computing split-level charge recovery logic.
International Workshop on Low Power Design, pages 177–182.

