


Statistical Inference

Jean Daunizeau


Wellcome Trust Centre for Neuroimaging
University College London

SPM Course
Edinburgh, April 2010

Image time-series




Spatial filter



Realignment


Smoothing

Normalisation




Anatomical
reference

Design matrix






General Linear Model




Parameter estimates

Statistical Parametric Map




Statistical
Inference

RFT



$p < 0.05$


Voxel-wise time series analysis



Overview

- ❑ Model specification and parameters estimation
- ❑ Hypothesis testing
- ❑ Contrasts
 - T -tests
 - F -tests
- ❑ Contrast estimability
- ❑ Correlation between regressors
 - Example(s)
- ❑ Design efficiency

Model Specification: The General Linear Model



N : number of scans, p : number of regressors

$$y = X\beta + \varepsilon$$

Sphericity assumption:
Independent and identically distributed (i.i.d.) error terms

$$\varepsilon \sim N(0, \sigma^2 I)$$

□ The General Linear Model is an equation that expresses the observed response variable in terms of a linear combination of explanatory variables X plus a well behaved error term. Each column of the design matrix corresponds to an effect one has built into the experiment or that may confound the results.

Parameter Estimation: Ordinary Least Squares

- Find $\hat{\beta}$ that minimises

$$\|y - X\beta\|^2 = \varepsilon^T \varepsilon$$

- The Ordinary Least Estimates are:

$$\hat{\beta} = (X^T X)^{-1} X^T y$$

- Under i.i.d. assumptions, the Ordinary Least Squares estimates are Maximum Likelihood.

$$\varepsilon \sim N(0, \sigma^2 I) \longrightarrow Y \sim N(X\beta, \sigma^2 I)$$

$$\hat{\sigma}^2 = \frac{\hat{\varepsilon}^T \hat{\varepsilon}}{N - p}$$

$$\hat{\beta} \sim N(\beta, \sigma^2 (X^T X)^{-1})$$

Hypothesis Testing

To test an hypothesis, we construct “test statistics”.

❑ The Null Hypothesis H_0

Typically what we want to disprove (no effect).


⇒ The Alternative Hypothesis H_A expresses outcome of interest.

❑ The Test Statistic T

The test statistic summarises evidence about H_0 .

Typically, test statistic is small in magnitude when the hypothesis H_0 is true and large when false.

⇒ We need to know the distribution of T under the null hypothesis.



Null Distribution of T

Hypothesis Testing


□ Significance level α :

Acceptable *false positive rate* α .

\Rightarrow threshold u_α

Threshold u_α controls the false positive rate

$$\alpha = p(T > u_\alpha \mid H_0)$$



Null Distribution of T

Observation of test statistic t , a realisation of T


□ The conclusion about the hypothesis:

We reject the null hypothesis in favour of the alternative hypothesis if $t > u_\alpha$

□ P-value:

A p -value summarises evidence against H_0 .

This is the chance of observing value more extreme than t under the null hypothesis.




Null Distribution of T

$$p(T > t \mid H_0)$$

Contrasts

- ❑ We are usually not interested in the whole β vector.
- ❑ A contrast selects a specific effect of interest:
 - \Rightarrow a contrast c is a vector of length p .
 - $\Rightarrow c^T \beta$ is a linear combination of regression coefficients β .



$$c^T = [1 \ 0 \ 0 \ 0 \ 0 \ \dots]$$

$$c^T \beta = 1 \times \beta_1 + 0 \times \beta_2 + 0 \times \beta_3 + 0 \times \beta_4 + 0 \times \beta_5 + \dots$$

$$c^T = [0 \ -1 \ 1 \ 0 \ 0 \ \dots]$$

$$c^T \beta = 0 \times \beta_1 + -1 \times \beta_2 + 1 \times \beta_3 + 0 \times \beta_4 + 0 \times \beta_5 + \dots$$


- ❑ Under i.i.d assumptions:

$$c^T \hat{\beta} \sim N(c^T \beta, \sigma^2 c^T (X^T X)^{-1} c)$$

T-test - one dimensional contrasts – SPM{t}

$$c^T = 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0$$

$\beta_1 \ \beta_2 \ \beta_3 \ \beta_4 \ \beta_5 \ \dots$



Question: box-car amplitude > 0 ?
=
 $\beta_1 = c^T \beta > 0 ?$

Null hypothesis:

$$H_0: c^T \beta = 0$$



*contrast of
estimated
parameters*


Test statistic:

$$T = \frac{\text{contrast of estimated parameters}}{\sqrt{\text{variance estimate}}}$$

$$T = \frac{c^T \hat{\beta}}{\sqrt{\text{var}(c^T \hat{\beta})}} = \frac{c^T \hat{\beta}}{\sqrt{\hat{\sigma}^2 c^T (X^T X)^{-1} c}} \sim t_{N-p}$$


T-contrast in SPM

□ For a given contrast c :




beta_???? images

$$\hat{\beta} = (X^T X)^{-1} X^T y$$




ResMS image

$$\hat{\sigma}^2 = \frac{\hat{\varepsilon}^T \hat{\varepsilon}}{N - p}$$



con_???? image

$$c^T \hat{\beta}$$




spmT_???? image

$$\text{SPM}\{t\}$$

T-test: a simple example

❑ Passive word listening versus rest


$$C^T = \begin{bmatrix} 1 & 0 \end{bmatrix}$$



Q: activation during listening ?

Null hypothesis: $\beta_1 = 0$

$$t = \frac{c^T \hat{\beta}}{Std(c^T \hat{\beta})}$$



SPMresults:

Height threshold $T = 3.2057$ $\{p < 0.001\}$



voxel-level			mm mm mm		
T	(Z)	$p_{\text{uncorrected}}$			
1	1	0.000000	1	1	1
2	2	0.000000	2	2	2
3	3	0.000000	3	3	3
4	4	0.000000	4	4	4
5	5	0.000000	5	5	5
6	6	0.000000	6	6	6
7	7	0.000000	7	7	7
8	8	0.000000	8	8	8
9	9	0.000000	9	9	9
10	10	0.000000	10	10	10
11	11	0.000000	11	11	11
12	12	0.000000	12	12	12
13	13	0.000000	13	13	13
14	14	0.000000	14	14	14
15	15	0.000000	15	15	15
16	16	0.000000	16	16	16
17	17	0.000000	17	17	17
18	18	0.000000	18	18	18
19	19	0.000000	19	19	19
20	20	0.000000	20	20	20
21	21	0.000000	21	21	21
22	22	0.000000	22	22	22
23	23	0.000000	23	23	23
24	24	0.000000	24	24	24
25	25	0.000000	25	25	25
26	26	0.000000	26	26	26
27	27	0.000000	27	27	27
28	28	0.000000	28	28	28
29	29	0.000000	29	29	29
30	30	0.000000	30	30	30
31	31	0.000000	31	31	31
32	32	0.000000	32	32	32
33	33	0.000000	33	33	33
34	34	0.000000	34	34	34
35	35	0.000000	35	35	35
36	36	0.000000	36	36	36
37	37	0.000000	37	37	37
38	38	0.000000	38	38	38
39	39	0.000000	39	39	39
40	40	0.000000	40	40	40
41	41	0.000000	41	41	41
42	42	0.000000	42	42	42
43	43	0.000000	43	43	43
44	44	0.000000	44	44	44
45	45	0.000000	45	45	45
46	46	0.000000	46	46	46
47	47	0.000000	47	47	47
48	48	0.000000	48	48	48
49	49	0.000000	49	49	49
50	50	0.000000	50	50	50
51	51	0.000000	51	51	51
52	52	0.000000	52	52	52
53	53	0.000000	53	53	53
54	54	0.000000	54	54	54
55	55	0.000000	55	55	55
56	56	0.000000	56	56	56
57	57	0.000000	57	57	57
58	58	0.000000	58	58	58
59	59	0.000000	59	59	59
60	60	0.000000	60	60	60
61	61	0.000000	61	61	61
62	62	0.000000	62	62	62
63	63	0.000000	63	63	63
64	64	0.000000	64	64	64
65	65	0.000000	65	65	65
66	66	0.000000	66	66	66
67</					

[illegible]

T-test: a few remarks

- T-test is a signal-to-noise measure (ratio of estimate to standard deviation of estimate).
- T-contrasts are simple combinations of the betas; the T-statistic does not depend on the scaling of the regressors or the scaling of the contrast.
- Unilateral test:
$$H_0: c^T \beta = 0 \quad \text{vs} \quad H_A: c^T \beta > 0$$

Scaling issue



$$T = \frac{c^T \hat{\beta}}{\sqrt{\text{var}(c^T \hat{\beta})}} = \frac{c^T \hat{\beta}}{\sqrt{\hat{\sigma}^2 c^T (X^T X)^{-1} c}}$$


- ❑ The T -statistic does not depend on the scaling of the regressors.
- ❑ The T -statistic does not depend on the scaling of the contrast.
- ❑ Contrast $c^T \hat{\beta}$ depends on scaling.
- Be careful of the interpretation of the contrasts $c^T \hat{\beta}$ themselves (eg, for a second level analysis):

sum \neq average

F-test - the extra-sum-of-squares principle

- Model comparison: *Full vs. Reduced model?*

Null Hypothesis H_0 : True model is X_0 (reduced model)




F-test - multidimensional contrasts – SPM{ F }

❑ Tests multiple linear hypotheses:

H_0 : True model is X_0

$H_0: \beta_3 = \beta_4 = \dots = \beta_9 = 0$






test $H_0: c^T \beta = 0$?



Full model?


Reduced model?

$$c^T = \begin{matrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{matrix}$$




SPM{ $F_{6,322}$ }

F-contrast in SPM




beta_???? images

$$\hat{\beta} = (X^T X)^{-1} X^T y$$




ResMS image

$$\hat{\sigma}^2 = \frac{\hat{\varepsilon}^T \hat{\varepsilon}}{N - p}$$



ess_???? images




$$(RSS_0 - RSS)$$




spmF_???? images

$$SPM\{F\}$$

F-test example: movement related effects



Multidimensional contrasts



Think of it as constructing 3 regressors from the 3 differences and complement this new design matrix such that data can be fitted in the same exact way (same error, same fitted data).

***F*-test: a few remarks**

- ❑ *F*-tests can be viewed as testing for the additional variance explained by a larger model wrt a simpler (nested) model \Rightarrow Model comparison.
- ❑ *F* tests a weighted sum of squares of one or several combinations of the regression coefficients β .
- ❑ In practice, we don't have to explicitly separate X into $[X_1 X_2]$ thanks to multidimensional contrasts.

❑ Hypotheses:

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Null Hypothesis $H_0 : \beta_1 = \beta_2 = \beta_3 = 0$

Alternative Hypothesis $H_A : \text{at least one } \beta_k \neq 0$

- ❑ In testing uni-dimensional contrast with an *F*-test, for example $\beta_1 - \beta_2$, the result will be the same as testing $\beta_2 - \beta_1$. It will be exactly the square of the *t*-test, testing for both positive and negative effects.


Estimability of a contrast

- If X is not of **full rank** then we can have $X\beta_1 = X\beta_2$ with $\beta_1 \neq \beta_2$ (different parameters).
- The parameters are **not** therefore 'unique', 'identifiable' or '**estimable**'.
- For such models, $X^T X$ is not invertible so we must resort to generalised inverses (SPM uses the **pseudo-inverse**).

□ Example:


$[1 \ 0 \ 0]$, $[0 \ 1 \ 0]$, $[0 \ 0 \ 1]$ are not estimable.

$[1 \ 0 \ 1]$, $[0 \ 1 \ 1]$, $[1 \ -1 \ 0]$, $[0.5 \ 0.5 \ 1]$ are estimable.



Design orthogonality

Statistical analysis: Design orthogonality



- For each pair of columns of the design matrix, the orthogonality matrix depicts the magnitude of the **cosine of the angle** between them, with the range 0 to 1 mapped from white to black.
- The cosine of the angle between two vectors a and b is obtained by:

$$\cos \alpha = \frac{a \cdot b}{\|a\| \|b\|}$$


- If both vectors have **zero mean** then the cosine of the angle between the vectors is the same as the **correlation** between the two variates.

Multicollinearity

$$\text{Var}(c^T \hat{\beta}) = \sigma^2 c^T (X^T X)^{-1} c$$

□ Orthogonal regressors (=uncorrelated):

By varying each separately, one can predict the combined effect of varying them jointly.






$(X^T X)^{-1}$ is diagonal



□ Non-orthogonal regressors (=correlated):


When testing for the first regressor, we are effectively removing the part of the signal that can be accounted for by the second regressor \Rightarrow *implicit orthogonalisation*.



$$x_2^\perp = x_2 - x_1 \cdot x_2 \cdot x_1$$


It does not reduce the predictive power or reliability of the model as a whole.

Shared variance




Orthogonal regressors.

Shared variance



*Testing for the **green**:*




Correlated regressors, for example:


green: subject age

yellow: subject score

Shared variance




*Testing for the **red**:*




Correlated regressors.

Shared variance





Testing for the *green*:



Highly correlated.
Entirely correlated \Rightarrow non estimable

Shared variance

Testing for the green and yellow



If significant, can be G and/or Y

Examples

A few remarks

- ❑ We implicitly test for an additional effect only, be careful if there is correlation
 - Orthogonalisation = decorrelation : not generally needed
 - Parameters and test on the non modified regressor change

- ❑ It is always simpler to have orthogonal regressors and therefore designs.

- ❑ In case of correlation, use F-tests to see the overall significance. There is generally no way to decide to which regressor the « common » part should be attributed to.

- ❑ Original regressors may not matter: it's the contrast you are testing which should be as decorrelated as possible from the rest of the design matrix

Design efficiency

- The aim is to minimize the standard error of a t -contrast (i.e. the denominator of a t -statistic).

$$T = \frac{c^T \hat{\beta}}{\sqrt{\text{var}(c^T \hat{\beta})}}$$

$$\text{var}(c^T \hat{\beta}) = \hat{\sigma}^2 c^T (X^T X)^{-1} c$$

- This is equivalent to maximizing the efficiency e :

$$e(\hat{\sigma}^2, c, X) = (\hat{\sigma}^2 c^T (X^T X)^{-1} c)^{-1}$$

Noise variance

Design variance


- If we assume that the noise variance is independent of the specific design:

$$e(c, X) = (c^T (X^T X)^{-1} c)^{-1}$$

- This is a relative measure: all we can really say is that one design is more efficient than another (for a given contrast).

Design efficiency

- ❑ The efficiency of an estimator is a measure of how reliable it is and depends on error variance (the variance not modeled by explanatory variables in the design matrix) and the design variance (a function of the explanatory variables and the contrast tested).
- ❑ $X^T X$ represents covariance of regressors in design matrix; high covariance increases elements of $(X^T X)^{-1}$.
- ❑ High correlation between regressors leads to low sensitivity to each regressor alone.



$$\begin{bmatrix} 1 & -0.9 \\ -0.9 & 1 \end{bmatrix}$$

$$c^T (X^T X)^{-1} c$$


$$c^T = [1 \ 0]: \quad 5.26$$

$$c^T = [1 \ 1]: \quad 20$$

$$c^T = [1 \ -1]: \quad 1.05$$

Bibliography:

- ❑ *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. Elsevier, 2007.
- ❑ *Plane Answers to Complex Questions: The Theory of Linear Models*. R. Christensen, Springer, 1996.
- ❑ *Statistical parametric maps in functional imaging: a general linear approach*. K.J. Friston et al, Human Brain Mapping, 1995.
- ❑ *Ambiguous results in functional neuroimaging data analysis due to covariate correlation*. A. Andrade et al., NeuroImage, 1999.
- ❑ Estimating efficiency a priori: a comparison of blocked and randomized designs. A. Mechelli et al., NeuroImage, 2003.



With many thanks to G. Flandin, J.-B. Poline and Tom Nichols for slides.