

Bayesian inference

J. Daunizeau

Brain and Spine Institute, Paris, France

Wellcome Trust Centre for Neuroimaging, London, UK

Overview of the talk

1 Probabilistic modelling and representation of uncertainty

1.1 Bayesian paradigm

1.2 Hierarchical models

1.3 Frequentist versus Bayesian inference

2 Numerical Bayesian inference methods

2.1 Sampling methods

2.2 Variational methods (ReML, EM, VB)

3 SPM applications

3.1 aMRI segmentation

3.2 Decoding of brain images

3.3 Model-based fMRI analysis (with spatial priors)

3.4 Dynamic causal modelling

Overview of the talk

1 Probabilistic modelling and representation of uncertainty

1.1 Bayesian paradigm

1.2 Hierarchical models

1.3 Frequentist versus Bayesian inference

2 Numerical Bayesian inference methods

2.1 Sampling methods

2.2 Variational methods (ReML, EM, VB)

3 SPM applications

3.1 aMRI segmentation

3.2 Decoding of brain images

3.3 Model-based fMRI analysis (with spatial priors)


3.4 Dynamic causal modelling

Bayesian paradigm

probability theory: basics

Degree of *plausibility* desiderata:

- should be represented using real numbers (D1)
- should conform with intuition (D2)
- should be consistent (D3)



• normalization: $\sum_a P(a) = 1$


• marginalization: $P(b) = \sum_a P(a, b)$

• **conditioning :**
(Bayes rule)

$$P(a, b) = P(a|b) P(b)$$
$$= P(b|a) P(a)$$


Bayesian paradigm

deriving the likelihood function



Bayesian paradigm


likelihood, priors and the model evidence



Likelihood: $p(y|\theta, m)$


Prior: $p(\theta|m)$

Bayes rule:
$$p(\theta|y, m) = \frac{p(y|\theta, m) p(\theta|m)}{p(y|m)}$$



Bayesian paradigm

forward and inverse problems





Bayesian paradigm

model comparison

Principle of parsimony :


« plurality should not be assumed without necessity »



Model evidence:


$$p(y|m) = \int p(y|\theta, m) p(\theta|m) d\theta$$

“Occam’s razor” :




Hierarchical models

principle




\vdots

$$p(\theta_2 | \theta_3, m)$$



$$p(\theta_1 | \theta_2, m)$$



$$p(y | \theta_1, m)$$

inference







causality



Hierarchical models

directed acyclic graphs (DAGs)



$$p(\theta_1 | \theta_2, u, m)$$


$$p(y | \theta_1, \sigma^2, m)$$

$$p(\theta | m) = \prod_j p(\theta_j | \text{par}(\theta_j), m)$$

Frequentist versus Bayesian inference

a (quick) note on hypothesis testing

- define the null, e.g.: $H_0 : \theta = 0$



- estimate parameters (obtain test stat.)
- apply decision rule, i.e.:

if $P(t > t^* | H_0) \leq \alpha$ then reject H_0

classical (null) hypothesis testing


Family-level inference

trading model resolution against statistical power


model selection error risk:

$$P(e = 1|y) = 1 - \max_m P(m|y) \\ = 0.3$$


$$P(m_1|y) = 0.04$$




$$P(m_2|y) = 0.25$$



$$P(m_2|y) = 0.01$$




$$P(m_2|y) = 0.7$$




Family-level inference

trading inference resolution against statistical power



$$P(f_1|y) = 0.05$$



$$P(f_2|y) = 0.95$$

model selection error risk:

$$P(e=1|y) = 1 - \max_m P(m|y) = 0.3$$


family inference
(pool statistical evidence)

$$P(f|y) = \sum_{m \in f} P(m|y)$$

$$P(e=1|y) = 1 - \max_f P(f|y) = 0.05$$


Group-level model comparison

preliminary: Polya's urn



$$\begin{cases} m_i = 1 & \rightarrow i^{\text{th}} \text{ marble is blue} \\ m_i = 0 & \rightarrow i^{\text{th}} \text{ marble is purple} \end{cases}$$

r = proportion of blue marbles in the urn



→ (binomial) probability of drawing a set of n marbles:

$$p(m|r) = \prod_{i=1}^n r^{m_i} (1-r)^{1-m_i}$$


Thus, our belief about the proportion of blue marbles is:

$$p(r|m) \propto p(r) \prod_{i=1}^n r^{m_i} (1-r)^{1-m_i} \quad \xRightarrow{p(r) \propto 1} \quad E[r|m] = \frac{1}{n} \sum_{i=1}^n m_i$$


Group-level model comparison

what if we are colour blind?

At least, we can measure how likely is the i^{th} subject's data under each model!



$$p(y_1|m_1) \quad p(y_2|m_2) \quad \dots \quad p(y_i|m_i) \quad \dots \quad p(y_n|m_n)$$



$$p(r, m|y) \propto p(r) \prod_{i=1}^n p(y_i|m_i) p(m_i|r)$$

Our belief about the proportion of models is:

$$p(r|y) = \sum_m p(r, m|y)$$

Exceedance probability: $\varphi_k = P(r_k > r_{k' \neq k} | y)$

Overview of the talk

1 Probabilistic modelling and representation of uncertainty

1.1 Bayesian paradigm

1.2 Hierarchical models

1.3 Frequentist versus Bayesian inference

2 Numerical Bayesian inference methods

2.1 Sampling methods

2.2 Variational methods (ReML, EM, VB)

3 SPM applications

3.1 aMRI segmentation


3.2 Decoding of brain images

3.3 Model-based fMRI analysis (with spatial priors)


3.4 Dynamic causal modelling

Sampling methods

MCMC example: Gibbs sampling



$$p(\theta_i|y, \theta_{j \neq i}) \propto \underbrace{p(\theta_i|par(\theta_i))}_{\text{prior}} \underbrace{\prod_{j=ch(i)} p(\theta_j|par(\theta_j))}_{\text{likelihood}}$$




$$\frac{1}{N} \sum_{n=1}^N p(y|\theta^{(n)}, m) \approx p(y|m)$$

Variational methods

VB / EM / ReML

$$\ln p(y|m) = \underbrace{\langle \ln p(\theta, y|m) \rangle_q}_{\text{free energy } F(q)} + S(q) + D_{KL}(q(\theta); p(\theta|y, m))$$

→ **VB** : maximize the **free energy** $F(q)$ w.r.t. the **approximate posterior** $q(\theta)$ under some (e.g., *mean field*, *Laplace*) simplifying constraint



Overview of the talk

1 Probabilistic modelling and representation of uncertainty

1.1 Bayesian paradigm

1.2 Hierarchical models

1.3 Frequentist versus Bayesian inference

2 Numerical Bayesian inference methods

2.1 Sampling methods

2.2 Variational methods (ReML, EM, VB)

3 SPM applications

3.1 aMRI segmentation

3.2 Decoding of brain images

3.3 Model-based fMRI analysis (with spatial priors)






3.4 Dynamic causal modelling

segmentation
and normalisation

posterior probability
maps (PPMs)

dynamic causal
modelling


multivariate
decoding




realignment

smoothing


general linear model



normalisation




template



statistical
inference




Gaussian
field theory

$p < 0.05$





aMRI segmentation

mixture of Gaussians (MoG) model




Decoding of brain images


recognizing brain states from fMRI



log-evidence of X-Y sparse mappings:
effect of lateralization





log-evidence of X-Y bilateral mappings:
effect of spatial deployment




fMRI time series analysis



spatial priors and model comparison




short-term memory
design matrix (X)




PPM: regions best explained
by short-term memory model



long-term memory
design matrix (X)






PPM: regions best explained
by long-term memory model





Dynamic Causal Modelling

network structure identification



DCMs and DAGs

a note on causality



I thank you for your attention.

A note on statistical significance

lessons from the Neyman-Pearson lemma


- **Neyman-Pearson lemma**: the likelihood ratio (or Bayes factor) test

$$\Lambda = \frac{p(y|H_1)}{p(y|H_0)} \geq u$$

is the most powerful test of size $\alpha = p(\Lambda \geq u | H_0)$ to test the null.

- what is the threshold u , above which the Bayes factor test yields a error I rate of 5%?

ROC analysis



MVB (Bayes factor)
 $u=1.09$, power=56%

CCA (F-statistics)
 $F=2.20$, power=20%