



- 29/317 nucleoside–triphosphatase regulator
- 4/79 unfolded protein binding
- 7/55 oxidoreductase activity, acting on a sulfur group of donors
- 11/152 gated channel
- 8/160 actin binding
- 26/502 DNA–binding transcription factor
- 2/14 metalloaminopeptidase
- 4/32 tumor necrosis factor receptor superfamily binding
- 5/38 oxidation–driven active transmembrane transporter
- 12/124 structural constituent of ribosome
- 5/29 myosin complex
- 40/582 plasma membrane region
- 3/22 intraciliary transport particle
- 5/82 peptidase complex
- 12/95 endoplasmic reticulum protein–containing complex
- 10/213 transferase complex, transferring phosphorus–containing groups
- 6/59 respiratory chain complex
- 13/92 cytosolic ribosome
- 2/28 spliceosomal tri–snRNP complex
- 1/15 response to anoxia
- 24/305 G protein–coupled receptor signaling pathway
- 20/133 response to topologically incorrect protein
- 9/89 cytoplasmic translation
- 6/90 oxidative phosphorylation
- 4/29 intraciliary transport involved in cilium assembly
- 3/14 folic acid metabolic process
- 3/15 cellular response to exogenous dsRNA
- 0/17 negative regulation of DNA–templated transcription, elongation