

# Methods of Mathematical Physics

## — Lecture 5 Singularities, Residue Theory —

Lei Du

dulei@dlut.edu.cn

<http://faculty.dlut.edu.cn/dulei>

School of Mathematical Sciences  
Dalian University of Technology

April, 2022

- 1 Zeros of Analytic Functions
- 2 Singular Points
- 3 Residue Theorem

1 Zeros of Analytic Functions

2 Singular Points

3 Residue Theorem

# Zeros

In this lecture, we introduce certain basic results on the "zeros" of an analytic function. But, first we have the following definition:

## Definition

A zero of an analytic function  $f(z)$  is any value of  $z$  for which  $f(z)$  vanishes.

# Zeros

In this lecture, we introduce certain basic results on the "zeros" of an analytic function. But, first we have the following definition:

## Definition

A zero of an analytic function  $f(z)$  is any value of  $z$  for which  $f(z)$  vanishes.

Let  $f(z)$  be analytic in a domain  $D$  and let  $a$  be a point of  $D$ . Then  $f(z)$  can be expanded as a Taylor series about  $z = a$  in the form

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n, \text{ where } a_n = \frac{f^{(n)}(a)}{n!}.$$

If  $f(a) = 0$ , i.e.,  $a$  is a zero of  $f(z)$ , we have  $a_0 = 0$ .

# Zeros

In this lecture, we introduce certain basic results on the "zeros" of an analytic function. But, first we have the following definition:

## Definition

A zero of an analytic function  $f(z)$  is any value of  $z$  for which  $f(z)$  vanishes.

Let  $f(z)$  be analytic in a domain  $D$  and let  $a$  be a point of  $D$ . Then  $f(z)$  can be expanded as a Taylor series about  $z = a$  in the form

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n, \text{ where } a_n = \frac{f^{(n)}(a)}{n!}.$$

If  $f(a) = 0$ , i.e.,  $a$  is a zero of  $f(z)$ , we have  $a_0 = 0$ .

It may also happen that more of the coefficients  $a_n$  vanish. If  $a_n = 0$  for  $n < m$ , but  $a_m \neq 0$ , then we have

$$\begin{aligned} f(z) &= \sum_{n=m}^{\infty} a_n(z-a)^n = \sum_{n=0}^{\infty} a_{n+m}(z-a)^{m+n} = (z-a)^m \sum_{n=0}^{\infty} a_{n+m}(z-a)^n \\ &= (z-a)^m \phi(z), \end{aligned}$$

where  $\phi(z) = \sum_{n=0}^{\infty} a_{n+m}(z-a)^n$  is analytic within the region of convergence of Taylor's expansion of  $f(z)$  and  $\phi(a) \neq 0$ .

# Zeros

In such case, we say that  $f(z)$  has a zero of order  $m$  at  $z = a$ . A zero of order one is said to be a simple zero. If  $a$  is a zero of  $f(z)$  of order  $m$ , then we have

$$f(a) = 0, \quad f'(a) = \dots = f^{(m-1)}(a) = 0,$$

but  $f^{(m)}(a) \neq 0$ . This is obvious from Taylor's expansion formula.

# Zeros

In such case, we say that  $f(z)$  has a zero of order  $m$  at  $z = a$ . A zero of order one is said to be a simple zero. If  $a$  is a zero of  $f(z)$  of order  $m$ , then we have

$$f(a) = 0, \quad f'(a) = \dots = f^{(m-1)}(a) = 0,$$

but  $f^{(m)}(a) \neq 0$ . This is obvious from Taylor's expansion formula. Now, we have an important result.

## Theorem

*Zeros are isolated points.*



# Zeros

In such case, we say that  $f(z)$  has a zero of order  $m$  at  $z = a$ . A zero of order one is said to be a simple zero. If  $a$  is a zero of  $f(z)$  of order  $m$ , then we have

$$f(a) = 0, \quad f'(a) = \cdots = f^{(m-1)}(a) = 0,$$

but  $f^{(m)}(a) \neq 0$ . This is obvious from Taylor's expansion formula. Now, we have an important result.

## Theorem

*Zeros are isolated points.*

## Proof.

Let  $f(z)$  be analytic in a domain  $D$ . Then we show that, unless  $f(z)$  is identically zero, there exists a neighborhood of each point in  $D$  throughout which the function has no zero, except possibly at the point itself. Suppose that  $f(z)$  has a zero of order  $m$  at  $a$ . Then, as above,

$$f(z) = (z - a)^m \sum_{n=0}^{\infty} a_{m+n}(z - a)^n = (z - a)^m \phi(z). \quad (1)$$

Now, we have

$$\phi(z) = \sum_{n=0}^{\infty} a_{m+n}(z - a)^n \text{ and } \phi(a) = a_m \neq 0.$$

Since the series in (1) is uniformly convergent and each term of the series is continuous at  $a$ , it follows that  $\phi(z)$ , being a sum function, is also continuous at  $a$ . Hence, for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that

$$|z - a| < \delta \implies |\phi(z) - \phi(a)| < \epsilon. \quad (2)$$

Take  $\epsilon = \frac{|a_m|}{2}$  and let  $\delta_0$  be the corresponding value of  $\delta$ . Then (2) gives

$$|z - a| < \delta_0 \implies |\phi(z) - a_m| = |\phi(z) - \phi(a)| < \frac{1}{2} |a_m|. \quad (3)$$

It follows that  $\phi(a) \neq 0$  at any point in the neighborhood  $|z - a| < \delta_0$ . For, if  $\phi(z) = 0$ , then the equality (3) is contradicted. The argument remains valid when  $m = 0$ . In this case, the two functions  $\phi$  and  $f$  are equal and  $f(a) \neq 0$ . □

1 Zeros of Analytic Functions

2 **Singular Points**

3 Residue Theorem

# Definitions

In this section, we introduce the functions which are analytic at all points of a bounded domain except at a finite number of points. Such exceptional points are called singular points or singularities.

# Definitions

In this section, we introduce the functions which are analytic at all points of a bounded domain except at a finite number of points. Such exceptional points are called singular points or singularities.

## Definition (Singular Points)

A singular point of a function  $f(z)$  is the point at which the function ceases to be analytic.

For example, the function  $f(z) = \frac{1}{z-1}$  has a singularity at  $z = 1$ .

# Definitions

In this section, we introduce the functions which are analytic at all points of a bounded domain except at a finite number of points. Such exceptional points are called singular points or singularities.

## Definition (Singular Points)

A singular point of a function  $f(z)$  is the point at which the function ceases to be analytic.

For example, the function  $f(z) = \frac{1}{z-1}$  has a singularity at  $z = 1$ .

## Definition (Isolated Singularities)

A point  $a$  is said to be an isolated singularity of function  $f(z)$  if  $f(z)$  is analytic at each point in some neighborhood  $|z - a| < \delta$  of  $a$ , except at the point  $a$  itself. Otherwise, it is called non-isolated.

**Examples:**

- 1 The function  $f(z) = \frac{z+1}{z(z^2+2)}$  possesses three isolated singular points  $z = 0$ ,  $z = \sqrt{2}i$  and  $z = -\sqrt{2}i$ .
- 2 The function  $\ln z$  has a singularity at the origin, but it is not isolated since every neighborhood of zero contains points on the negative real axis where  $\ln z$  ceases to be analytic.

# Definitions

Suppose  $f$  has an isolated singularity at  $z = a$ .

## Definition (Removable Singularities)

If there a function  $g$ , analytic at  $a$  and such that  $f(z) = g(z)$  for all  $x$  in some deleted neighborhood of  $a$ , we say that  $f$  has a removable singularity at  $a$  i.e., if the value of  $f$  is connected at the point  $z = a$ , it becomes analytic there.

## Definition (Poles)

If, for  $z = a$ ,  $f(z)$  can be written as  $f(z) = \frac{\phi(z)}{\psi(z)}$  where  $\phi$  and  $\psi$  are analytic at  $a$ ,  $\phi(a) \neq 0$ , and  $\psi(a) = 0$ , we say that  $f$  has a pole at  $a$ . In other words, if  $\psi$  has a zero of order  $m$  at  $a$ , we say that  $f$  has a pole of order  $m$ .

## Definition (Essential Singularities)

If  $f$  has neither a removable singularity nor a pole at  $a$ , we say that  $f$  has an essential singularity at  $a$ .

# Removable Singularities

Let  $z = a$  be an isolated singularity of a function  $f(z)$ . Since the singularity is isolated, there exists a deleted neighborhood  $N_a$  defined by

$$0 < |z - a| < \delta$$

in which  $f(z)$  is analytic. Then, by Laurent's theorem, we can expand  $f(z)$  in a series of non-negative and negative powers of  $(z - a)$  in  $N_a$ . Thus, with suitable definitions of  $a_n$  and  $b_n$  in the region  $N_a$ , we have

$$f(z) = \sum_{n=0}^{\infty} a_n(z - a)^n + \sum_{n=1}^{\infty} b_n(z - a)^{-n}.$$

The part  $b_n(z - a)^{-n}$  of Laurent's series is called the **principal part** of  $f(z)$  at  $z = a$ .



# Removable Singularities

Let  $z = a$  be an isolated singularity of a function  $f(z)$ . Since the singularity is isolated, there exists a deleted neighborhood  $N_a$  defined by

$$0 < |z - a| < \delta$$

in which  $f(z)$  is analytic. Then, by Laurent's theorem, we can expand  $f(z)$  in a series of non-negative and negative powers of  $(z - a)$  in  $N_a$ . Thus, with suitable definitions of  $a_n$  and  $b_n$  in the region  $N_a$ , we have

$$f(z) = \sum_{n=0}^{\infty} a_n(z - a)^n + \sum_{n=1}^{\infty} b_n(z - a)^{-n}.$$

The part  $b_n(z - a)^{-n}$  of Laurent's series is called the **principal part** of  $f(z)$  at  $z = a$ .

Now, there arise three distinct possibilities:

- 1 **Removable Singularity.** If the principal part of  $f(z)$  at  $z = a$  consists of no terms, then  $a$  is said to be a removable singularity of  $f(z)$ .

**Alternative Definition.** A singularity  $z = a$  is called a removable singularity of  $f(z)$  if  $\lim_{z \rightarrow a} f(z)$  exists finitely.

# Removable Singularities

Let  $z = a$  be an isolated singularity of a function  $f(z)$ . Since the singularity is isolated, there exists a deleted neighborhood  $N_a$  defined by

$$0 < |z - a| < \delta$$

in which  $f(z)$  is analytic. Then, by Laurent's theorem, we can expand  $f(z)$  in a series of non-negative and negative powers of  $(z - a)$  in  $N_a$ . Thus, with suitable definitions of  $a_n$  and  $b_n$  in the region  $N_a$ , we have

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n + \sum_{n=1}^{\infty} b_n(z-a)^{-n}.$$

The part  $b_n(z-a)^{-n}$  of Laurent's series is called the **principal part** of  $f(z)$  at  $z = a$ .

Now, there arise three distinct possibilities:

- 1 **Removable Singularity.** If the principal part of  $f(z)$  at  $z = a$  consists of no terms, then  $a$  is said to be a removable singularity of  $f(z)$ .

**Alternative Definition.** A singularity  $z = a$  is called a removable singularity of  $f(z)$  if  $\lim_{z \rightarrow a} f(z)$  exists finitely.

For example, the function  $f(z) = \frac{\sin z}{z}$  has a removable singularity at  $z = 0$  since

$$\frac{\sin z}{z} = \frac{1}{z} \left( z - \frac{z^3}{3!} + \frac{z^5}{5!} - \dots \right) = 1 - \frac{z^2}{3!} + \frac{z^4}{5!} - \dots$$

has no term containing negative powers of  $z$ . However, this singularity can be removed and the function be made analytic by defining  $\frac{\sin z}{z} = 1$  at  $z = 0$ .

- 2 **Pole.** If the principal part of a function  $f(z)$  at  $z = a$  consists of a finite number of terms, say  $m$ , we say that  $a$  is a pole of order  $m$  of  $f(z)$ . For example, if  $b_m$  is the last coefficient that does not vanish, then we have

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n + \frac{b_1}{z-a} + \frac{b_2}{(z-a)^2} + \cdots + \frac{b_m}{(z-a)^m}.$$

Poles of order 1 and 2 are called, respectively, single and double poles.

**Alternate Definition.** If there exists a positive integer  $m$  such that

$$\lim_{z \rightarrow a} (z-a)^m f(z) = b \neq 0, \text{ but } \lim_{z \rightarrow a} (z-a)^{m+1} f(z) = 0,$$

then  $z = a$  is called a pole of order  $m$ .

**Examples:**

- Let  $f(z) = \frac{1}{(z-1)^2(z-3)^5}$ . Then  $z = 1$  is a pole of order 2 and  $z = 3$  is a pole of order 5.
- $\csc^2 z$  has one double pole and an infinite number of simple poles.

# Isolated Essential Singularities

- 3 **Isolated Essential Singularity.** If the principal part of  $f(z)$  at  $z = a$  contains an infinite number of terms, then  $a$  is called an isolated essential singularity of  $f(z)$ . In such a case

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n + \sum_{n=1}^{\infty} \frac{b_n}{(z-a)^n},$$

the last series being convergent for all values of  $z$  in  $|z-a| < \delta$  except at  $z = a$ .

**Alternate Definition.** If there exists **no finite value of  $m$**  such that

$$\lim_{z \rightarrow a} (z-a)^m f(z) = b = \text{a finite non-zero constant},$$

then  $z = a$  is called an isolated essential singularity.

## Examples:

- $e^{1/z} = 1 + \frac{1}{z} + \frac{1}{2!z^2} + \frac{1}{3!z^3} + \cdots$  has an isolated essential singularity at  $z = 0$ .
- The function  $f(z) = (z-3) \sin \frac{1}{z+2}$  has Laurent's expansion  $f(z) = 1 - \frac{5}{z+2} - \frac{1}{6(z+2)^2} + \frac{5}{6(z+2)^3} + \frac{1}{120(z+2)^4} - \cdots$ . Thus  $z = -2$  is an essential singularity of  $f(z)$ .

# Isolated Essential Singularities

We must take utmost care while classifying a given point  $a$  as an isolated essential singularity of a function  $f(z)$  on the basis of Laurent's expansion of  $f(z)$  in which the series of negative powers of  $z - a$  does not terminate. It is important to bear in mind that the series should be convergent for all values of  $z$  in  $|z - a| < \delta$ , except at  $z = a$ , for some  $\delta > 0$ .

# Isolated Essential Singularities

We must take utmost care while classifying a given point  $a$  as an isolated essential singularity of a function  $f(z)$  on the basis of Laurent's expansion of  $f(z)$  in which the series of negative powers of  $z - a$  does not terminate. It is important to bear in mind that the series should be convergent for all values of  $z$  in  $|z - a| < \delta$ , except at  $z = a$ , for some  $\delta > 0$ .

For example, the following series contains an infinite number of terms in the principal part

$$\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} + \sum_{n=1}^{\infty} z^{-n}. \quad (4)$$

But on this ground alone, we should not declare that  $z = 0$  is an isolated essential singularity of the sum-function of the series (4). We must also test whether the series (4) converges in some deleted neighborhood of the origin, say,  $0 < |z| < \delta$ .

# Isolated Essential Singularities

We must take utmost care while classifying a given point  $a$  as an isolated essential singularity of a function  $f(z)$  on the basis of Laurent's expansion of  $f(z)$  in which the series of negative powers of  $z - a$  does not terminate. It is important to bear in mind that the series should be convergent for all values of  $z$  in  $|z - a| < \delta$ , except at  $z = a$ , for some  $\delta > 0$ .

For example, the following series contains an infinite number of terms in the principal part

$$\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} + \sum_{n=1}^{\infty} z^{-n}. \quad (4)$$

But on this ground alone, we should not declare that  $z = 0$  is an isolated essential singularity of the sum-function of the series (4). We must also test whether the series (4) converges in some deleted neighborhood of the origin, say,  $0 < |z| < \delta$ .

Since

$$\frac{1}{2-z} = \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \quad \frac{1}{z-1} = \sum_{n=1}^{\infty} z^{-n},$$

and the first series converges for  $|z| < 2$ , while the second series converges for  $|z| > 1$ . Thus, the domain of convergence of the series (4) is the annular region  $1 < |z| < 2$ , but it is not a neighborhood of the origin. Indeed, the sum-function of (4) in  $1 < |z| < 2$  is given by

$$f(z) = \frac{1}{z-1} + \frac{1}{2-z} = \frac{1}{3z-2-z^2}.$$

$f(z)$  is a function of which the only singularities are the simple poles at  $z = 1$  and  $z = 2$ .

# Isolated Essential Singularities

Let us consider another example of the series

$$f(z) = \sum_{n=1}^{\infty} (z-1)^{-n}, \quad (5)$$

which gives an impression, at first sight, that the point  $z = 1$  is an isolated essential singularity of the sum-function of the series (5). The crux of the matter is that the series converges for  $|z-1| > 1$  and this does not define a neighborhood of 1 .

Indeed, the sum-function of (5), in the domain of its convergence, is  $\frac{1}{z-2}$ , which is analytic at  $z = 1$  and of which the only singularity is the simple pole at  $z = 2$ .



# Classification of singularities via limits

Suppose  $z_0$  is an isolated singularity of  $f(z)$ . Then

$z_0$ is removable $\Leftrightarrow$	$\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0.$
$z_0$ is a pole $\Leftrightarrow$	(a) $\neg \left( \lim_{z \rightarrow z_0} (z - z_0)f(z) = 0 \right)$ and (b) $\exists n \in \mathbb{N}$ such that $\lim_{z \rightarrow z_0} (z - z_0)^{n+1}f(z) = 0.$ (The smallest such $n$ is called the order of the pole $z_0$ of $f$ .)
$z_0$ is essential $\Leftrightarrow$	$\forall n \in \mathbb{N} \neg \left( \lim_{z \rightarrow z_0} (z - z_0)^n f(z) = 0 \right).$

(Here  $\neg$  is the symbol for negation, to be read as "it is not the case that".)

# Classification via Laurent coefficients

Let

- 1  $z_0$  be an isolated singularity of  $f(z)$ , and
- 2  $f(z) = \sum_{n \in \mathbb{Z}} c_n (z - z_0)^n$  for  $0 < |z - z_0| < R$ , for some  $R > 0$ .

Then

$z_0$ is removable $\Leftrightarrow$	For all $n < 0$ , $c_n = 0$
$z_0$ is a pole $\Leftrightarrow$	There exists an $m \in \mathbb{N}$ such that (a) $c_{-m} \neq 0$ and (b) for all $n < -m$ , $c_n = 0$ Then the order of the pole $z_0$ is $m$ .
$z_0$ is essential $\Leftrightarrow$	There are infinitely many negative indices $n$ such that $c_n \neq 0$ .

1 Zeros of Analytic Functions

2 Singular Points

3 Residue Theorem

# Introduction

The inspiration behind this section is the desire to obtain possible values for the integrals  $\int_C f(z)dz$ , where  $f$  is analytic inside the closed curve  $C$  and on  $C$ , **except for  $a$  inside  $C$** .

- If  $f$  has a removable singularity at  $a$ , then it is clear that the integral will be zero.
- If  $z = a$  is a pole or an essential singularity, then the answer is not always zero, but can be found with little difficulty.

In this section, we show the very surprising fact that Cauchy's residue theorem yields a very elegant and simple method for evaluation of such integrals.

# The Residues at Singularities

We know that, in the neighborhood of an isolated essential singularity  $z = a$ , a single-valued analytic function  $f(z)$  can be expanded in Laurent's series

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n + \sum_{n=1}^{\infty} b_n(z-a)^{-n}.$$

Thus the coefficient  $b_1$ , which is called the residue of  $f(z)$  at  $z = a$ , is given by the formula

$$b_1 = \frac{1}{2\pi i} \int_{\gamma} f(z) dz,$$

where  $\gamma$  is any circle with center at  $a$ , which includes singularities of  $f(z)$ . We denote the residue of  $f(z)$  at  $z = a$  by

$$\operatorname{Res}_{z=a} f(z) \quad \text{or} \quad \operatorname{Res}(f, z_0).$$

If  $z = a$  is a single pole, then we also have

$$b_1 = \lim_{z \rightarrow \alpha} (z-a)f(z).$$

# The Residues at Singularities

A more general definition of the "residue" of a function  $f(z)$  at a point is the following.

If  $z = a$  is **the only singularity** of an analytic function  $f(z)$  inside a closed contour  $C$  and

$$\frac{1}{2\pi i} \int_C f(z) dz$$


has a definite value, then the value is called the residue of  $f(z)$  at  $z = a$ .

If  $C$  includes **a finite number of singularities** of  $f(z)$  which is an analytic function elsewhere inside and on  $C$ , then the **sum of the residues** at singularities is given by

$$\frac{1}{2\pi i} \int_C f(z) dz.$$

# The Residues at Singularities

If  $f(z)$  is analytic in a multiply connected region bounded by and including the contours  $C$  and  $C_1, C_2, \dots, C_n$  contained within  $C$  as shown in the following figure.



then the sum of the residues of  $f(z)$  at the included essential singularities is easily seen to be given by

$$\frac{1}{2\pi i} \left[ \int_C f(z) dz - \sum_{r=1}^n \int_{C_r} f(z) dz \right].$$

# Calculation of Residues in Some Special Cases

## 1 Residues at Simple Poles:

When  $f(z) = \frac{\phi(z)}{\psi(z)}$ , where  $\phi(a) \neq 0$  and  $\psi(z)$  has a simple zero at  $z = a$ .

Since  $\psi(z)$  has a simple zero at  $z = a$ ,  $\psi(a) = 0$ , but  $\psi'(a) \neq 0$ . Then it is evident that  $f(z)$  has a simple pole at  $z = a$ . Therefore, we have

$$\begin{aligned}\operatorname{Res}_{z=a} f(z) &= \operatorname{Res}_{z=a} \frac{\phi(z)}{\psi(z)} = \lim_{z \rightarrow a} (z - a) \frac{\phi(z)}{\psi(z)} \\ &= \lim_{z \rightarrow a} \frac{\phi(z)}{\frac{\psi(z) - \psi(a)}{z - a}} = \frac{\phi(a)}{\psi'(a)}.\end{aligned}$$



# Calculation of Residues in Some Special Cases

## 1 Residues at Simple Poles:

When  $f(z) = \frac{\phi(z)}{\psi(z)}$ , where  $\phi(a) \neq 0$  and  $\psi(z)$  has a simple zero at  $z = a$ .

Since  $\psi(z)$  has a simple zero at  $z = a$ ,  $\psi(a) = 0$ , but  $\psi'(a) \neq 0$ . Then it is evident that  $f(z)$  has a simple pole at  $z = a$ . Therefore, we have

$$\begin{aligned}\operatorname{Res}_{z=a} f(z) &= \operatorname{Res}_{z=a} \frac{\phi(z)}{\psi(z)} = \lim_{z \rightarrow a} (z-a) \frac{\phi(z)}{\psi(z)} \\ &= \lim_{z \rightarrow a} \frac{\phi(z)}{\frac{\psi(z)-\psi(a)}{z-a}} = \frac{\phi(a)}{\psi'(a)}.\end{aligned}$$

## 2 Residues of Poles of Order Greater than Unity:

When  $f(z)$  has a pole of order  $m(m > 1)$  at  $z = a$ .

Laurent's expansion of  $f(z)$  in the neighborhood of the point  $z = a$  is given by

$$f(z) = \frac{b_m}{(z-a)^m} + \dots + \frac{b_1}{z-a} + a_0 + a_1(z-a) + \dots.$$

Hence we have

$$\begin{aligned}(z-a)^m f(z) &= b_m + b_{m-1}(z-a) + \dots + b_1(z-a)^{m-1} \\ &\quad + a_0(z-a)^m + \dots\end{aligned}$$

Differentiating both sides with respect to  $z$ ,  $(m-1)$  times, we have

$$D^{m-1}[(z-a)^m f(z)] = (m-1)!b_1 + m(m-1) \dots 2a_0(z-a) + \dots$$

# Calculation of Residues in Some Special Cases

Taking the limit as  $z \rightarrow a$ , we have

$$\begin{aligned}(m-1)!b_1 &= \lim_{z \rightarrow a} \left[ D^{m-1} \{ (z-a)^m f(z) \} \right] \\ &= \lim_{z \rightarrow a} \left[ \phi^{(m-1)}(z) \right] \\ &= \phi^{(m-1)}(a),\end{aligned}$$

where  $f(z) = \frac{\phi(z)}{(z-a)^m}$ . Hence we have

$$\operatorname{Res}_{z=a} f(z) = b_1 = \frac{\phi^{(m-1)}(a)}{(m-1)!}$$

In particular, if  $\frac{\phi(z)}{(z-a)^2}$ , then we have

$$\operatorname{Res}_{z=a} f(z) = \phi'(a).$$

If  $\frac{\phi(z)}{(z-a)^3}$ , then we have

$$\operatorname{Res}_{z=a} f(z) = \frac{\phi''(a)}{2!}$$

and so on.

# Calculation of Residues in Some Special Cases

## 3 Another Method:

Since residue at  $z = a$  is the coefficient of  $\frac{1}{z-a}$  in Laurent's expansion of  $f(z)$ , it follows that the residue is the coefficient of  $1/t$  in the expansion of  $f(a+t)$  as a power series where  $t$  is considered sufficiently small.

When  $f(z) = \frac{\phi(z)}{z\psi(z)}$ , where the numerator and the denominator have no common factor while  $\psi(0) \neq 0$ .

In this case,  $f(z)$  has a simple pole at the origin, due to the factor  $\frac{1}{z}$ , and  $f(z)$  also has a number of simple poles arising from the zeros of  $\psi(z)$ . Hence we have

$$\operatorname{Res}_{z=0} f(z) = \frac{\phi(0)}{\psi(0)}.$$

Suppose that  $z = a_m$  is a simple pole of  $\frac{1}{\psi(z)}$ . Then we have

$$\begin{aligned}\operatorname{Res}_{z=a_m} f(z) &= \lim_{z \rightarrow a_m} \left[ (z - a_m) \frac{\phi(z)}{z\psi(z)} \right] \\ &= \frac{\phi(a_m)}{a_m \psi'(a_m)}\end{aligned}$$

provided  $a_m \neq 0$ .

# Definition: Residues at Infinity

The definition of residue can be extended to include the point at infinity. If  $f(z)$  is analytic or has an **isolated essential singularity at infinity** and  $C$  is a circle enclosing within it all other singularities of  $f(z)$  in the finite regions of the  $z$ -plane, then the residue at infinity is defined by

$$\frac{1}{2\pi i} \int_C f(z) dz$$

where the integral is taken round  $C$  in the negative sense (clockwise direction), provided that this integral has a definite value.

If we take the integral round  $C$  in an anti-clockwise direction, then the residue at infinity is  $-\frac{1}{2\pi i} \int_C f(z) dz$ .

# Calculation: Residues at Infinity

By means of the substitution  $z = \frac{1}{Z}$ , the integral defining the residue at infinity takes the form

$$\frac{1}{2\pi i} \int \left\{ -f\left(\frac{1}{Z}\right) \right\} \frac{dZ}{Z^2}$$

taken in a counterclockwise direction round a sufficiently small circle with center at the origin. It follows that, if

$$\lim_{Z \rightarrow 0} \left\{ -f\left(\frac{1}{Z}\right) Z^{-1} \right\} \quad \text{or} \quad \lim_{z \rightarrow \infty} \{ -zf(z) \}$$

has a definite value, then that value is the residue of  $f(z)$  at infinity.

# Some Residue Theorems

## Theorem (Cauchy's Residue Theorem)

If  $f(z)$  is regular, except at a finite number of poles  $z_0, z_2, \dots, z_n$  within a closed contour  $C$  where its residues are  $R_1, R_2, \dots, R_n$ , respectively, and continuous on the boundary  $C$ , then

$$\int_C f(z) dz = 2\pi i (R_1 + R_2 + \dots + R_n)$$

or

$$\int_C f(z) dz = 2\pi i \text{ (sum of residues at the poles within) } C.$$

# Some Residue Theorems

## Theorem (Cauchy's Residue Theorem)

If  $f(z)$  is regular, except at a finite number of poles  $z_0, z_2, \dots, z_n$  within a closed contour  $C$  where its residues are  $R_1, R_2, \dots, R_n$ , respectively, and continuous on the boundary  $C$ , then

$$\int_C f(z) dz = 2\pi i (R_1 + R_2 + \dots + R_n)$$

or

$$\int_C f(z) dz = 2\pi i \text{ (sum of residues at the poles within) } C.$$

## Theorem

If a single-valued function has only a finite number of singularities, then the sum of residues at these singularities, including the residue at infinity, is zero.