

# Soil organic carbon and nitrogen in a Mollisol in central Ohio as affected by tillage and land use

P. Puget, R. Lal\*

*The Ohio State University, School of Natural Resources, Kottman Hall, 2021 Coffey Road, Columbus, OH 43212, USA*

Received 23 December 2002; received in revised form 19 March 2004; accepted 25 March 2004

## Abstract

Minimum tillage practices are known for increasing soil organic carbon (SOC). However, not all environmental situations may manifest this potential change. The SOC and N stocks were assessed on a Mollisol in central Ohio in an 8-year-old tillage experiment as well as under two relatively undisturbed land uses; a secondary forest and a pasture on the same soil type. Cropped systems had  $51 \pm 4$  (equiv. mass)  $\text{Mg ha}^{-1}$  lower SOC and lower  $3.5 \pm 0.3$  (equiv. mass)  $\text{Mg ha}^{-1}$  N in the top 30 cm soil layer than under forest. Being a secondary forest, the loss in SOC and N stocks by cultivation may have been even more than these reported herein. No differences among systems were detected below this depth. The SOC stock in the pasture treatment was  $29 \pm 3 \text{ Mg ha}^{-1}$  greater in the top 10 cm layer than in cultivated soils, but was similar to those under forest and no-till (NT). Among tillage practices (plow, chisel and NT) only the 0–5 cm soil layer under NT exhibited higher SOC and N concentrations. An analysis of the literature of NT effect on SOC stocks, using meta-analysis, suggested that NT would have an overall positive effect on SOC sequestration rate but with a greater variability of what was previously reported. The average sequestration rate of NT was  $330 \text{ kg SOC ha}^{-1} \text{ year}^{-1}$  with a 95% confidence interval ranging from 47 to  $620 \text{ kg SOC ha}^{-1} \text{ year}^{-1}$ . There was no effect of soil texture or crop rotation on the SOC sequestration rate that could explain this variability. The conversion factor for SOC stock changes from plow to NT was equal to 1.04. This suggests that the complex mechanisms and pathways of SOC accrual warrant a cautious approach when generalizing the beneficial changes of NT on SOC stocks.

© 2004 Elsevier B.V. All rights reserved.

**Keywords:** Soil carbon sequestration; Nitrogen stock; No till; Pasture; Forest; Meta-analysis; Soil quality; Conservation tillage

## 1. Introduction

Conversion of natural ecosystems to agriculture as well as increasing intensity of tillage are known to decrease soil organic matter (SOM) levels and contribute significantly to the increase in atmospheric  $\text{CO}_2$  concentration (Lal et al., 1998). Between 1850 and 1995, SOM mineralization emitted  $136 \pm 55 \times 10^{15} \text{ g of C}$

to the atmosphere (Houghton, 1995; Watson et al., 2001). Changes in SOM by different agricultural land uses and practices have been extensively reviewed by Mann (1986), Davidson and Ackerman (1993), Guo and Gifford (2002), and Murty et al. (2002). Guo and Gifford (2002) reported that soils lost 42 and 59% of their soil organic carbon (SOC) stock upon conversion from forest to crop and from grassland to crop, respectively. In contrast, Mann (1986) calculated losses <20% of the initial stock, which was corroborated by Kern and Johnson (1993) and Murty et al. (2002) who evaluated the decrease of SOC stock in the major

\* Corresponding author. Tel.: +1-614-292-9069;

fax: +1-614-292-7932.

E-mail address: [lal.1@osu.edu](mailto:lal.1@osu.edu) (R. Lal).

US cropland soils at ca. 16 and 22–25%, respectively. However, Davidson and Ackerman (1993) argued that calculations made on a fixed soil depth instead of an equivalent mass of soil underestimated the changes in SOC stock. With the objective of reducing net emission of CO<sub>2</sub> and other greenhouse gases (GHGs), soil C sink capacity has been debated (Swift, 2001; West and Marland, 2002), i.e., can the “lost C” be re-sequestered in soils and if so, how and to what extent?

Three main strategies for SOC sequestration in the biosphere are: (i) enhancing the re-growth of perennial vegetation through conversion of cropland to grassland or forestland, (ii) increasing the net primary productivity and therefore the amount of residue returned to the soil, mainly through intensification of agricultural inputs: fertilizer, irrigation, manuring, and (iii) adopting conservation tillage especially no tillage (NT). Post and Kwon (2000) reported that changes in land use to perennial vegetation may potentially sequester C at a rate of 300–350 kg C ha<sup>-1</sup> year<sup>-1</sup>. However, they acknowledged that this increase was too small to counteract the “anthropogenically released” C in the global C cycle. Several studies have shown that intensification of farming systems augments SOC levels mainly by increasing the amount of biomass produced and the amount of residues returned (Larson et al., 1972; Zielke and Christenson, 1986; Havlin et al., 1990; Campbell and Zetner, 1993; Follett, 2001). However, some researchers have pointed out that these approaches may not be as efficient as perceived when fluxes of all GHGs, especially the hidden C costs of agricultural intensification, are taken into account (Robertson et al., 2000; Schlesinger, 2000a,b).

Conservation tillage practices, especially NT, can improve SOC and other soil nutrient stocks. Conservation tillage was developed primarily to combat soil erosion by developing a protective mulch layer of crop residues at the soil surface. Consequently, the repetitive use of this technique profoundly affects the incorporation and depth distribution of SOC, as well as the whole ecology and functioning of soil (Kladivko, 2001). No tillage impacts SOC stock in two ways: (i) by reducing disturbance which favors the formation of soil aggregates and protects SOC encapsulated inside these stable aggregates from rapid oxidation (Chaney et al., 1985; Elliott, 1986; Beare et al., 1994; Six et al., 2000) and (ii) by modifying the local edaphic envi-

ronment: bulk density, pore size distribution, temperature, water and air regime that might also restrict SOC biodegradation (Mielke et al., 1986; Kay and VandenBygaart, 2002). The literature is replete with studies that show an increase in SOC stock with conversion to NT, at least in the surface soil (e.g. Blevins et al., 1983; Dick, 1983; Dick et al., 1986, 1991; Doran, 1980; Lal et al., 1990). Paustian et al. (1997) and Lal et al. (1998) summarized the rate of accumulation of SOC stock under NT at 300–800 kg SOC ha<sup>-1</sup> year<sup>-1</sup>. However, under some conditions, especially in fine-textured soils, there may be little or no increase in SOC stock, especially at a depth greater than the plow layer (Mielke et al., 1986; Dalal, 1988; Angers et al., 1995, 1997; Wander et al., 1998).

The present study was undertaken to determine how contrasting agricultural land uses and tillage practices might affect SOC sequestration in a fine-textured Mollisol in central Ohio. We determined the distribution and stocks of SOC and N through the soil profile and compared our data with those reported in the literature in which studies were carried out on paired experiments of NT versus plow till.

## 2. Materials and methods

A tillage experiment was initiated in Autumn 1993 to determine the effect of three tillage practices [chisel till (CT), moldboard plow till (PT) and NT] on corn (*Zea mays*) yield and plant nutrients. The experiment was located at the Don Scott Experimental Farm, Columbus, OH (40°04'30"N; 83°04'W) on a Kokomo silty clay loam (somewhat poorly drained, fine, mixed, mesic, Typic Argiaquolls) (McLoda and Parkinson, 1980). The experimental plot was divided in two blocks (west and east) on which corn and soybean (*Glycine max*) were cropped in alternate years. The above-ground crop residue returned was 8–10 Mg ha<sup>-1</sup> year<sup>-1</sup> during the corn cycle and 2–3 Mg ha<sup>-1</sup> during the soybean cycle. Each block was divided into 12 subplots of 17.5 m × 22.5 m corresponding to three tillage treatments replicated four times. Soil samples were collected in December 2001 in the west block after corn harvest following 8 years of consecutive tillage treatment. A soil core (7.6 cm in diameter) was taken from the center of each subplot, using a truck-mounted hydraulic probe

to 70 or 80 cm depth. The core was divided into 5 cm increments to 40 cm depth and 10 cm increments thereafter. Two adjacent areas, located within 1 km of the tillage experiment on the same soil type, were also sampled: one managed as grazed pasture for at least 15 years and the second as a woodlot (ca. 80-year-old secondary forest of mixed hardwood). Predominant species included a mixture of orchard grass (*Dactylis glomerata*) and red clover (*Trifolium pratense*) in pasture, and maple (*Acer* spp.), oak (*Quercus* spp.) and hickory (*Carya glabra*) in the woodlot. Three distinct soil cores were taken from these plots as pseudo field replicates.

The coarse crop residue corresponding to 2001 harvest was not included in the soil samples. Similarly, the fresh litter in the forest and the green grass in the pasture plot were also removed before taking soil samples. Soil samples were brought to the lab, weighed (field moist weight), dried at 60 °C and weighed again. Sub-samples were taken and weighed after 48 h at 105 °C for computing the bulk density. Total C and N concentrations (dry combustion method on a Thermoquest NC soil 1100) were determined after sieving the samples through a 2-mm sieve and fine grinding using a ball-mill grinder. No residues were removed. No carbonates were detected when an acid-drop test was used on the samples. Thus total C was assumed to be equivalent to SOC. The C and N stocks were then calculated by multiplying the C or N concentration by the bulk density and expressed in  $\text{Mg ha}^{-1}$ .

### 2.1. Data analysis

Analysis of variance (ANOVA) was used to assess the treatment effects (land use and tillage) and depth, or treatment only, on soil bulk density, SOC and N concentrations and stocks. Tukey's mean difference test was used at the 5% probability level to separate treatments (SAS Institute, 1999).

Comparison of SOC stock between NT and PT was made for 28 published studies reporting 56-paired experiments. We selected studies reporting SOC stock on an area basis or including data of both bulk density and SOC concentration. A linear regression analysis was performed between SOC stock under PT and NT by using the linear model procedure in SAS with a fixed intercept equal to zero.

A meta-analysis of literature data was conducted to provide a comprehensive evaluation of the effect of NT on SOC. Meta-analysis allows synthesis of quantitatively independent studies to answer a specific question in different environments (Gurevitch and Hedges, 1999, 2001). This tool has already been used in the field of soil science to address questions related to SOC stock changes under different land uses and management (Johnson and Curtis, 2001; Guo and Gifford, 2002).

Because of the diverse situations (e.g., duration of the experiment, cropping rotation, soil type, initial SOC stock, location, etc.) the rate of SOC sequestration (difference in SOC stock between NT and PT divided by the duration of the experiment in years) was chosen as the response variable. Most of the studies did not report consistent measures of variance, thus an 'unweighted' meta-analysis (i.e., equal variance) was conducted. The mean and 95% confidence interval were calculated using Meta-Win 2 software (Rosenberg et al., 2000) after a bias corrected bootstrapping procedure (Dixon, 2001; Manly, 1997). The dataset was partitioned into categories corresponding to soil type and cropping intensity. Two soil type categories were fine-textured soils (clay content >30%) and coarse-textured soils. Two cropping intensity categories were based on the amount of crop residue biomass returned: (1) low; wheat (*Triticum aestivum*)–fallow rotation, continuous soybean (*Glycine max*) or corn silage, and (2) high; continuous corn or sorghum (*Sorghum bicolor*).

## 3. Results and discussion

### 3.1. Soil bulk density

Tillage treatment had no impact on soil bulk density. Lower bulk density was observed in forest compared to cultivated plots and the pasture to 30 cm depth (Fig. 1). Unger (1982) reported that bulk density of a Pullman clay loam in Texas was not impacted by 36 years of different tillage treatments (mainly disk tillage). In contrast, other physical parameters of soil quality (dry and water-stable aggregate size distribution, resistance to rupture and penetration) were modified. Similarly, Lal (1999) also reported that tillage did not have any effect




Fig. 1. Soil profile bulk density under plow till, chisel till, no till, pasture and forest. Error bars represent standard error of  $n = 4$  for tilled soils and  $n = 3$  for pasture and forest soils.




Fig. 2. Soil profile organic carbon concentration under plow till, chisel till, no till, pasture and forest. Error bars represent standard error of  $n = 4$  for tilled soils and  $n = 3$  for pasture and forest soils.

on soil bulk density of a Mollic Ochraqualf in Ohio.

### 3.2. Soil organic carbon and nitrogen concentrations

Distribution of C and N concentrations in the soil profile are shown in Figs. 2 and 3. Both SOC and N distributions were strongly stratified with depth in the forest soil, which had the highest C and N concentrations in the top 30 cm. Differences among land uses and tillage practices diminished below 30 cm and no differences were observed below 50 cm. In comparison to cropland, pasture soil contained greater C and

N but only in the 0–5 and 5–10 cm soil layers. Among the three tillage treatments, SOC and N concentrations were significantly different, but only in the 0–5 cm soil layer (Table 1). Between CT and PT, neither SOC nor N differed at any depth.

Similar results showing different SOC and N concentrations among tillage systems in the top 10 cm of clayey soils have been reported. For instance, Dick (1983) and Lal et al. (1990) reported for a Mollic Ochraqualf in northwestern Ohio that even after 18 and 12 years of continuous cultivation with NT, an increase in SOC concentration by 15% compared with PT occurred only in the first 5 cm layer. Higher but




Fig. 3. Soil profile nitrogen concentration under plow till, chisel till, no till, pasture and forest. Error bars represent standard error of  $n = 4$  for tilled soils and  $n = 3$  for pasture and forest soils.




Fig. 4. Soil profile C:N ratio under plow till, chisel till, no till, pasture and forest. Error bars represent standard error of  $n = 4$  for tilled soils and  $n = 3$  for pasture and forest soils.

non-significant concentrations were observed in NT versus PT between 5 and 20 cm depths and no differences between the two treatments below 20 cm depth. Wander et al. (1998) and Needelman et al. (1999) found similar results in a large number of clayey soils in Illinois where the top 0–5 cm layer under NT contained greater C and N concentrations than under PT. Similar results have been reported in fine-textured soils of different pedoclimatic conditions, e.g., for a Grundy silty clay soil in Kansas (Havlin et al., 1990); the cool and humid environment in Canada (Carter and

Rennie, 1982; Angers et al., 1995, 1997) and in Vertisols in Australia under tropical and subtropical climate (Dalal, 1988; Chan et al., 2002; Haynes and Beare, 1996).

The C:N ratio was affected by both land use and tillage (Fig. 4). Forest soil had the highest C:N ratio for all sampling depths, and pasture the lowest values (except for 0–5 cm depth). This effect was similar to that reported by Franzluebbers et al. (2000) and values mainly reflected different qualities of organic residues entering the SOM pool and could be attributed to

Table 1  
Cumulative stock of SOC and N in plow till, chisel till, no till, pasture and forest

| Treatment<br>Depth (cm) | Soil organic carbon stock (Mg ha <sup>-1</sup> ) |             |         |         |        | Stock of N (Mg ha <sup>-1</sup> ) |             |         |         |         |
|-------------------------|--|-------------|---------|---------|--------|-----------------------------------|-------------|---------|---------|---------|
|                         | Plow till  | Chisel till | No till | Pasture | Forest | Plow till                         | Chisel till | No till | Pasture | Forest  |
| 0–5                     | 16 c   | 16 c        | 19 bc   | 44 a    | 27 b   | 1.4 b                             | 1.4 b       | 1.6 b   | 3.7 a   | 2.1 b   |
| 0–10                    | 32 b   | 30 b        | 35 b    | 61 a    | 54 a   | 2.8 b                             | 2.7 b       | 3.1 b   | 5.3 a   | 4.3 a   |
| 0–15                    | 47 b   | 44 b        | 50 b    | 75 a    | 76 a   | 4.2 b                             | 4.1 b       | 4.5 b   | 6.7 a   | 6.1 a   |
| 0–20                    | 62 c   | 59 c        | 65 bc   | 88 ab   | 97 a   | 5.5 b                             | 5.4 b       | 5.9 b   | 7.9 a   | 7.9 a   |
| 0–25                    | 76 b   | 73 b        | 81 b    | 99 ab   | 114 a  | 6.8 b                             | 6.8 b       | 7.4 ab  | 9.0 a   | 9.3 a   |
| 0–30                    | 88 b   | 84 b        | 94 b    | 107 b   | 127 a  | 7.9 NS                            | 7.8 NS      | 8.6 NS  | 9.7 NS  | 10.4 NS |
| 0–35                    | 98 b   | 91 b        | 103 b   | 112 b   | 139 a  | 8.8 NS                            | 8.5 NS      | 9.5 NS  | 10.3 NS | 11.2 NS |
| 0–40                    | 105 b  | 98 b        | 110 b   | 119 b   | 147 a  | 9.5 NS                            | 9.1 NS      | 10.3 NS | 11.0 NS | 12.0 NS |
| 0–50                    | 112 b  | 108 b       | 120 b   | 127 b   | 161 a  | 10.3 NS                           | 10.3 NS     | 11.3 NS | 12.1 NS | 13.3 NS |
| 0–60                    | 120 b  | 117 b       | 128 b   | 136 b   | 173 a  | 11.2 NS                           | 11.4 NS     | 12.3 NS | 13.1 NS | 14.7 NS |
| 0–70                    | 137 b  | 126 b       | 135 b   | 147 b   | 178 a  | 13.1 NS                           | 12.6 NS     | 13.1 NS | 14.3 NS | 15.2 NS |
| 0–80                    | 139 NS   | 134 NS      | 143 NS  | 149 NS  |        | 13.3 NS                           | 13.5 NS     | 14.1 NS | 14.6 NS |         |

Different letters in the same row represent a significant difference in stock among treatments, according to Tukey's mean significance test at  $P < 0.05$ . NS is not significant.

contrasting vegetation covers. Tillage treatment also altered soil C:N ratio with lowest associated with minimum tillage practices. The lower C:N ratio in CT and NT than in PT may have been due to a relatively higher contribution of root than aboveground inputs. The high potential N mineralization and microbial biomass under minimum tillage, as reported by other researchers (Doran, 1980; Franzluebbers and Arshad, 1997), may have also led to this change in the C:N ratio.

### 3.3. Soil organic carbon and nitrogen stocks

There were no differences in stocks of SOC and N among the three tillage systems throughout the soil profile (Table 1). In the whole soil profile (0–80 cm depth) this Mollisol stored an average of 139 Mg C ha<sup>-1</sup> and 13.6 Mg N ha<sup>-1</sup> when cultivated.

There were differences in SOC and N stocks between cultivated and uncultivated soils (Table 1). Pasture soil contained the highest SOC and N stocks in the top 5 cm layer reflecting the larger grass root density found in this layer. Pasture soil had 44 Mg C ha<sup>-1</sup> and 3.7 Mg N ha<sup>-1</sup>, which was more than twice the SOC and N stocks in cultivated soils and more than 1.5 times that in the forest soil. The difference in SOC and N stocks between pasture and cultivated soils decreased with increasing sampling depth. The contribution of the deeper soil layers to the SOC and N stocks was small and similar to that in cropped soils below 10 cm depth. In forest, the soil layers between 10 and 40 cm depth contributed almost equally to the SOC and N stocks as the 0–5 cm soil layer. Thus, in the whole soil profile (0–80 cm depth or 70 cm under forest soil) the soil under forest contained significantly greater SOC stock than that under pasture and the cultivated soils: 178 Mg C ha<sup>-1</sup> in forest versus 136 ± 8 Mg C ha<sup>-1</sup> in the other treatments. When SOC and N stocks were computed on an equivalent soil mass rather than on a fixed depth basis, similar differences among treatments occurred (Table 2). In 0–30 cm soil depth cropped soils had an average of 63% of the SOC stock as that under forest. Below 30 cm, no further N or C losses were observed in comparison to the reference stock under forest, suggesting that changes in SOC and N stocks upon cultivation occurred mainly in the A horizon. Being a secondary forest, the loss in SOC and N stocks by

Table 2  
Stock of SOC on an equivalent soil mass referenced to plow till at 0–20 cm and at 0–30 cm depth

|             | Equivalent soil layer depth of 0–20 cm in plow till |                           |                                  |                                | Equivalent soil layer depth of 0–30 cm in plow till |                                  |                                  |                                |
|-------------|---|---------------------------|----------------------------------|--------------------------------|---|----------------------------------|----------------------------------|--------------------------------|
|             | Soil mass (Mg ha <sup>-1</sup> )                    | Equivalent thickness (cm) | SOC stock (Mg ha <sup>-1</sup> ) | N stock (Mg ha <sup>-1</sup> ) | Percentage of forest SOC stock (%)                  | Percentage of forest N stock (%) | SOC stock (Mg ha <sup>-1</sup> ) | N stock (Mg ha <sup>-1</sup> ) |
| Plow till   | 2688  | 20                        | 62 (±3) b                        | 6 (±0.2) b                     | 53  | 57                               | 88.5 (±2.6) b                    | 7.9 (±0.2) b                   |
| Chisel till | 2688  | 20                        | 59 (±2) b                        | 5 (±0.1) b                     | 50  | 56                               | 83.3 (±2.5) b                    | 7.7 (±0.2) b                   |
| No till     | 2688  | 19.5                      | 64 (±1) b                        | 6 (±0.1) b                     | 54  | 60                               | 90.9 (±2.7) b                    | 8.3 (±0.2) b                   |
| Pasture     | 2688  | 17.9                      | 83 (±12) ab                      | 7 (±0.9) a                     | 71  | 78                               | 108.4 (±13.0) b                  | 9.9 (±1.1) b                   |
| Forest      | 2688  | 26.4                      | 117 (±10) a                      | 10 (±0.8) a                    | 100   | 100                              | 138.3 (±9.4) a                   | 11.5 (±0.7) a                  |

Different letters in the same columns express a significant difference among treatment at  $P < 0.05$  according to Tukey's mean difference significance test. Values in parenthesis represent standard error of  $n = 4$  for tilled soils and  $n = 3$  for pasture and forest soil.



cultivation may have been greater than that presented in these comparisons. There was some improvement in SOC in pasture as reported previously (Conant et al., 2001), but the beneficial effect of pasture was limited to the surface 15–20 cm of soil (Table 1). Pasture soil had  $29 \pm 3 \text{ Mg ha}^{-1}$  greater SOC in the top 10 cm layer than the cultivated soils. Furthermore, the non-particulate fraction of C was greater under pasture than NT (Franzluebbers and Stuedemann, 2002).

Within the surface 20 cm, results of C and N were very consistent. Below 20 cm, minor differences in N occurred, but treatment effects were similar to SOC. Similar results were reported by Murty et al. (2002).

Change in SOC stock from cropping compared with forest land was less than that reported by Mikhailova et al. (2000) in a Russian Chernozem after 50 years of continuous cultivation and hay production. They observed significant reductions to 100 cm depth with an average loss of 35% for SOC and 55% for N concentration. However, the lack of differences among tillage treatments in SOC and N stocks was similar to observations made previously such as on a Mollisol in the dryland US northern Great Plains (Halvorson et al., 2002), on soils in Alberta, Saskatchewan and eastern Canada (Carter and Rennie, 1982; Angers et al., 1995, 1997), on soils in Illinois (Wander et al., 1998); and in Vertisols in Australia (Mielke et al., 1986; Dalal, 1988). This lack of beneficial effect of NT may have been due to the high initial level of SOC and clay contents, which may have contributed to form very stable and time-resilient soil aggregates. Additional SOM

may not be protected within those already highly stable aggregates. This hypothesis refers to the concept of saturation capacity of soil to sequester SOC (Hassink, 1997; Hassink and Whitmore, 1997; Six et al., 2002). The initial SOC concentration seems to be an important parameter towards predicting the magnitude of SOC loss due to cultivation. However, Mann (1986) found that fractional loss of SOC (i.e. difference of SOC between cultivated and uncultivated situation, divided by the initial SOC content) increased with increasing initial SOC concentration. Davidson and Ackerman (1993) did not observe any “initial SOC” effect in their literature survey of SOC changes due to cultivation.

### 3.4. Comparison of soil organic carbon stock in no till and plow till soils

Several researchers have reported similar SOC stock under NT versus PT even in coarse-textured soils (Powlson and Jenkinson, 1981; Yang and Wander, 1999; Yang and Kay, 2001). Based on the approach of Paustian et al. (1997) who analyzed 27-paired experiments of NT versus PT, we expanded this approach to include the results of 56-paired (including the present study) experimental comparisons of SOC stocks reported for NT and PT treatments (Table 3). Among the 56 pairs, 42 had a positive increase of SOC under NT and 11 had a negative response. Among the 42 pairs with SOC gain, statistically significant differences were reported in only




Fig. 5. Soil organic carbon sequestration rate (mean and 95% confidence interval) under NT in response to soil type and intensity of cropping. The number of studies is in parenthesis.

Table 3  
References included in the dataset for meta-analysis

| Location                | Soil type            | Crop rotation          | Duration of NT (year) | Sampling depth (cm) | Rate of sequestration kg SOC ha <sup>-1</sup> year <sup>-1</sup> | References   |
|-------------------------|----------------------|------------------------|-----------------------|---------------------|--|--|
| St. Lambert, Que.       | Aeric Haplaquept     | Silage C               | 11                    | 24                  | 345  | Angers et al. (1995)                                   |
| Harrington, PEI         | Haplorthod           | W–B–B–SB               | 8                     | 60                  | –995   | Angers et al. (1997)                                   |
| La Pocatière, Que.      | Humaquept            | C–SB                   | 6                     | 60                  | –3380  | Angers et al. (1997)                                   |
| Normandin, Que.         | Humaquept            | C–SB                   | 3                     | 60                  | 900  | Angers et al. (1997)                                   |
| Ottawa, Ont.            | Eutrochrept          | C                      | 5                     | 60                  | 1196   | Angers et al. (1997)                                   |
| Ottawa, Ont.            | Eutrochrept          | C–W                    | 5                     | 60                  | 2966   | Angers et al. (1997)                                   |
| Dehli, Ont.             | Psamment             | C                      | 4                     | 60                  | –770   | Angers et al. (1997)                                   |
| Harrow, Ont.            | Haplaquoll           | C                      | 11                    | 60                  | –85  | Angers et al. (1997)                                   |
| Boigneville, France     | Typic Eutrochrep     | C                      | 18                    | Eq 30               | 139  | Balesdent et al. (1990)                                |
| Sydney, NB              | Pachic Haplustoll    | SW                     | 22                    | 20                  | 306  | Cambardella and Elliott (1992)                         |
| Swift Current SAS       | Typic Haploboroll    | W–F                    | 12                    | 15                  | 31   | Campbell et al. (1999)                                 |
| Swift Current SAS       | Typic Haploboroll    | W                      | 12                    | 15                  | 251  | Campbell et al. (1999)                                 |
| Lethbridge, ALB         | Typic Boroll         | W–F                    | 16                    | 14                  | –508   | Carter and Rennie (1982)                               |
| Melfort, SAS            | Udic Boroll          | W–F                    | 12                    | 20                  | 719  | Carter and Rennie (1982)                               |
| Watrous, SAS            | Typic Boroll         | W                      | 4                     | 10                  | –1052  | Carter and Rennie, 1982                                |
| Scott, SAS              | Typic Boroll         | W–W–W–W–2y of oilseeds | 2                     | 10                  | 2734   | Carter and Rennie (1982)                               |
| Warwick, Australia      | Udic Pellustert      | W or B                 | 14                    | 30                  | 128  | Dalal (1988)   |
| Hoytville, OH           | Mollic Epiaqualf     | C                      | 18                    | 20                  | 333  | Dick et al. (1986) recalculated from Lal et al. (1998) |
| Hoytville, OH           | Mollic Epiaqualf     | C–SB                   | 18                    | 20                  | 778  | Dick et al. (1986) recalculated from Lal et al. (1998) |
| Crossville, AL          | Typic Hapludult      | C (W as covercrop)     | 10                    | 20                  | 1717   | Edwards et al. (1992)                                  |
| Crossville, AL          | Typic Hapludult      | SB (W as covercrop)    | 10                    | 20                  | 1138   | Edwards et al. (1992)                                  |
| Crossville, AL          | Typic Hapludult      | C–SB (W as covercrop)  | 10                    | 20                  | 441  | Edwards et al. (1992)                                  |
| Horseshoe, GA           | Typic Rhodudult      | SG–W or SB             | 6                     | 21                  | 330  | Groffman (1984)  |
| Mandan, ND              | Typic Haplustoll     | SW–F                   | 12                    | 30.4                | –225   | Halvorson et al. (2002)                                |
| Mandan, ND              | Typic Haplustoll     | SW–WW–Sun              | 12                    | 30.4                | 542  | Halvorson et al. (2002)                                |
| Manhattan, KS           | Cumulic Hplustoll    | SB                     | 11                    | 15                  | 155  | Havlin and Kissel (1997)                               |
| Manhattan, KS           | Cumulic Hplustoll    | SG–SB                  | 11                    | 15                  | 345  | Havlin and Kissel (1997)                               |
| Manhattan, KS           | Cumulic Hplustoll    | SG                     | 11                    | 15                  | 109  | Havlin and Kissel (1997)                               |
| Griffin Farm, GA        | Typic Kanhapludult   | S or SG–WW–rye         | 17                    | 20                  | 588  | Hendricks et al. (1998)                                |
| Horseshoe, GA           | Rhodic Khanhapludult | S–WW                   | 17                    | 21                  | 353  | Hendricks et al. (1998)                                |
| Canterbury, New Zealand | Typic Eutrochept     | W–P–Cl–RG              | 9                     | 15                  | 3083   | Hermawan and Cameron (1993)                            |
| Lexington, KY           | Typic Paleudalf      | C (RG as covercrop)    | 20                    | 30                  | 165  | Ismail et al. (1994)                                   |
| Elwood, IL              | Aeric Ochraqualf     | C                      | 6                     | 30                  | 987  | Mielke et al. (1986)                                   |
| Waseca, MN              | Aquic Hapludoll      | C                      | 11                    | 15                  | 591  | Mielke et al. (1986)                                   |



|                                    |  |             |    |       |      |                              |
|------------------------------------|--|-------------|----|-------|------|------------------------------|
| Waseca, MN                         | Typic Endoaquoll                       | C           | 6  | 30    | –3   | Mielke et al. (1986)         |
| Lincoln, NB                        | Pachic Argiustoll                      | C           | 6  | 30    | 2523 | Mielke et al. (1986)         |
| Sydney, NB                         | Aridic Argiustoll                      | W–F         | 12 | 30    | 37   | Mielke et al. (1986)         |
| Corpus Christi, TX                 | Typic Orchraqualf                      | 4y Cot–4y C | 15 | 20    | 347  | Potter et al. (1998)         |
| Rothamsted, UK                     | Batcombe series in UK classification   | WW          | 5  | Eq 25 | –88  | Powlson and Jenkinson (1981) |
| Boxworth, UK                       | Hanslope series in UK classification   | WW          | 6  | Eq 25 | 507  | Powlson and Jenkinson (1981) |
| Headley Hall Leed’s University, UK | Wothersome series in UK classification | SB          | 8  | Eq 25 | 270  | Powlson and Jenkinson (1981) |
| Penicuik, Scotland                 | Macmery series in UK classification    | SB          | 10 | Eq 25 | –136 | Powlson and Jenkinson (1981) |
| Kellogs, BS, MI                    | Typic Hapludalf                        | C–SB–W      | 9  | 7.5   | 333  | Robertson et al. (2000)      |
| Sydney, NB                         | Pachic Haplustoll                      | WW–F        | 27 | 20    | 193  | Six et al. (1999)            |
| Wooster, OH                        | Typic Fragiudalf                       | C           | 34 | 20    | 125  | Six et al. (1999)            |
| Kellogs, BS, MI                    | Typic Hapludalf                        | C–S–W       | 34 | 20    | 63   | Six et al. (1999)            |
| Lexington, KY                      | Typic Paleudalf                        | C           | 25 | 20    | 247  | Six et al. (1999)            |
| Perry, IL                          | Acquic Argiudoll                       | C–SB        | 11 | 30    | 327  | Wander et al. (1998)         |
| Monmouth, IL                       | Acquic Hapludoll                       | C–SB        | 11 | 30    | 991  | Wander et al. (1998)         |
| Delkab, IL                         | Typic Haplaquall                       | C–SB        | 11 | 30    | 18   | Wander et al. (1998)         |
| Guelph, Ont.                       | Typic Hapludalf                        | Alf         | 20 | 20    | 263  | Yang and Kay (2001)          |
| Columbus, OH Don Scott Exp.        | Typic Argiaquoll                       | C–SB        | 8  | Eq 30 | 268  | This study                   |

C: corn; SB: soybean; WW: winter wheat; SW: spring wheat; F: fallow; B: barley; SG: sorghum; Sun: sunflower; Cot: cotton; Cl: clover; RG: ryegrass; Alf: alfafa, Eq: equivalent soil mass corresponding to this sampling depth under PT.

10 pairs. No studies actually indicated a significant decrease under NT. The increase of SOC stock upon conversion to NT averaged  $3.8 \text{ Mg ha}^{-1}$  but with large scatter ranging from  $-20$  to  $+27 \text{ Mg ha}^{-1}$ . In comparison with the SOC sequestration rates under NT of  $300\text{--}800 \text{ kg C ha}^{-1} \text{ year}^{-1}$  reported in the literature, (Paustian et al., 1997; Lal et al., 1998; Follett, 2001), we calculated an average rate of  $330 \text{ kg C ha}^{-1} \text{ year}^{-1}$  (Figs. 5 and 7). The 95% confidence interval of the sequestration rate was  $47\text{--}620 \text{ kg SOC ha}^{-1} \text{ year}^{-1}$  suggesting that conversion to NT had an overall positive effect on SOC sequestration (Fig. 5). Significantly positive sequestration rates were observed in coarse-textured soils and in systems with high residue inputs. As pointed out by Franzluebbers and Steiner (2002) climatic factors may be of primary importance in determining the SOC storage capacity in NT.

When SOC stocks in NT were plotted against those in PT (Fig. 6), the increase of SOC stock after conversion to NT was 4.6% of the initial SOC stock under PT with a 95% confidence interval of 1–8%. This coefficient of increase in SOC stock under NT was smaller than the one calculated by West and Post (2002). They concluded that conversion to NT could increase SOC stocks by 16%. Two uncertainties, or sources of error, can be pointed out in our dataset: (i) most of the studies reported stock for the same soil depth, therefore, leading to an overestimation of SOC stock under NT if bulk density was greater under NT; and (ii) few studies accounted for the litter residues in NT, this on the other hand, could lead to an underestimation of SOC stock under NT (Yang and Wander, 1999; Mann et al., 2002). There was no apparent effect of




Fig. 6. Soil organic carbon stock under NT plotted against SOC stock under PT. Linear regression plot (solid line) is represented with 95% confidence interval (long dotted line).

texture (as confirmed by the meta-analysis) or of the antecedent SOC stock (i.e., SOC under PT) effect on SOC, as reflected in Fig. 7.

#### 4. Conclusions

Our experimental study did not show a strong positive effect of NT on the overall SOC and N sequestration potential in this Mollisol. However, an increase in SOC concentration in the 0–5 cm soil layer of NT compared with PT was observed. One of the reasons for this lack of difference may have been the short duration of the experiment (e.g., 8 years). However, most of the changes in SOC are likely to occur within the first decade of conversion to NT. The mechanisms and pathways of SOC accrual are proba-




Fig. 7. Percent difference in stock under no till vs. plow till in relationship to SOC stock in plow tilled plots.

bly more complex and more dependent on site-specific pedo-climatic conditions and therefore less universal than simply reversing the trajectory of SOC loss by cultivation. From the literature analysis, we suggest a cautious approach be used to generalize the effect of NT on SOC stock. The magnitudal effect of NT on SOC stock was highly variable. Regardless, the conversion of conventional tillage to minimum or NT can still be promoted as a strategy for minimizing SOC losses during cropping.

## Acknowledgements

The authors wish to thank Dr. G. Hall, Dr. P. Jacinthe and J. Alder for their help during sampling as well as Dr. P. Curtis for his helpful discussions on statistical analysis and for his encouragement. The senior author was financially supported by the US Department of Energy, through the C-site project, managed by Drs. W.M. Post, J. Jastrow and C. Izauralde. We are grateful for their support.

## References

- Angers, D.A., Voroney, R.J.P., Cote, D., 1995. Dynamics of soil organic matter and corn residues affected by tillage practices. *Soil Sci. Soc. Am. J.* 59, 1311–1315.
- Angers, D.A., Bolinder, M.A., Carter, M.R., Gregorich, E.G., Drury, C.F., Liang, B.C., Voroney, R.P., Simard, R.R., Donald, R.G., Beyaert, R.P., Martel, J., 1997. Impact of tillage practices on organic carbon and nitrogen storage in cool, humid soils of eastern Canada. *Soil Till. Res.* 41, 191–201.
- Balesdent, J., Mariotti, A., Boisgontier, D., 1990. Effect of tillage on soil organic carbon mineralization estimated from  $^{13}\text{C}$  natural abundance. *J. Soil Sci.* 41, 587–596.
- Beare, M.H., Cabrera, M.L., Hendrix, P.F., Coleman, D.C., 1994. Aggregate-protected and unprotected organic matter pools in conventional and no-tillage soils. *Soil Sci. Soc. Am. J.* 58, 787–795.
- Blevins, R.L., Thomas, G.W., Smith, M.S., Frye, W.W., Cornelius, P.L., 1983. Changes in soil properties after 10 years continuous non-tilled and conventionally tilled corn. *Soil Till. Res.* 3, 135–146.
- Cambardella, C.A., Elliott, E.T., 1992. Particulate soil organic matter changes across a grassland cultivation sequence. *Soil Sci. Soc. Am. J.* 56, 777–783.
- Campbell, C.A., Zetner, R.P., 1993. Soil organic matter as influenced by crop rotations and fertilization. *Soil Sci. Soc. Am. J.* 57, 1034–1040.
- Campbell, C.A., Biederbeck, V.O., McConkey, B.G., Curtin, D., Zentner, R.P., 1999. Soil quality effect of tillage and fallow frequency. Soil organic matter quality as influenced by tillage and fallow frequency in a silt loam in southwestern Saskatchewan. *Soil Bio. Biochem.* 31, 1–7.
- Carter, M.R., Rennie, D.A., 1982. Changes in soil quality under zero tillage farming systems: Distribution of microbial biomass and mineralizable C and N potentials. *Can. J. Soil Sci.* 62, 587–597.
- Chan, K.Y., Heenan, D.P., Oates, A., 2002. Soil fractions and relationship to soil quality under different tillage and stubble management. *Soil Till. Res.* 63, 133–139.
- Chaney, K., Hodgson, D.R., Braim, M.A., 1985. The effect of direct drilling, shallow cultivation and ploughing on some physical properties in a long-term experiment on spring barley. *J. Agric. Sci.* 104, 125–133.
- Conant, R.T., Paustian, K., Elliott, E.T., 2001. Grassland management and conversion into grassland: Effects on soil carbon. *Ecol. Appl.* 11, 343–355.
- Dalal, R.C., 1988. Long-term effects of no-tillage, crop residues, and nitrogen application on properties of a Vertisol. *Soil Sci. Soc. Am. J.* 53, 1511–1515.
- Davidson, E.A., Ackerman, I.L., 1993. Changes in soil carbon inventories following cultivation of previously untilled soils. *Biogeochemistry* 20, 161–193.
- Dick, W.A., 1983. Organic carbon, nitrogen and phosphorus concentrations and pH in soil profiles as affected by tillage intensity. *Soil Sci. Soc. Am. J.* 47, 102–107.
- Dick, W.A., Van Doren Jr., D.M., Triplett Jr., G.J.B., Henry, J.E., 1986. Influence of long-term tillage and rotation combination on crop yields and selected soil parameters: results obtained for a Mollic Ochraqualf soil. *OARDC Research Bulletin*, Wooster, OH, 30 pp.
- Dick, W.A., McCoy, E.L., Edwards, W.M., Lal, R., 1991. Continuous application of no-tillage to Ohio soils. *Agron. J.* 83, 65–73.
- Dixon, P.M., 2001. The bootstrap and the jackknife: describing the precision of ecological studies. In: Scheiner, S.M., Gurevitch, J. (Eds.), *Design and Analysis of Ecological Experiments*, 2nd ed. Oxford University Press, Oxford, UK, pp. 267–288.
- Doran, J.W., 1980. Soil microbial and biochemical changes associated with reduced tillage. *Soil Sci. Soc. Am. J.* 44, 771–780.
- Edwards, J.H., Wood, C.W., Thurlow, D.L., Ruf, M.E., 1992. Tillage and crop rotation effects on fertility status of a Hapludult soil. *Soil Sci. Soc. Am. J.* 56, 1577–1582.
- Elliott, E.T., 1986. Aggregate structure and carbon, nitrogen and phosphorus in native and cultivated soils. *Soil Sci. Soc. Am. J.* 50, 627–633.
- Follett, R.F., 2001. Soil management concepts and carbon sequestration in cropland soils. *Soil Till. Res.* 61, 77–92.
- Franzluebbers, A.J., Arshad, M.A., 1997. Particulate organic carbon content and potential mineralization as affected by tillage and texture. *Soil Sci. Soc. Am. J.* 61, 1382–1386.
- Franzluebbers, A.J., Steiner, J.L., 2002. Climatic influences on soil organic carbon storage with no tillage. In: Kimble, J.M., et al. (Eds.), *Agricultural Practices and Policies for Carbon Sequestration in Soil*. Lewis Publishers, Boca Raton, FL, pp. 71–86.

- Franzluebbers, A.J., Stuedemann, J.A., 2002. Paniculate and non-particulate fractions of soil organic carbon under pastures in the southern Piedmont, USA. *Environ. Pollut.* 116 (Suppl.), S53–S62.
- Franzluebbers, A.J., Stuedemann, J.A., Schanberg, H.H., Wilkinson, S.R., 2000. Soil organic C and N pools under long-term pasture management in the southern Piedmont, USA. *Soil Biol. Biochem.* 32, 469–478.
- Groffman, P.M., 1984. Nitrification and denitrification in conventional and no-tillage soils. *Soil Sci. Soc. Am. J.* 49, 329–334.
- Guo, L.B., Gifford, R.M., 2002. Soil carbon stocks and land use change: a meta analysis. *Glob. Change Biol.* 8, 345–360.
- Gurevitch, J., Hedges, L.V., 1999. Statistical issues in ecological meta analysis. *Ecology* 80, 1142–1149.
- Gurevitch, J., Hedges, L.V., 2001. Meta-analysis: combining the results of independent experiments. In: Scheiner, S.M., Gurevitch, J. (Eds.), *Design and Analysis of Ecological Experiments*. Oxford University Press, Oxford, UK, pp. 347–369.
- Halvorson, A., Wienhold, B.J., Black, A.L., 2002. Tillage nitrogen and cropping system effects on soil carbon sequestration. *Soil Sci. Soc. Am. J.* 66, 906–912.
- Hassink, J., 1997. The capacity of soils to preserve organic C and N by their association with clay and silt particles. *Plant and Soil* 191, 77–87.
- Hassink, J., Whitmore, A.P., 1997. A model of the physical protection of soil organic matter in soils. *Soil Sci. Soc. Am. J.* 61, 131–138.
- Havlin, J.L., Kissel, D.E., Maddux, L.D., Claassen, M.M., Long, J.H., 1990. Crop rotation and tillage effects on soil organic carbon and nitrogen. *Soil Sci. Soc. Am. J.* 54, 448–452.
- Havlin, J.L., Kissel, D.E., 1997. Management effects on soil organic carbon and nitrogen in the east central plains of Kansas. In: Paul, E.A., et al. (Eds.), *Soil Organic Matter in Temperate Agroecosystems. Long-term Experiments in North America*. CRC Press, Boca Raton, FL, pp. 381–386.
- Haynes, R.J., Beare, M.H., 1996. Aggregation and soil organic matter storage in meso-thermal, humid soils. In: Carter, M.R., Stewart, B.A. (Eds.), *Structure and Organic Matter Storage in Agricultural Soils*. Lewis–CRC Press, Boca Raton, FL, pp. 213–262.
- Hendricks, P.F., Franzluebbers, A.J., McCracken, D.V., 1998. Management effects on C accumulation and loss in soils of the southern Appalachian Piedmont of Georgia. *Soil Till. Res.* 47, 245–251.
- Hermawan, D.J., Cameron, K.C., 1993. Structural changes in a silt loam under long-term conventional or minimum tillage. *Soil Till. Res.* 26, 139–150.
- Houghton, R.A., 1995. Land use change and the carbon cycle. *Glob. Change Biol.* 1, 275–287.
- Ismail, I., Blevins, R.L., Frye, W.W., 1994. Long-term no-tillage effects on soil properties and continuous corn yields. *Soil Sci. Soc. Am. J.* 58, 193–198.
- Johnson, D.W., Curtis, P.S., 2001. Effects of forest management on soil C and N storage: meta analysis. *For. Ecol. Manage.* 140, 227–238.
- Kay, B.D., VandenBygaart, A.J., 2002. Conservation tillage and depth stratification of porosity and soil organic matter. *Soil Till. Res.* 66, 107–118.
- Kern, I.S., Johnson, M.G., 1993. Conservation tillage impacts on national soil and atmospheric carbon levels. *Soil Sci. Soc. Am. J.* 57, 200–210.
- Kladivko, E.J., 2001. Tillage systems and soil ecology. *Soil Till. Res.* 61, 61–76.
- Lal, R., 1999. Soil compaction and tillage effects on soil physical properties of a Mollic Ochraqualf in northwest Ohio. *J. Sust. Agric.* 14, 53–65.
- Lal, R., Logan, T., Fausey, N.R., 1990. Long-term tillage effects on a Mollic Ochraqualf in Northwest Ohio. III. Soil nutrient profile. *Soil Till. Res.* 15, 351–382.
- Lal, R., Follett, R.F., Kimble, J.M., Cole, C.V., 1998. *The Potential of US Cropland to Sequester Carbon and Mitigate the Greenhouse Effect*. Lewis Publisher, Boca Raton, FL, 128 pp.
- Larson, W.E., Clapp, C.E., Pierre, W.R., Morachan, Y.B., 1972. Effects of increasing amount of organic residues on continuous corn. II. Organic carbon, nitrogen, phosphorus and sulfur. *Agron. J.* 64, 204.
- Manly, B.F., 1997. *Randomization, bootstrap and Monte Carlo methods in biology*, 2nd ed. Chapman & Hall, London, UK, 399 pp.
- Mann, L., Tolbert, V., Cushman, J., 2002. Review: potential environmental effects of corn (*Zea mays* L.) stover removal with emphasis on soil organic matter and erosion. *Agric. Ecosyst. Environ.* 89, 149–166.
- Mann, L.K., 1986. Changes in soil carbon storage after cultivation. *Soil Sci.* 142, 279–288.
- McLoda, N.A., Parkinson, R.J., 1980. *Soil Survey of Franklin County, Ohio*. United States Department of Agriculture, Soil Conservation Service.
- Mielke, L.N., Doran, J.W., Richards, K.A., 1986. Physical environment near the surface of plowed and no-tilled soils. *Soil Till. Res.* 7, 355–366.
- Mikhailova, E.A., Bryant, R.B., Vassenev, I.I., Schwager, S.J., Post, C.J., 2000. Cultivation effects on soil carbon and nitrogen contents at depth in the Russian Chernozem. *Soil Sci. Soc. Am. J.* 64, 738–745.
- Murty, D., Kirschbaum, M.U.F., McMurtie, R.E., McGilvray, A., 2002. Does conversion of forest to agricultural land change soil carbon and nitrogen? A review of the literature. *Glob. Change Biol.* 8, 105–123.
- Needelman, B.A., Wander, M.M., Bollero, G.A., Boast, C.W., Sims, G.K., Bullock, D.G., 1999. Interaction of tillage and soil texture: biologically active soil organic matter in Illinois. *Soil Sci. Soc. Am. J.* 63, 1326–1334.
- Paustian, K., Collins, H.P., Paul, E.A., 1997. Management controls on soil carbon. In: Paul, E.A., et al. (Eds.), *Soil Organic Matter in Temperate Agroecosystems: Long-Term Experiments in North America*. CRC Press, Boca Raton, FL, pp. 15–42.
- Post, W.M., Kwon, K.C., 2000. Soil carbon sequestration and land-use change: processes and potential. *Glob. Change Biol.* 6, 317–327.
- Potter, K.N., Torbert, H.A., Jones, O.R., Matocha, J.E., Morrisson, J.E.J., Unger, P.W., 1998. Distribution and amount of soil

- organic C in long-term management systems in Texas. *Soil Till. Res.* 47, 309–321.
- Powlson, D.S., Jenkinson, D.S., 1981. A comparison of the organic matter, biomass adenosine triphosphate and mineralizable nitrogen contents of ploughed and direct-drilled soils. *J. Agric. Sci.* 97, 713–721.
- Robertson, G.P., Paul, E.A., Harwood, R.R., 2000. Greenhouse gases in intensive agriculture contributions of individual gases to the radiative forcing of the atmosphere. *Science* 289, 1922–1925.
- Rosenberg, M.S., Adams, D.C., Gurevitch, J., 2000. MetaWin: Statistical Software for Meta-Analysis, Version 2.0. Sinauer Associates, Inc., Sunderland, MA.
- SAS Institute, 1999. SAS/STAT. Version 8.02 for Windows. SAS Institute, Inc., Cary, NC, USA.
- Schlesinger, W.H., 2000a. Carbon sequestration in soils: some cautions admits optimism. *Agric. Ecosyst. Environ.* 82, 121–127.
- Schlesinger, W.H., 2000b. Response to carbon cost of applying nitrogen fertilizer. *Science* 288, 809.
- Six, J., Elliott, E.T., Paustian, K., 1999. Aggregate and soil organic matter dynamics under conventional and no-tillage systems. *Soil Sci. Soc. Am. J.* 63, 1350–1358.
- Six, J., Elliott, E.T., Paustian, K., 2000. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Bio. Biochem.* 32, 2099–2103.
- Six, J., Conant, R.T., Paul, E.A., Paustian, K., 2002. Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant and Soil* 241, 155–176.
- Swift, R.S., 2001. Sequestration of carbon by soil. *Soil Sci.* 166, 858–871.
- Unger, P.W., 1982. Surface soil physical properties after 36 years of cropping to winter wheat. *Soil Sci. Soc. Am. J.* 46, 796–801.
- Wander, M.M., Bidart, M.G., Aref, S., 1998. Tillage impact on depth distribution of total and particulate organic matter in three Illinois soils. *Soil Sci. Soc. Am. J.* 62, 1704–1710.
- Watson, R.T., Noble, I.R., Bolin, B., Ravindranath, N.H., Verardo, D.J., Dokken, D.J., 2001. IPCC Special Report on Land-Use, Land-Use Change and Forestry. IPCC. Cambridge University Press, Cambridge, UK.
- West, T.O., Marland, G., 2002. A synthesis of carbon sequestration, carbon emissions, and net carbon flux in agriculture: comparing tillage practices in the United States. *Agric. Ecosyst. Environ.* 91, 217–232.
- West, T.O., Post, W.M., 2002. Soil organic carbon sequestration rates by tillage and crop rotation: a global data analysis. *Soil Sci. Soc. Am. J.* 66, 1930–1946.
- Yang, X.-M., Wander, M.M., 1999. Tillage effects on soil organic carbon distribution and storage in a silt loam soil in Illinois. *Soil Till. Res.* 52, 1–9.
- Yang, X.-M., Kay, B.D., 2001. Rotation and tillage effects on soil organic carbon sequestration in a typic Hapludalf in southeastern Ontario. *Soil Till. Res.* 59, 107–114.
- Zielke, R.C., Christenson, D.R., 1986. Organic carbon and nitrogen changes in soil under selected cropping systems. *Soil Sci. Soc. Am. J.* 50, 363–367.