

# EXAMINATION SCHEDULING

---

Alexander Eckl, Maximilian Fiedler, Mickael Grima,  
Roland Halbig

May 25, 2016

Technische Universität München

Problem

Modeling The Problem

Next Steps

# PROBLEM

---

Find a good examination schedule for the exam period of the TUM.

# CRITERIA FOR A GOOD EXAMINATION SCHEDULE

- Each exam is planned in exactly one period

# CRITERIA FOR A GOOD EXAMINATION SCHEDULE

- Each exam is planned in exactly one period
- In each room there is only one exam at a time

# CRITERIA FOR A GOOD EXAMINATION SCHEDULE

- Each exam is planned in exactly one period
- In each room there is only one exam at a time
- There are enough seats for each exam

# CRITERIA FOR A GOOD EXAMINATION SCHEDULE

- Each exam is planned in exactly one period
- In each room there is only one exam at a time
- There are enough seats for each exam
- No student has to write two exams at the same time



# CRITERIA FOR A GOOD EXAMINATION SCHEDULE

- Each exam is planned in exactly one period
- In each room there is only one exam at a time
- There are enough seats for each exam
- No student has to write two exams at the same time
- Rooms for an exam are minimized

# CRITERIA FOR A GOOD EXAMINATION SCHEDULE

- Each exam is planned in exactly one period
- In each room there is only one exam at a time
- There are enough seats for each exam
- No student has to write two exams at the same time
- Rooms for an exam are minimized
- Time between exams is maximized

# MODELING THE PROBLEM

---

$$x_{i,k,l} := \begin{cases} 1, & \text{if exam } i \text{ is written in period } l \text{ in room } k \\ 0, & \text{otherwise} \end{cases}$$

$$x_{i,k,l} := \begin{cases} 1, & \text{if exam } i \text{ is written in period } l \text{ in room } k \\ 0, & \text{otherwise} \end{cases}$$

$$y_{i,l} := \begin{cases} 1, & \text{if exam } i \text{ is written in period } l \\ 0, & \text{otherwise} \end{cases}$$

- Connecting the variables  $x$  and  $y$ :

- Connecting the variables  $x$  and  $y$ :

$$(1) \sum_{\text{rooms } k} x_{i,k,l} \leq y_{i,l} \cdot M \quad \forall \text{ exams } i, \forall \text{ periods } l$$

- Connecting the variables  $x$  and  $y$ :

$$(1) \sum_{\text{rooms } k} x_{i,k,l} \leq y_{i,l} \cdot M \quad \forall \text{ exams } i, \forall \text{ periods } l$$

$$(2) \sum_{\text{rooms } k} x_{i,k,l} \geq y_{i,l} \quad \forall \text{ exams } i, \forall \text{ periods } l$$



# CONSTRAINTS

- Each exam is planned in exactly one period:

- Each exam is planned in exactly one period:

$$(3) \sum_{\text{periods } l} y_{i,l} = 1 \quad \forall \text{ exams } i$$

- Each exam is planned in exactly one period:

$$(3) \quad \sum_{\text{periods } l} y_{i,l} = 1 \quad \forall \text{ exams } i$$

$$\text{c.f. (2)} \quad \sum_{\text{rooms } k} x_{i,k,l} \geq y_{i,l} \quad \forall \text{ exams } i, \forall \text{ periods } l$$

- There are enough seats for the students in the exam rooms:

- There are enough seats for the students in the exam rooms:

$$(4) \quad \sum_{\substack{\text{periods } l, \\ \text{rooms } k}} c_k \cdot x_{i,k,l} \geq s_i \quad \forall \text{ exams } i$$

$s_i :=$  # students taking exam  $i$

$c_k :=$  # seats in room  $k$

- In every room there is only one exam at a given time:

- In every room there is only one exam at a given time:

$$(5) \sum_{\text{exams } i} x_{i,k,l} \leq 1 \quad \forall \text{ rooms } k, \forall \text{ periods } l$$

- There are no conflicts for students taking multiple exams:



- There are no conflicts for students taking multiple exams:

$$(6) \quad \sum_{\substack{\text{exams } j, \\ j \text{ conflicts with } i}} y_{j,l} \leq (1 - y_{i,l}) \cdot M \quad \forall \text{ exams } i, \forall \text{ periods } l$$

- Minimize the total number of rooms:

$$(a) \min \sum_{\substack{\text{exams } i, \\ \text{rooms } k, \\ \text{periods } l}} x_{i,k,l}$$

- Maximize the time between two exams:

$$(b) \max \sum_{\substack{\text{exams } i, j \\ i \text{ conflicts with } j}} d_{i,j}$$

$d_{i,j} :=$  distance between exams  $i$  and  $j$

# OBJECTIVE FUNCTION

- Combine the previous two objective functions using a weighting factor  $\gamma > 0$ :

$$(a + b) \min \sum_{\substack{\text{exams } i, \\ \text{rooms } k, \\ \text{periods } l}} x_{i,k,l} - \gamma \cdot \sum_{\substack{\text{exams } i, j \\ i \text{ conflicts with } j}} d_{i,j}$$

## NEXT STEPS


---

# FINDING A FEASIBLE STARTING SOLUTION

Compute planning  
order of exams

# FINDING A FEASIBLE STARTING SOLUTION


Compute planning  
order of exams



```
graph TD; A[Compute planning order of exams] --> B[Plan exam most difficult to plan];
```


Plan exam most  
difficult to plan

# FINDING A FEASIBLE STARTING SOLUTION







# FINDING A FEASIBLE STARTING SOLUTION




# FINDING A FEASIBLE STARTING SOLUTION



# FINDING A FEASIBLE STARTING SOLUTION



# FINDING A FEASIBLE STARTING SOLUTION



- Add Clique Constraints:

$$(7) \quad \sum_{i \text{ in clique}} y_{i,1} \leq 1 \quad \forall \text{ cliques}$$

- Add Clique Constraints:

$$(7) \quad \sum_{i \text{ in clique}} y_{i,l} \leq 1 \quad \forall \text{ cliques}$$

- Remove absolute value in the objective function.

- Add Clique Constraints:

$$(7) \quad \sum_{i \text{ in clique}} y_{i,l} \leq 1 \quad \forall \text{ cliques}$$

- Remove absolute value in the objective function.
- Improve running time using heuristics, pre-solving, etc.

- Add Clique Constraints:

$$(7) \quad \sum_{i \text{ in clique}} y_{i,1} \leq 1 \quad \forall \text{ cliques}$$

- Remove absolute value in the objective function.
- Improve running time using heuristics, pre-solving, etc.
- Use a path based model and column generation





Figure 1: Server, hopefully somewhere in Germany