

EXAMINATION SCHEDULING

Alexander Eckl, Maximilian Fiedler, Mickael Grima, Roland Halbig

March 21, 2016

Technische Universität München

Problem

ILP Formulation

Next Steps and Improvements

PROBLEM

Find a good examination schedule for the exam period of the TUM

1. Each exam is planed in exactly one period

MOST IMPORTANT CRITERIA FOR AN EXAMINATION SCHEDULE

1. Each exam is planed in exactly one period
2. There are enough seats

MOST IMPORTANT CRITERIA FOR AN EXAMINATION SCHEDULE

1. Each exam is planed in exactly one period
2. There are enough seats
3. In each room there is only one exam at a time

MOST IMPORTANT CRITERIA FOR AN EXAMINATION SCHEDULE

1. Each exam is planed in exactly one period
2. There are enough seats
3. In each room there is only one exam at a time
4. No student has to write two exams at the same time

MOST IMPORTANT CRITERIA FOR AN EXAMINATION SCHEDULE

1. Each exam is planed in exactly one period
2. There are enough seats
3. In each room there is only one exam at a time
4. No student has to write two exams at the same time
5. Rooms for an exam are minimized

MOST IMPORTANT CRITERIA FOR AN EXAMINATION SCHEDULE

1. Each exam is planed in exactly one period
2. There are enough seats
3. In each room there is only one exam at a time
4. No student has to write two exams at the same time
5. Rooms for an exam are minimized
6. Time between exams is maximized

ILP FORMULATION

$$x_{i,k,l} := \begin{cases} 1, & \text{if exam } i \text{ is written in period } l \text{ in room } k \\ 0, & \text{sonst} \end{cases}$$

$$y_{i,l} := \begin{cases} 1, & \text{if exam } i \text{ is written in period } l \\ 0, & \text{sonst} \end{cases}$$

0. Connecting variables x and y

$$(1) \quad \sum_{\forall \text{ rooms } k} x_{i,k,l} \leq y_{i,l} \cdot M \quad \forall \text{ exams } i, \forall \text{ periods } l$$

0. Connecting variables x and y

$$(1) \quad \sum_{\forall \text{ rooms } k} x_{i,k,l} \leq y_{i,l} \cdot M \quad \forall \text{ exams } i, \forall \text{ periods } l$$

$$(2) \quad \sum_{\forall \text{ rooms } k} x_{i,k,l} \geq y_{i,l} \quad \forall \text{ exams } i, \forall \text{ periods } l$$

1. Each exam is planned in exactly one period

$$(3) \quad \sum_{\forall \text{ periods } l} y_{i,l} = 1 \quad \forall \text{ exams } i$$

1. Each exam is planned in exactly one period

$$(3) \quad \sum_{\forall \text{ periods } l} y_{i,l} = 1 \quad \forall \text{ exams } i$$

$$(2) \quad \sum_{\forall \text{ rooms } k} x_{i,k,l} \geq y_{i,l} \quad \forall \text{ exams } i, \forall \text{ periods } l$$

2. There must be enough seats for the students in the chosen rooms

$$\sum_{\substack{\forall \text{ periods } l, \\ \forall \text{ rooms } k}} c_k x_{i,k,l} \geq s_i \quad \forall \text{ exams } i$$

$s_i :=$ # students taking exam i .

$c_k :=$ # seats in room k .

3. In each room there is only one exam at a time

$$\sum_{\forall \text{ exams } i} x_{i,k,l} \leq t_{k,l} \quad \forall \text{ rooms } k, \forall \text{ periods } l$$

4. No student has to write two exams at the same time

$$\sum_{\substack{\forall \text{ exams } j: \\ i,j \text{ have a conflict}}} y_{j,l} \leq (1 - y_{i,l}) * M \quad \forall \text{ exams } i, \forall \text{ periods } l$$

IMPROVEMENTS AND NEXT STEPS

s_i := Number of students signed up for exam i .

c_k := Number of available seats in the lecture room k .

Q := Kollisionsmatrix

$q_{i,j} := \begin{cases} 0, & \text{falls Prüfung } i \text{ und } j \text{ gleichzeitig stattfinden können} \\ 1, & \text{sonst} \end{cases}$

T := Sperrmatrix

$t_{i,j} := \begin{cases} 1, & \text{falls Raum } k \text{ zum Zeitintervall } l \text{ geöffnet ist} \\ 0, & \text{sonst} \end{cases}$

h_l := Anzahl der Stunden von Periode l nach Beginn des Prüfungszeitraumes

$$x_{i,k,l} := \begin{cases} 1, & \text{wenn Prüfung } i \text{ zum Zeitpunkt } l \text{ in Raum } k \text{ stattfindet} \\ 0, & \text{sonst} \end{cases}$$

$$y_{i,l} := \begin{cases} 1, & \text{wenn Prüfung } i \text{ im Zeitintervall } l \text{ stattfindet} \\ 0, & \text{sonst} \end{cases}$$

Dimensionen:

n : Anzahl der Prüfungen

r : Anzahl der Räume

p : Anzahl der Zeitintervalle

1. Verknüpfung der Variablen

$$\sum_{k=1}^r x_{i,k,l} \leq y_{i,l} \cdot r \quad \forall i \in [n] \forall l \in [p]$$

$$\sum_{k=1}^r x_{i,k,l} \geq y_{i,l} \quad \forall i \in [n] \forall l \in [p]$$

2. Jede Prüfung wird auf genau einem Zeitintervall eingeplant

$$\sum_{l=1}^p y_{i,l} = 1 \quad \forall i \in [n]$$

3. Konfliktvermeidung

$$\sum_{j=1, j>i}^n q_{i,j} y_{j,l} \leq (1 - y_{i,l}) \sum_{\nu=1}^n q_{i,\nu} \quad \forall i \in [n], \forall l \in [p]$$

5. Alle Studierenden bekommen einen Platz

$$\sum_{l=1}^p \sum_{k=1}^r c_{k,i,k,l} x_{i,k,l} \geq s_i \quad \forall i \in [n]$$

6. Jedem Raum wird je Zeit maximal eine Prüfung zugeteilt

$$\sum_{i=1}^n x_{i,k,l} \leq t_{k,l} \quad \forall k \in [r], \forall l \in [p]$$

7. Clique Constraints

$$\sum_{j \text{ in clique}} y_{i,l} \leq 1 \quad \forall l \in [p]$$

$$\min \sum_{i=1}^n \sum_{k=1}^r \sum_{l=1}^p s_i x_{i,k,l} - \gamma \min_{j>i: q_{i,j}>0} |\Delta h_{i,j}|$$

where $\Delta h_{i,j} := \sum_{l=1}^p h_l(y_{i,l} - y_{j,l})$. Resolving abs:

$$\begin{aligned} \min \quad & \sum_{i=1}^n \sum_{k=1}^r s_i x_{i,k} - \gamma w \\ \text{s.t.} \quad & z_{i,j} \leq \Delta h_{i,j} + \delta_{i,j}(h_p - h_1) \quad \forall i, j \in [n] \\ & z_{i,j} \leq -\Delta h_{i,j} + (1 - \delta_{i,j})(h_p - h_1) \quad \forall i, j \in [n] \\ & z_{i,j} \geq \Delta h_{i,j} \quad \forall i, j \in [n] \\ & z_{i,j} \geq -\Delta h_{i,j} \quad \forall i, j \in [n] \\ & w \geq z_{i,j} \quad \forall i, j \in [n] \end{aligned}$$

<http://lpsolve.sourceforge.net/5.1/absolute.htm>