


THE PUMPING LEMMA

THE PUMPING LEMMA

Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that

- (1) $x = uvw$
- (2) $|uv| \leq n$
- (3) $|v| \geq 1$
- (4) for all $i \geq 0$: $uv^i w \in L$.




THE PUMPING LEMMA

Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that

- (1) $x = uvw$
- (2) $|uv| \leq n$
- (3) $|v| \geq 1$
- (4) for all $i \geq 0$: $uv^i w \in L$.


$\xleftarrow{\hspace{1cm} n \hspace{1cm}}$



THE PUMPING LEMMA

Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that



- (1) $x = uvw$
- (2) $|uv| \leq n$
- (3) $|v| \geq 1$
- (4) for all $i \geq 0$: $uv^i w \in L$.



THE PUMPING LEMMA

Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that



- (1) $x = uvw$
- (2) $|uv| \leq n$
- (3) $|v| \geq 1$
- (4) for all $i \geq 0$: $uv^i w \in L$.



THE PUMPING LEMMA

Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that



- (1) $x = uvw$
- (2) $|uv| \leq n$
- (3) $|v| \geq 1$
- (4) for all $i \geq 0$: $uv^i w \in L$.



THE PUMPING LEMMA

Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that



- (1) $x = uvw$
- (2) $|uv| \leq n$
- (3) $|v| \geq 1$
- (4) for all $i \geq 0$: $uv^i w \in L$.



THE PUMPING LEMMA

Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that

- (1) $x = uvw$
- (2) $|uv| \leq n$
- (3) $|v| \geq 1$
- (4) for all $i \geq 0$: $uv^i w \in L$.



THE PUMPING LEMMA


Theorem. For any regular language L there exists an integer n , such that for all $x \in L$ with $|x| \geq n$, there exist $u, v, w \in \Sigma^*$, such that

$$(1) \quad x = uvw$$

$$(2) \quad |uv| \leq n$$

$$(3) \quad |v| \geq 1$$

$$(4) \quad \text{for all } i \geq 0: uv^i w \in L.$$



PROOF OF P.L. (SKETCH)

PROOF OF P.L. (SKETCH)

Let M be a DFA for L . Take n be the number of states of M plus 1.

PROOF OF P.L. (SKETCH)

Let M be a DFA for L . Take n be the number of states of M plus 1.

Take any $x \in L$ with $|x| \geq n$. Consider the path (from start state to an accepting state) in M that corresponds to x . The length of this path is $|x| \geq n$.

PROOF OF P.L. (SKETCH)

Let M be a DFA for L . Take n be the number of states of M plus 1.

Take any $x \in L$ with $|x| \geq n$. Consider the path (from start state to an accepting state) in M that corresponds to x . The length of this path is $|x| \geq n$.


Since M has at most $n - 1$ states, some state must be visited twice or more in the first n steps of the path.

PROOF OF P.L. (SKETCH)

Let M be a DFA for L . Take n be the number of states of M plus 1.

Take any $x \in L$ with $|x| \geq n$. Consider the path (from start state to an accepting state) in M that corresponds to x . The length of this path is $|x| \geq n$.

Since M has at most $n - 1$ states, some state must be visited twice or more in the first n steps of the path.




PROOF OF P.L. (SKETCH)

Let M be a DFA for L . Take n be the number of states of M plus 1.

Take any $x \in L$ with $|x| \geq n$. Consider the path (from start state to an accepting state) in M that corresponds to x . The length of this path is $|x| \geq n$.

Since M has at most $n - 1$ states, some state must be visited twice or more in the first n steps of the path.



$$\begin{aligned} x &= uw \in L \\ &= uvw \in L \\ &= uvw \in L \\ &= uvvw \in L \\ &\dots \end{aligned}$$

USING PUMPING LEMMA TO PROVE NON-REGULARITY

| | | |
|-----------------|--------------|--------------------------|
| L regular | \implies | L satisfies P.L. |
| L non-regular | \implies | ? |
| L non-regular | \Leftarrow | L doesn't satisfy P.L. |

Negation:

$\exists n \in \mathbb{N} \ \forall x \in L \text{ with } |x| \geq n$

$\exists u, v, w \in \Sigma^*$

all of these hold:

$$(1) \ x = uvw$$

$$(2) \ |uv| \leq n$$

$$(3) \ |v| \geq 1$$

$$(4) \ \forall i \geq 0: uv^i w \in L.$$

USING PUMPING LEMMA TO PROVE NON-REGULARITY

L regular $\implies L$ satisfies P.L.

L non-regular $\implies ?$

L non-regular $\iff L$ doesn't satisfy P.L.

Negation:

$\exists n \in \mathbb{N} \ \forall x \in L \text{ with } |x| \geq n$

$\exists u, v, w \in \Sigma^*$

all of these hold:

$$(1) \ x = uvw$$

$$(2) \ |uv| \leq n$$

$$(3) \ |v| \geq 1$$

$$(4) \ \forall i \geq 0: uv^i w \in L.$$

USING PUMPING LEMMA TO PROVE NON-REGULARITY

L regular $\implies L$ satisfies P.L.

L non-regular $\implies ?$

L non-regular $\iff L$ doesn't satisfy P.L.

Negation:

$\exists n \in \mathbb{N} \forall x \in L$ with $|x| \geq n$

$\exists u, v, w \in \Sigma^*$

all of these hold:

(1) $x = uvw$

(2) $|uv| \leq n$

(3) $|v| \geq 1$

(4) $\forall i \geq 0: uv^i w \in L.$

$\forall n \in \mathbb{N} \exists x \in L$ with $|x| \geq n$

$\forall u, v, w \in \Sigma^*$

not all of these hold:

(1) $x = uvw$

(2) $|uv| \leq n$

(3) $|v| \geq 1$

(4) $\forall i \geq 0: uv^i w \in L.$

USING PUMPING LEMMA TO PROVE NON-REGULARITY

L regular $\implies L$ satisfies P.L.

L non-regular $\implies ?$

L non-regular $\iff L$ doesn't satisfy P.L.

Negation:

$\exists n \in \mathbb{N} \forall x \in L$ with $|x| \geq n$

$\exists u, v, w \in \Sigma^*$

all of these hold:

$\forall n \in \mathbb{N} \exists x \in L$ with $|x| \geq n$

$\forall u, v, w \in \Sigma^*$

not all of these hold:

(1)

Equivalently:

(2)

$(1) \wedge (2) \wedge (3) \Rightarrow \text{not}(4)$

(3)

where **not**(4) is:

$\exists i : uv^i w \notin L$

(4)

$\checkmark \quad v \subseteq \dots \omega \omega \omega \subset L.$

(1) $x = uvw$

(2) $|uv| \leq n$

(3) $|v| \geq 1$

(4) $\forall i \geq 0: uv^i w \in L.$

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If (1), (2), (3) hold then $x = 0^n 1^n = uvw$ with $|uv| \leq n$ and $|v| \geq 1$.

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If (1), (2), (3) hold then $x = 0^n 1^n = uvw$ with $|uv| \leq n$ and $|v| \geq 1$.

So, $u = 0^s$, $v = 0^t$, $w = 0^p 1^n$ with

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If (1), (2), (3) hold then $x = 0^n 1^n = uvw$ with $|uv| \leq n$ and $|v| \geq 1$.

So, $u = 0^s$, $v = 0^t$, $w = 0^p 1^n$ with

$$s + t \leq n, \quad t \geq 1, \quad p \geq 0, \quad s + t + p = n.$$

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If (1), (2), (3) hold then $x = 0^n 1^n = uvw$ with $|uv| \leq n$ and $|v| \geq 1$.

So, $u = 0^s$, $v = 0^t$, $w = 0^p 1^n$ with

$$s + t \leq n, \quad t \geq 1, \quad p \geq 0, \quad s + t + p = n.$$

But then (4) fails for $i = 0$:

EXAMPLE 1

Prove that $L = \{0^i 1^i : i \geq 0\}$ is NOT regular.

Proof. Show that P.L. doesn't hold (*note: showing P.L. holds doesn't mean regularity*).

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^n 1^n$

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If (1), (2), (3) hold then $x = 0^n 1^n = uvw$ with $|uv| \leq n$ and $|v| \geq 1$.

So, $u = 0^s$, $v = 0^t$, $w = 0^p 1^n$ with

$$s + t \leq n, \quad t \geq 1, \quad p \geq 0, \quad s + t + p = n.$$

But then (4) fails for $i = 0$:

$$uv^0w = uw = 0^s 0^p 1^n = 0^{s+p} 1^n \notin L, \quad \text{since } s + p \neq n$$

IN PICTURE

$\exists u, v, w$ such that

(1) $x = uvw$

(2) $|uv| \leq n$

(3) $|v| \geq 1$

(4) $\forall i \in \mathbb{N} : uv^i w \in L.$

IN PICTURE

$\exists u, v, w$ such that

$$(1) \ x = uvw$$

$$(2) \ |uv| \leq n$$

$$(3) \ |v| \geq 1$$

$$(4) \ \forall i \in \mathbb{N} : uv^i w \in L.$$



IN PICTURE

$\exists u, v, w$ such that


(1) $x = uvw$

(2) $|uv| \leq n$

(3) $|v| \geq 1$

(4) $\forall i \in \mathbb{N} : uv^i w \in L.$

non-empty



IN PICTURE

$\exists u, v, w$ such that


$$(1) \ x = uvw$$

$$(2) \ |uv| \leq n$$

$$(3) \ |v| \geq 1$$

$$(4) \ \forall i \in \mathbb{N} : uv^i w \in L.$$

non-empty



If (1), (2), (3) hold then (4) fails: it is not the case that for all i , $uv^i w$ is in L .

In particular, let $i = 0$. $uw \notin L$.

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^m$ where $m \geq n + 2$ is prime.

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^m$ where $m \geq n + 2$ is prime.

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^m$ where $m \geq n + 2$ is prime.

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^m$ where $m \geq n + 2$ is prime.

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If 0^m is written as $0^m = uvw$, then $0^m = 0^{|u|}0^{|v|}0^{|w|}$.

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^m$ where $m \geq n + 2$ is prime.

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If 0^m is written as $0^m = uvw$, then $0^m = 0^{|u|}0^{|v|}0^{|w|}$.

If $|uv| \leq n$ and $|v| \geq 1$, then consider $i = |v| + |w|$:

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^m$ where $m \geq n + 2$ is prime.

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If 0^m is written as $0^m = uvw$, then $0^m = 0^{|u|}0^{|v|}0^{|w|}$.

If $|uv| \leq n$ and $|v| \geq 1$, then consider $i = |v| + |w|$:

$$\begin{aligned} uv^i w &= 0^{|v|} 0^{|v|} (|v| + |w|) 0^{|w|} \\ &= 0^{(|v|+1)(|v|+|w|)} \notin L \end{aligned}$$

EXAMPLE 2

Prove that $L = \{0^i : i \text{ is a prime}\}$ is NOT regular.

Proof. We show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Now let $x = 0^m$ where $m \geq n + 2$ is prime.

$x \in L$ and $|x| \geq n$, so by P.L. $\exists u, v, w$ such that (1)–(4) hold.

We show that $\forall u, v, w$ (1)–(4) don't all hold.

If 0^m is written as $0^m = uvw$, then $0^m = 0^{|u|}0^{|v|}0^{|w|}$.

If $|uv| \leq n$ and $|v| \geq 1$, then consider $i = |v| + |w|$:

$$\begin{aligned} uv^i w &= 0^{|v|} 0^{|v|} (|v| + |w|) 0^{|w|} \\ &= 0^{(|v|+1)(|v|+|w|)} \notin L \end{aligned}$$

Both factors ≥ 2

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Let us consider $x = 0^n 0^n \in L$. Obviously $|x| \geq n$.

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Let us consider $x = 0^n 0^n \in L$. Obviously $|x| \geq n$.

Can $0^n 0^n$ be written as $0^n 0^n = uvw$ such that $|uv| \leq n$ $|v| \geq 1$ and that for all i : $uv^i w \in L$?

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Let us consider $x = 0^n 0^n \in L$. Obviously $|x| \geq n$.

Can $0^n 0^n$ be written as $0^n 0^n = uvw$ such that $|uv| \leq n$ $|v| \geq 1$ and that for all i : $uv^i w \in L$?

YES! Let $u = \epsilon$, $v = 00$, and $w = 0^{2n-2}$.

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Let us consider $x = 0^n 0^n \in L$. Obviously $|x| \geq n$.

Can $0^n 0^n$ be written as $0^n 0^n = uvw$ such that $|uv| \leq n$ $|v| \geq 1$ and that for all i : $uv^i w \in L$?

YES! Let $u = \epsilon$, $v = 00$, and $w = 0^{2n-2}$.

Then $\forall i$, $uv^i w$ is of the form $0^{2k} = 0^k 0^k$.

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Let us consider $x = 0^n 0^n \in L$. Obviously $|x| \geq n$.

Can $0^n 0^n$ be written as $0^n 0^n = uvw$ such that $|uv| \leq n$ $|v| \geq 1$ and that for all i : $uv^i w \in L$?

YES! Let $u = \epsilon$, $v = 00$, and $w = 0^{2n-2}$.

Then $\forall i$, $uv^i w$ is of the form $0^{2k} = 0^k 0^k$.

Does this mean that L is regular?

EXAMPLE 3

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Again we try to show that P.L. doesn't hold.

If L is regular, then by P.L. $\exists n$ such that ...

Let us consider $x = 0^n 0^n \in L$. Obviously $|x| \geq n$.

Can $0^n 0^n$ be written as $0^n 0^n = uvw$ such that $|uv| \leq n$ $|v| \geq 1$ and that for all i : $uv^i w \in L$?

YES! Let $u = \epsilon$, $v = 00$, and $w = 0^{2n-2}$.

Then $\forall i$, $uv^i w$ is of the form $0^{2k} = 0^k 0^k$.

Does this mean that L is regular?

NO. We have chosen a bad string x . To show that L fails the P.L., we only need to exhibit some x that cannot be “pumped” (and $|x| \geq n$).

EXAMPLE 3, 2ND ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

EXAMPLE 3, 2ND ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = (01)^n(01)^n$. Obviously $x \in L$ and $|x| \geq n$.

EXAMPLE 3, 2ND ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = (01)^n(01)^n$. Obviously $x \in L$ and $|x| \geq n$.

Q: Can x be “pumped” for some choice of u, v, w with $|uv| \leq n$ and $|v| \geq 1$?

EXAMPLE 3, 2ND ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = (01)^n(01)^n$. Obviously $x \in L$ and $|x| \geq n$.

Q: Can x be “pumped” for some choice of u, v, w with $|uv| \leq n$ and $|v| \geq 1$?

A: Yes! Take $u = \epsilon$, $v = 0101$, $w = (01)^{2n-2}$.

EXAMPLE 3, 2ND ATTEMPT

Prove that $L = \{yy : y \in \{0,1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = (01)^n(01)^n$. Obviously $x \in L$ and $|x| \geq n$.

Q: Can x be “pumped” for some choice of u, v, w with $|uv| \leq n$ and $|v| \geq 1$?

A: Yes! Take $u = \epsilon$, $v = 0101$, $w = (01)^{2n-2}$.

Another bad choice of x !

EXAMPLE 3, 3RD ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

EXAMPLE 3, 3RD ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = 0^n 1 0^n 1$. Again $x \in L$ and $|x| \geq n$.

EXAMPLE 3, 3RD ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = 0^n 1 0^n 1$. Again $x \in L$ and $|x| \geq n$.

$\forall u, v, w$ such that $0^n 1 0^n 1 = uvw$ and $|uv| \leq n$ and $|v| \geq 1$:

EXAMPLE 3, 3RD ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = 0^n 1 0^n 1$. Again $x \in L$ and $|x| \geq n$.

$\forall u, v, w$ such that $0^n 1 0^n 1 = uvw$ and $|uv| \leq n$ and $|v| \geq 1$: must have uv contained in the first group of 0^n . Thus consider

EXAMPLE 3, 3RD ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = 0^n 1 0^n 1$. Again $x \in L$ and $|x| \geq n$.

$\forall u, v, w$ such that $0^n 1 0^n 1 = uvw$ and $|uv| \leq n$ and $|v| \geq 1$: must have uv contained in the first group of 0^n . Thus consider

$$uv^0 w = 0^{n-|v|} 1 0^n 1.$$

EXAMPLE 3, 3RD ATTEMPT

Prove that $L = \{yy : y \in \{0, 1\}^*\}$ is NOT regular.

Given n from the P.L., let $x = 0^n 1 0^n 1$. Again $x \in L$ and $|x| \geq n$.

$\forall u, v, w$ such that $0^n 1 0^n 1 = uvw$ and $|uv| \leq n$ and $|v| \geq 1$: must have uv contained in the first group of 0^n . Thus consider

$$uv^0 w = 0^{n-|v|} 1 0^n 1.$$

Since $|v|$ is at least 1, this is clearly not of the form yy .