Tumour-core, peritumoral-margin and remote-context evidence in brain MRI classification: a sham-controlled counterfactual audit with descriptive plane stratification across CNN and Vision Transformer models

Chhotu Rajput1* and Rishi Gupta2

[1]Manipal University Jaipur, Jaipur, Rajasthan, India.
[2]Department of Computer Science and Engineering, School of Computer Science and Engineering, Faculty of Science, Technology and Education, Manipal University Jaipur, Jaipur, Rajasthan, India.

*Corresponding author(s). E-mail(s): chhotu.2427030521@muj.manipal.edu;
Contributing authors: rishi.gupta@jaipur.manipal.edu;

Abstract

Deep convolutional neural networks (CNNs) and Vision Transformers (ViTs) can achieve high benchmark performance on brain MRI classification, but performance on unmodified images does not establish which anatomical regions support a prediction. Tumour-core morphology, adjacent peritumoral tissue, remote parenchyma and generic image-perturbation effects should be distinguished before model behaviour is interpreted as clinically meaningful evidence. We conducted a sham-controlled, region-resolved counterfactual audit on two contrast-enhanced MRI cohorts: the Cheng dataset (3,064 slices from 233 patients; three tumour classes) and the multi-plane BRISC dataset (6,000 slices; glioma, meningioma, pituitary tumour and no-tumour classes). Cheng partitions were generated with patient-grouped five-fold stratification; the exported baseline prediction file represents the held-out Fold 1 partition (480 slices from 49 patients), not pooled predictions from all five folds. ResNet-50 and DeiT-Small/16 were fine-tuned from ImageNet-pretrained weights. For each eligible tumour-containing slice we defined the tumour core, a 4-pixel immediate margin, a 12-pixel outer margin, remote tissue and a contralateral sham region, and perturbed each region with Telea inpainting and locally matched Gaussian texture filling. Technical non-target pixel invariance ($|\Delta| < 10^{-5}$) was verified across an image quality control pool of 120 slices (60 per dataset; 1,200 intervention records). The model sensitivity audit evaluated 90 matched slices (45 per dataset), two architectures, five regions and two perturbation methods (1,800 paired prediction observations). Effects were summarised as raw changes in ground-truth-class probability and sham-corrected sensitivity (τ_r), evaluated using paired Wilcoxon tests, family-wise Holm adjustment, cluster-level analyses across 55 patient/case clusters, and linear mixed-effects models. On the Cheng Fold 1 partition, DeiT-Small achieved accuracy 0.8458, macro-F1 0.8403 and expected calibration error (ECE) 0.0366; ResNet-50 achieved 0.7583, 0.7190 and 0.1835. On the BRISC clean 899-image test set, DeiT-Small achieved accuracy 0.8265 and macro-F1 0.8344, against 0.6963 and 0.6991 for ResNet-50. For DeiT-Small on Cheng, Telea removal of the tumour core yielded a mean sham-corrected sensitivity of 0.1282 (slice-level Wilcoxon $p = 9.13\times10^{-6}$; condition-level Holm-adjusted $p = 3.65\times10^{-5}$; patient-clustered two-sided Wilcoxon $W = 8.0$, $p = 0.0488$ [one-sided $W = 47.0$, $p = 0.0244$] across $N = 10$ patient clusters); the corresponding ResNet-50 value was 0.0311 (condition-level Holm-adjusted $p = 2.62\times10^{-3}$; patient-clustered $p = 0.0488$). On BRISC, DeiT-Small core sensitivity was 0.0517 (raw two-sided $p = 0.0320$; condition-level Holm-adjusted $p = 0.0639$; 32-test family Holm-adjusted $p = 0.6072$, not significant after family-wise error rate control under either strategy). In the pooled comparison across 55 clusters, DeiT-Small showed greater core reliance than ResNet-50 (difference +0.0738; slice-level $p = 0.0123$; cluster-level Wilcoxon $W = 483.0$, $p = 0.0162$; paired t-test $t(54) = 3.29$, $p = 0.00177$; Linear

Mixed Model fixed effect $\beta = +0.074$, SE = 0.020, $z = 3.756$, $p < 0.001$). Remote-region effects were not statistically distinguishable from sham in primary comparisons. In the evaluated BRISC archive, 159 exact duplicate pairs and 101 cross-split test matches were identified; this archive-specific finding was addressed by evaluating both official (N = 1,000) and clean (N = 899) partitions. These experiments provide a retrospective mechanistic audit of spatial reliance in single-run baseline models, not clinical validation or evidence of diagnostic deployability.

Keywords: brain tumour MRI, counterfactual intervention, peritumoral margin, shortcut learning, ResNet-50, DeiT-Small, Vision Transformer, data leakage, calibration

1 Introduction

Automated classification of primary intracranial neoplasms from contrast-enhanced magnetic resonance imaging (CE-MRI) is a foundational task in computer-aided neuro-oncology. Discriminating between the three most prevalent primary brain tumours (meningiomas, gliomas and pituitary adenomas) directly governs neurosurgical approach selection, stereotactic radiosurgery boundaries and adjuvant chemotherapy regimens. Over the past decade, deep learning systems based on convolutional neural networks (CNNs) and Vision Transformers (ViTs) have repeatedly reported benchmark classification accuracies exceeding 90% to 95% on public MRI datasets [2, 5].

Despite these nominal scores, translational deployment of medical imaging AI into clinical radiology workflows remains impeded by vulnerability to shortcut learning and the "Clever Hans" effect [1, 11]. Wallis and Buvat [1] showed that deep classifiers trained on a widely used public brain tumour MRI dataset achieved high diagnostic accuracy even when the entire intracranial brain compartment was occluded. The models were inadvertently exploiting acquisition-site background noise, skull thickness signatures and scanner artefacts rather than pathological tissue. Furthermore, unaddressed patient-level data leakage and slice-to-slice correlation across cross-validation splits distort reported metrics and create a false impression of clinical generalisation [6]. Interpretability claims for high-stakes imaging models also need to be tested rather than assumed [13].

Existing methodologies for auditing spatial reliance suffer from two main limitations:

1. **Coarse binary masking and out-of-distribution shift.** Prior occlusion studies predominantly replace entire cerebral hemispheres or rectangular tiles with zero-intensity black patches or uniform noise [3, 4]. This introduces sharp, high-frequency boundary cliffs and out-of-distribution (OOD) domain shift, making it impossible to determine whether a performance collapse is caused by the loss of diagnostic evidence or by network failure on unnatural artefacts.
2. **Absence of anatomically matched controls.** Standard ablation studies lack matched sham controls. Without assessing the effect of identical image perturbations in normal contralateral brain parenchyma, one cannot isolate tumour-specific diagnostic evidence from non-specific perturbation sensitivity.

Research objectives and hypotheses. To address these challenges, we design and execute a literature-informed, sham-controlled counterfactual audit with descriptive plane stratification across CNN (ResNet-50) and Vision Transformer (DeiT-Small/16) architectures (Fig. 1). The study is a mechanistic audit of spatial reasoning in two research-grade classifiers; it is not a diagnostic tool and makes no claim of clinical deployability. We investigate five hypotheses:

- **Hypothesis 1 (spatial evidence gradient).** Lesion-centred perturbations are expected to produce larger sham-corrected changes than remote-region perturbations; the magnitude and ordering may vary by cohort, model and perturbation operator.
- **Hypothesis 2 (transitional margin contribution).** The immediate 4-pixel margin may contribute predictive evidence beyond the matched sham response; this is tested separately by cohort and with multiplicity correction.

- **Hypothesis 3 (architectural inductive bias).** ResNet-50 and DeiT-Small may exhibit different regional sensitivity and calibration profiles; the direction and robustness of any difference are empirical questions.
- **Hypothesis 4 (perturbation method concordance).** Regional sensitivity estimates may be concordant across Telea inpainting and locally matched texture filling; this is evaluated rather than assumed.
- **Hypothesis 5 (benchmark leakage and prior distortion).** The locally evaluated BRISC archive may contain exact duplicate matches across partitions; any finding is specific to the inspected archive and is not presumed to apply to every distribution.

2 Related work

2.1 Shortcut learning in medical image analysis

Shortcut learning occurs when machine learning models achieve high nominal accuracy by exploiting spurious decision rules that fail under distributional shift [11]. In radiographic COVID-19 detection, DeGrave et al. [14] demonstrated that deep networks relied heavily on hospital-specific radiopaque markers and scanner-bed textures rather than pulmonary consolidations. In neuro-oncology, Wallis and Buvat [1] showed that skull and background features alone supported high classification accuracy, so benchmark performance on public MRI datasets cannot be accepted without region-resolved attribution audits.

2.2 Spatial interventions and counterfactual attribution

Attribution by input perturbation has evolved from simple occlusion sensitivity to generative counterfactuals. Goyal et al. [12] formalised counterfactual visual explanations by identifying minimal edits required to alter class predictions. Rossi et al. [3] introduced TACE (Tumour-Aware Counterfactual Explanations) as a preprint framework to evaluate classifier behaviour under generative lesion insertion and removal. Sourget et al. [4] investigated model sensitivity to unexpected anatomical regions in radiographic images. Among the studies we reviewed, prior medical imaging intervention studies evaluated CNNs rather than Vision Transformers and did not use matched contralateral controls to cancel generic inpainting artefacts.

2.3 Brain-tumour MRI classification and benchmark leakage audits

The brain tumour dataset published by Cheng et al. [2] ($N = 3{,}064$ slices) has served as a foundational benchmark for over a decade. Recently, Fateh et al. [5] introduced the BRISC benchmark ($N = 6{,}000$ multi-plane slices). Public medical datasets frequently suffer from duplicate contamination and patient split leakage. As demonstrated by Yagis et al. [6], random slice-level cross-validation without patient grouping results in near-perfect slice memorisation and inflated performance estimates.

3 Datasets, provenance and data leakage audit

3.1 Datasets and cohort overview

We evaluated two contrast-enhanced T1-weighted brain MRI datasets:

1. **The Cheng dataset [2].** 3,064 CE-T1 slices from 233 unique patients across three tumour classes: 708 meningiomas, 1,426 gliomas and 930 pituitary adenomas. Each slice contains radiologist-delineated tumour boundary contours.
2. **The BRISC dataset [5].** 6,000 skull-stripped CE-MRI slices partitioned into an official training set ($N = 5{,}000$, split into 4,000 train and 1,000 validation) and an official test set ($N = 1{,}000$) across four classes (glioma, meningioma, pituitary tumour and no-tumour controls), imaged in axial, coronal and sagittal planes. Usable segmentation masks are available for 860 test neoplasm slices.

3.2 Sample accounting and selection rules

To ensure transparency and reproducibility across all reported tables and figures, the sample flow and selection rules are as follows (Fig. 1).




Fig. 1 Study design and sham-controlled counterfactual intervention pipeline: (A) ingestion and provenance audit of the Cheng ($N = 3{,}064$) and BRISC ($N = 6{,}000$) cohorts; (B) five-region topological decomposition; (C) dual perturbation engines; (D) paired model evaluation across ResNet-50 and DeiT-Small.

- **Baseline evaluation partitions.**
  - For Cheng (N = 3,064 slices across 233 patients), a five-fold patient-isolated cross-validation split was created. The saved prediction files in predictions/ (oof_predictions_resnet50_cheng.csv and oof_predictions_deit_small_patch16_224_cheng.csv) contain exactly $N = 480$ records, all belonging to Fold 1 (evaluated on 49 unseen patients). Slices from folds 2–5 are not included in these exported prediction logs; this partition is therefore designated the **patient-isolated held-out Fold 1 test partition ($N = 480$)** rather than an aggregated all-fold out-of-fold set.
  - For BRISC, the complete official test predictions (N = 1,000) are exported in predictions/test_predictions_resnet50_brisc.csv and predictions/test_predictions_deit_small_patch16_224_brisc.csv, while the clean de-duplicated predictions (N = 899) are exported in the corresponding *_brisc_clean.csv files. The mask-supported neoplasm evaluation subset (N = 480; 254 gliomas, 226 meningiomas) is preserved in predictions/test_predictions_*_neoplasm_480.csv.
- **Counterfactual intervention QC manifest ($N = 1{,}200$).** Phase 6 candidate pool, recorded in manifests/intervention_qc.csv across 120 unique slices (60 Cheng, 60 BRISC), five regions (R_{core}, $R_{\text{immediate}}$, R_{outer}, R_{remote}, R_{sham}) and two perturbation engines (Method A: Telea inpainting; Method B: Gaussian texture fill), giving 120 × 5 × 2 = 1,200 verified records. All 90 evaluated slices (45 Cheng + 45 BRISC) have complete, verified QC records in this manifest, confirming zero non-target corruption ($|\Delta| < 10^{-5}$). The remaining 30 candidate slices were not part of the balanced evaluation protocol below, which retains 15 slices per tumour class in Cheng and 15 per acquisition plane in BRISC.
- Paired counterfactual observations (N = 1,800) and sample accounting. Two distinct sample sets were constructed: (i) an Image Quality Control (QC) pool of 120 unique slices (60 Cheng, 60 BRISC; 1,200 records in manifests/intervention_qc.csv) to verify 100% mathematical non-target pixel invariance ($|\Delta| < 10^{-5}$ outside edited masks); and (ii) a primary Model Evaluation pool of 90 balanced slices (45 Cheng, 45 BRISC; 1,800 paired predictions in predictions/intervention_effects.csv across 2 models × 5 regions × 2 methods). Slices for model evaluation were selected deterministically in sequential index order of the test manifests, taking the first 15 mask-containing slices meeting technical invariance per tumour class for Cheng (15 gliomas, 15 meningiomas, 15 pituitary adenomas, originating from N = 10 patient clusters) and the first 5 slices per (class × plane) cell for BRISC (5 glioma + 5

meningioma + 5 pituitary across axial, coronal, and sagittal planes = exactly 15 slices per plane). No post-hoc cherry-picking or manual reordering was applied.

3.3 Duplicate audit and split policy

- **Cheng patient-isolated five-fold cross-validation.** Embedded MATLAB struct headers were parsed to extract patient identifiers (PID). A five-fold split was established using StratifiedGroupKFold (n_splits = 5, shuffle = True, random_state = 42) on patient ID. Automated set intersection verified zero patient overlap across folds (intersection = ∅).
- **BRISC duplicate discovery and provenance.** In the Kaggle development distribution of BRISC used here (manifests/duplicate_pairs.csv), perceptual hashing (pHash, dHash, aHash) and decoded SHA-256 pixel-matrix verification identified 159 exact duplicate image pairs (identical decoded SHA-256 pixel matrices). Of these, 110 pairs cross the official train–test split boundary, involving 101 unique test slices (10.1% of the 1,000-slice test set) that are identical copies of training/validation images (96 meningiomas and 5 pituitary adenomas). However, the peer-reviewed data descriptor by Fateh et al. [5] (*Scientific Data*, 2026) reports that de-duplication was conducted before dataset partitioning. We therefore present this finding as an empirical observation specific to the archive release we evaluated, not as a verdict on the published benchmark design. The clean subset serves as a sensitivity check on the official results. We report both the official benchmark (N = 1,000) and the local de-duplicated clean subset (N = 899; 759 mask-supported clean slices) side by side.
- **Cross-dataset overlap.** De-duplication screening between Cheng (3,064 slices) and BRISC (6,000 slices) found 0 exact duplicate pairs and structural similarity index (SSIM) values between 0.28 and 0.58. Exact hashing and SSIM cannot exclude perceptual near-duplicates, particularly after skull-stripping and resizing, and BRISC was collated from public sources [5]; the cohorts are therefore treated as separate collections whose independence is not proven.

4 Materials and methods

4.1 Image preprocessing and geometric standardisation

All slices were standardised through an audited pipeline (Fig. 2).




Fig. 2 Preprocessing pipeline and mask geometric alignment verification: raw slice, percentile intensity scaling [P_1, P_{99}], aspect-ratio-preserving letterbox padding (224 × 224) and binary mask boundary alignment.

1. Percentile intensity clamping. Tissue intensities were mapped to [0, 1]:

Contralateral sham controls were constructed via vertical midline reflection into the contralateral normal brain parenchyma, followed by affine in-tissue translation wherever mirrored contours overlapped the peritumoral exclusion zone or intracranial tissue boundaries. Across all 120 evaluated QC slices, 100% of sham interventions (120/120) utilized the patient's lesion-derived mirrored contour template (22.5% direct vertical reflection, 77.5% affine translated contour; 0% circular fallback). Across all 120 slices, 95.8% (115/120) achieved exact 1.0000 geometric parity with the lesion core, while 4.2% (5/120) experienced peripheral clipping against intracranial tissue masks

(minimum area ratio 0.7525), yielding an overall mean sham-to-core area ratio of 0.9975 ± 0.0227 and mean template Dice overlap of 0.9986 ± 0.0130.

where P_1 and P_{99} denote the 1st and 99th percentiles of intracranial tissue.

2. Aspect-ratio-preserving letterbox padding. Slices were scaled along their maximum axis to 224 pixels and symmetrically padded with zeros to 224 × 224. Nearest-neighbour interpolation was enforced for segmentation masks. Across all 3,064 Cheng and 860 BRISC test slices, 0 masks disappeared, area retention matched theoretical scaling, and centroid drift was bounded within 0.0035.

3. Channel standardisation. Slices were replicated to three channels and normalised using ImageNet parameters (μ = [0.485, 0.456, 0.406], σ = [0.229, 0.224, 0.225]). Horizontal and vertical flips were omitted to preserve hemispheric anatomical orientation.

4.2 Model architectures and training protocols

Models were configured in strict concordance with configs/training.yaml.

- **ResNet-50 (CNN).** 23.51 million parameters; initialised with IMAGENET1K_V2 weights [7]; linear classification head; dropout $p = 0.2$; initial learning rate 1×10^{-4}.
- **DeiT-Small/16 (ViT).** 21.66 million parameters; initialised with ImageNet-1k weights [8]; 12 Transformer blocks, 384 embedding dimensions, 6 heads; class-token projection head; dropout $p = 0.1$; initial learning rate 5×10^{-5}.
- **Optimisation.** Both architectures were optimised using AdamW with weight decay 1×10^{-2} and batch size 16. Training ran for up to 5 epochs per fold/split with early stopping (patience of 3 epochs) on validation macro-F1, following a one-epoch linear warm-up and cosine annealing schedule. Standard cross-entropy loss was minimised with random seed 42. The baselines are therefore single-run, short-schedule fine-tuned models, and the Cheng results use Fold 1 only.
- **Computational environment.** All experiments used Python 3.10.11, PyTorch 2.5.1 and CUDA 12.1 on an NVIDIA RTX 4050 GPU (6 GB VRAM). Hardware, operating system, software versions, git revision and SHA-256 hashes of the model checkpoints are recorded in environment_manifest.json.

4.3 Five-region topological decomposition

For each test slice x with tumour mask M and intracranial tissue support S, five regions were defined:

1. Tumour core (R_{core}). The lesion mask M.

2. Immediate peritumoral margin ($R_{\text{immediate}}$). A 4-pixel dilation ring:

$$R_{\text{immediate}} = (\text{dilate}(M, 4) - M) \cap S \tag{2}$$

3. Outer peritumoral margin (R_{outer}). A 12-pixel dilation ring:

$$R_{\text{outer}} = (\text{dilate}(M, 16) - \text{dilate}(M, 4)) \cap S \tag{3}$$

4. Remote context (R_{remote}). A tissue patch located more than 32 pixels from the tumour boundary within S.

5. Matched contralateral sham control (R_{sham}). An area- and contour-matched control placed in contralateral normal parenchyma and disjoint from the lesion-centred regions:

$$R_{\text{sham}} \cap (R_{\text{core}} \cup R_{\text{immediate}} \cup R_{\text{outer}}) = \emptyset \tag{4}$$

4.4 Dual counterfactual perturbation engines

- **Method A (Telea PDE inpainting).** Solves the fast-marching-method boundary problem (cv2.INPAINT_TELEA, radius 3 px) [10] to propagate surrounding healthy parenchyma inward.

- **Method B (locally matched Gaussian texture fill).** Estimates the local surrounding mean μ_{local} and variance σ_{local}^2 from a 5-pixel dilated border, and fills the region with Normal(μ_{local}, σ_{local}) smoothed via a 2-pixel cosine border taper. The fill therefore matches the local first- and second-order intensity statistics of the surroundings but not their spatial texture structure, and it is treated as a complementary probe rather than a substitute for inpainting.
- **Non-target pixel invariance.** Both implementations preserve original pixel values outside the edited mask; the maximum absolute difference was checked in the intervention QC records:

$$x_{edit}(u, v) = x(u, v) \text{ for all } (u, v) \notin R \tag{5}$$

4.5 Sensitivity estimands and statistical inference

Raw evidence drop (Δ_r), **sham-corrected region sensitivity** (τ_r) and **top-1 prediction flip** (F_r) are defined as

$$\Delta_r(x) = p_y(x) - p_y(x_{edit,r}) \tag{6}$$

$$\tau_r(x) = \Delta_r(x) - \Delta_{sham}(x) \tag{7}$$

$$F_r(x) = 1 \text{ if } \operatorname{argmax} p(x_{edit,r}) \neq \operatorname{argmax} p(x), \text{ and 0 otherwise} \tag{8}$$

where p_y is the predicted probability of the ground-truth class.

- **Hypothesis testing.** Two-sided paired Wilcoxon signed-rank tests assess whether region sensitivity differs from the sham control; Holm correction is applied across regions.
- **Patient-clustered analysis.** For the Cheng cohort, where 45 slices originate from 10 patients, patient-level mean sensitivities were computed and evaluated with clustered Wilcoxon and clustered one-sample *t*-tests. For the pooled multi-cohort sample ($N = 90$), clustered inference across 55 independent clusters (10 Cheng patients + 45 BRISC cases) was conducted alongside linear mixed-effects modelling.
- **Effect size.** Cohen's d_z:

$$d_z = \text{mean}(\tau_r) / \text{SD}(\tau_r) \tag{9}$$

5 Results

5.1 Baseline classification performance and probability calibration

Baseline models showed moderate-to-good discrimination on both benchmarks, with DeiT-Small clearly stronger than ResNet-50 (Table 1; Fig. 3). Both are short single-run baselines and not tuned to benchmark-level performance. DeiT-Small outperformed ResNet-50 in macro-F1 (Cheng Fold 1, $N = 480$: 0.8403 vs 0.7190; BRISC clean four-class test, $N = 899$: 0.8344 vs 0.6991) and in probability calibration. On Cheng Fold 1, DeiT-Small achieved a 10-bin expected calibration error (ECE) [9] of 0.0366, compared with 0.1835 for ResNet-50.




Fig. 3 Reliability diagrams and calibration curves: empirical calibration curves and 10-bin expected calibration error (ECE) for ResNet-50 and DeiT-Small on the Cheng patient-isolated held-out Fold 1 partition and the BRISC clean test set.

Table 1 Baseline classification performance and probability calibration. Panel A: full-label evaluation (Cheng, three classes, Fold 1 held-out partition; BRISC, four classes). Panel B: BRISC mask-supported neoplasm batch, in which only glioma and meningioma are present; macro-F1 is computed over the two present labels.

Dataset	Model	Evaluation partition	Accuracy	Balanced acc.	Macro-F1	ECE (10-bin)	Brier score
Panel A. Full-label evaluation (Cheng, three classes; BRISC, four classes)							
Cheng	ResNet-50 (CNN)	Patient-isolated held-out Fold 1 ($N = 480$)	0.7583	0.7448	0.7190	0.1835	0.3692
Cheng	DeiT-Small (ViT)	Patient-isolated held-out Fold 1 ($N = 480$)	0.8458	0.8394	0.8403	0.0366	0.2136
BRISC	ResNet-50 (CNN)	Official test ($N = 1{,}000$; full 4-class)	0.7040	0.6870	0.7095	0.3875	0.6510
BRISC	ResNet-50 (CNN)	Clean test ($N = 899$; de-duplicated)	0.6963	0.6815	0.6991	0.3790	0.6505
BRISC	DeiT-Small (ViT)	Official test ($N = 1{,}000$; full 4-class)	0.7880	0.8094	0.8030	0.0719	0.2864
BRISC	DeiT-Small (ViT)	Clean test ($N = 899$; de-duplicated)	0.8265	0.8330	0.8344	0.0901	0.2448
Panel B. BRISC mask-supported neoplasm batch (two classes present; macro-F1 over present labels)							
BRISC	ResNet-50 (CNN)	Official neoplasm batch ($N = 480$; glioma and meningioma)	0.6479	0.6510	0.6919	0.3312	0.6591
BRISC	ResNet-50 (CNN)	Clean neoplasm batch ($N = 397$)	0.6247	0.6349	0.6574	0.3061	0.6595
BRISC	DeiT-Small (ViT)	Official neoplasm batch ($N = 480$; glioma and meningioma)	0.6875	0.6813	0.7131	0.0421	0.3938
BRISC	DeiT-Small (ViT)	Clean neoplasm batch ($N = 397$)	0.7456	0.7294	0.7431	0.0718	0.3303

Note: For Cheng, evaluations reflect the 480 held-out slices of Fold 1 (49 patients unseen in training), not pooled five-fold out-of-fold predictions. For BRISC, official and clean benchmark scores are computed across the full four-class test partitions ($N = 1{,}000$ and $N = 899$; test_predictions_*_brisc.csv and test_predictions_*_brisc_clean.csv). Panel B reports the mask-supported neoplasm subset ($N = 480$ official and $N = 397$ clean; test_predictions_*_neoplasm_480.csv) with the label set fixed to the classes present in each subset.

5.2 Quality control of counterfactual interventions

All 1,200 generated counterfactual interventions met the quality-control gates (Fig. 4):




Fig. 4 Counterfactual intervention and residual montage gallery: an eight-panel diagnostic display showing the original slice, the five-colour segmentation topology, Method A inpainted core and sham, Method B texture-filled core and sham, and residual difference maps.

- **Non-target preservation.** The maximum absolute difference outside the edit mask was 0.00 across all 1,200 edits.
- **Intra-mask magnitude.** The mean absolute difference (MAD) inside the core mask averaged 0.156–0.232 (Method A) and 0.216–0.259 (Method B).
- **Evaluated coverage.** All 90 evaluated slices in predictions/intervention_effects.csv have verified QC records in manifests/intervention_qc.csv.
- **Edge discontinuity.** Boundary gradient offsets were bounded (0.16–0.24); this measure does not imply that the edited regions are artefact-free.

5.3 Primary counterfactual sensitivity: core versus margin evidence

Evaluation of the 1,800 paired counterfactual observations in predictions/intervention_effects.csv revealed spatial sensitivities that differ across architectures and intervention methods (Table 2; Fig. 5).




Fig. 5 Regional evidence sensitivity across architectures (Method A, inpainting), pooled across the Cheng and BRISC intervention subsets (N = 90 slices): sham-corrected sensitivity (τ_r) for the tumour core, immediate

margin (4 px), outer margin (12 px) and remote context. Error bars show the standard error of the mean; asterisks mark paired Wilcoxon significance against sham (* $p < 0.05$, *** $p < 0.001$; n.s., not significant).

Table 2 Regional evidence sensitivity and hypothesis testing. Shown are 12 of the 40 dataset × model × method × region estimates: all four regions for DeiT-Small with Method A in each cohort, plus core sensitivity (and the immediate margin for ResNet-50 Method A) for the remaining Cheng model–method combinations. Values for Δ_r and τ_r are mean ± SD. Holm adjustment is applied across the four regions within each dataset–model–method set.

Dataset	Model	Method	Region	Raw sensitivity (Δ_r)	Sham-corrected (τ_r)	Flip rate	Raw Wilcoxon p	Holm-adj. p	Cohen's d_z
Cheng (N = 45)	DeiT-Small	A (Telea)	Core	0.1212 ± 0.245	0.1282 ± 0.245	13.3%	9.13×10^{-6}	3.65×10^{-5}	0.518
Cheng (N = 45)	DeiT-Small	B (Texture)	Core	0.1577 ± 0.241	0.1449 ± 0.214	8.9%	3.09×10^{-8}	1.24×10^{-7}	0.669
Cheng (N = 45)	DeiT-Small	A (Telea)	Immediate (4 px)	0.0143 ± 0.082	0.0213 ± 0.096	8.9%	0.0267	0.0802	0.222
Cheng (N = 45)	DeiT-Small	A (Telea)	Outer (12 px)	−0.0007 ± 0.091	0.0063 ± 0.098	8.9%	0.2866	0.2866	0.064 (n.s.)
Cheng (N = 45)	DeiT-Small	A (Telea)	Remote context	0.0058 ± 0.030	0.0128 ± 0.065	2.2%	0.0801	0.1603	0.196 (n.s.)
Cheng (N = 45)	ResNet-50	A (Telea)	Core	0.0264 ± 0.041	0.0311 ± 0.056	8.9%	6.54×10^{-4}	2.62×10^{-3}	0.548
Cheng (N = 45)	ResNet-50	B (Texture)	Core	0.0844 ± 0.134	0.0603 ± 0.150	31.1%	0.0173	0.0691	0.396
Cheng (N = 45)	ResNet-50	A (Telea)	Immediate (4 px)	0.0079 ± 0.036	0.0125 ± 0.050	6.7%	0.0621	0.1242	0.248
BRISC (N = 45)	DeiT-Small	A (Telea)	Core	0.0367 ± 0.096	0.0517 ± 0.120	8.9%	0.0320	0.0639 [0.6072]*	0.432
BRISC (N = 45)	DeiT-Small	A (Telea)	Immediate (4 px)	0.0089 ± 0.048	0.0239 ± 0.063	6.7%	0.0152	0.0455	0.381
BRISC (N = 45)	DeiT-Small	A (Telea)	Outer (12 px)	0.0150 ± 0.065	0.0301 ± 0.071	11.1%	0.0049	0.0197	0.425
BRISC (N = 45)	DeiT-Small	A (Telea)	Remote context	−0.0010 ± 0.018	0.0140 ± 0.056	6.7%	0.2815	0.2815	0.249 (n.s.)

*Note: Raw Wilcoxon two-sided p-values with step-down Holm-Bonferroni correction within each 4-region hypothesis family. *For BRISC DeiT Core, condition-level Holm adjustment yields p = 0.0639 (Rank 3/4, multiplier 2); under the complete 32-test study-wide family (Table S1), p = 0.6072; both confirm non-significance at α = 0.05. n.s.: not significant.*

Non-monotonic spatial dynamics and patient-clustered analysis.

1. **Core evidence dominance versus method dependency.** Core dominance is strongly evident in Vision Transformers under PDE inpainting on Cheng, where core sensitivity ($\tau_{core} = 0.1282$) far exceeds margin effects ($\tau_{immediate} = 0.0213$, $\tau_{outer} = 0.0063$, $\tau_{remote} = 0.0128$). However, spatial decay is not universally monotonic: under Method B (locally matched Gaussian texture fill) and in skull-stripped BRISC scans, outer-margin effects frequently equal or approach core sensitivity (e.g. BRISC DeiT outer $\tau = 0.0301$ vs core $\tau = 0.0517$). The outer ring covers more tissue than the immediate ring and may include perilesional signal and boundary texture. The experiment does not isolate oedema as the cause of measured sensitivity.
2. Patient-clustered statistical inference. In Cheng, the 45 intervention slices originate from N = 10 unique patient clusters. While slice-level Wilcoxon tests yield $p = 9.13 \times 10^{-6}$ for DeiT-Small and $p = 6.54 \times 10^{-4}$ for ResNet-50, patient-clustered analysis across the 10 patient-mean sensitivities yields clustered Wilcoxon W = 8.0, two-sided p = 0.0488 (one-sided W = 47.0, p = 0.0244) for both architectures (clustered one-sample t-test: DeiT t = 2.22, p = 0.0534, ResNet t = 2.00, p = 0.0771). Both levels of inference confirm core sensitivity above sham controls, but patient clustering reveals the true effective degrees of freedom (N = 10).
3. **Remote context invariance.** Remote-region effects were non-significant in the highlighted comparisons (BRISC DeiT Telea $p = 0.2815$; Cheng DeiT Telea $p = 0.0801$ before multiplicity correction). Under Method B, which fills the region with noise matched to the local mean and variance of its surroundings, DeiT-Small remote sensitivity was likewise low and non-

significant (τ_{remote} = 0.0052, p = 0.4571), whereas ResNet-50 showed higher remote-parenchyma sensitivity under the same perturbation. Non-significance is not proof of zero effect or of complete shortcut independence.

4. **Region area and per-pixel sensitivity.** Regions differ in size and only the sham is area-matched to the core, so differences between regions may partly reflect area. The outer margin was on average 2.2 times larger than the tumour core (mean 2,239 vs 1,013 px). Normalised per 1,000 edited pixels, sensitivity ranged from 0.049 to 0.136 for the core and from 0.015 to 0.036 for margin and remote regions, so evidence density was higher in the core in the summarised comparisons.

5.4 Architectural inductive bias: ViT versus CNN

Paired comparison between DeiT-Small and ResNet-50 showed architectural divergences (Fig. 6).




Fig. 6 Architectural inductive bias, ViT versus CNN regional sensitivity profile (Method A): difference plot ($\tau_{ViT} - \tau_{CNN}$) highlighting the greater reliance of DeiT-Small on intra-lesion core and transitional margin evidence.

- Intra-lesion core reliance. In the pooled multi-cohort comparison (N = 90, Method A), DeiT-Small showed higher mean sensitivity to core ablations than ResNet-50 (τcore,ViT = 0.0900 vs τcore,CNN = 0.0161; difference +0.0738; slice-level paired Wilcoxon p = 0.0123; Cohen's dz = 0.362). Crucially, under cluster-level inference accounting for the 55 independent clusters (10 Cheng patient clusters + 45 independent BRISC cases), the architectural superiority of Vision Transformers remains robust: cluster-level Wilcoxon W = 483.0, two-sided p = 0.0162, cluster-level paired t-test t(54) = 3.29, p = 0.00177, and Linear Mixed-Effects Modeling fixed effect βViT = +0.074 (SE = 0.020, Wald z = 3.756, p < 0.001). In the Cheng-specific subset (N = 45), the core sensitivity difference was 0.1282 − 0.0311 = +0.0971 (slice-level Wilcoxon p = 0.156; patient-clustered Wilcoxon p = 0.232, clustered t-test p = 0.148). Statistical significance for architectural divergence is thus achieved when pooling across the combined multi-cohort (N = 90, 55 clusters).
- **Transitional margin reliance.** DeiT-Small displayed moderately higher sensitivity to the immediate peritumoral margin across the pooled cohort (τ = 0.0270 vs 0.0104; difference +0.0166; p = 0.1124) and to the outer margin (difference +0.0118).
- **Remote-region comparison.** The pooled difference between architectures for remote-region sensitivity was small (difference +0.0072; p = 0.9087). This audit did not show a detectable architectural difference; it does not prove that either model is free of background shortcuts.

5.5 Perturbation method concordance

The two perturbation engines did not produce uniformly concordant effects (Fig. 7). The correlation between Telea and texture-fill core sensitivities was $r = 0.748$ (Spearman $\rho = 0.541$) when pooled across models on Cheng, and $r = 0.300$ ($\rho = 0.332$) on BRISC. Across the region-by-dataset comparisons, correlations varied substantially, including weak or near-zero associations for several margin and remote regions. This indicates that measured sensitivity is partly intervention-dependent. Texture filling should therefore be interpreted as a complementary perturbation, not as an interchangeable substitute for inpainting. The results do not establish that one architecture is intrinsically more sensitive to a particular artefact mechanism.




Fig. 7 Perturbation method concordance, Telea inpainting versus matched texture fill: paired sensitivity comparison between Telea boundary inpainting and locally matched Gaussian texture filling across ResNet-50 and DeiT-Small.

5.6 Imaging plane stratification (BRISC)

Regional sensitivity varied across acquisition planes (Table 3; Fig. 8). Tumour-core sensitivity was highest in each plane, but the ordering of the surrounding regions was not uniform. In particular, axial remote-context sensitivity exceeded immediate-margin sensitivity (0.0312 vs 0.0283), yielding the axial ordering: Core (0.0492) > Outer margin (0.0469) > Remote context (0.0312) > Immediate margin (0.0283). In coronal slices, ordering was Core (0.0370) > Outer (0.0175) > Immediate (0.0118) > Remote (0.0027); and in sagittal slices, Core (0.0690) > Immediate (0.0317) > Outer (0.0259) > Remote (0.0082). The unweighted mean across the three planes exactly reproduces the overall BRISC DeiT inpainting sensitivities reported in Table 2: Core ((0.0492 + 0.0370 + 0.0690)/3 = 0.0517), Immediate margin ((0.0283 + 0.0118 + 0.0317)/3 = 0.0239), Outer margin ((0.0469 + 0.0175 + 0.0259)/3 = 0.0301), and Remote context ((0.0312 + 0.0027 + 0.0082)/3 = 0.0140). These plane-stratified findings are descriptive and do not establish statistically significant differences between acquisition planes.




Fig. 8 Core sensitivity by acquisition plane (BRISC, DeiT-Small, Method A): axial ($N = 15$), coronal ($N = 15$) and sagittal ($N = 15$).

Table 3 Core sensitivity by acquisition plane (BRISC, DeiT-Small, Method A).

Acquisition plane	Sample count (N)	Mean core τcore	Mean immediate margin τimmed	Mean outer margin τouter	Mean remote τremote
Axial	15	0.0492	0.0283	0.0469	0.0312
Coronal	15	0.0370	0.0118	0.0175	0.0027
Sagittal	15	0.0690	0.0317	0.0259	0.0082

Note: Descriptive sample; no inferential test of plane differences is claimed. The mean of the three plane values (0.0517) matches Table 2.

5.7 Exploratory tumour class heterogeneity

Analysing sensitivity across Cheng tumour classes revealed marked differences in regional reliance (Table 4; Fig. 9). In Cheng (DeiT-Small, Method A; N = 15 per class from 10 patient clusters), Meningiomas demonstrated pronounced core sensitivity (τ_core = 0.3438) and modest margin sensitivity (τ_immed = 0.0338, ratio 9.8%). Gliomas exhibited τ_core = 0.0398 and τ_immed = 0.0268 (ratio 67.2%), while Pituitary adenomas showed circumscribed presentation (τ_core = 0.0009, τ_immed = 0.0034, τ_outer = 0.0392). The unweighted class mean sensitivity across the 45 Cheng slices reproduces Table 2 exactly: Core ((0.0398 + 0.3438 + 0.0009)/3 = 0.1282), Immediate margin ((0.0268 + 0.0338 + 0.0034)/3 = 0.0213), Outer margin ((-0.0066 - 0.0136 + 0.0392)/3 = 0.0063), and Remote context ((0.0218 + 0.0165 + 0.0001)/3 = 0.0128). These exploratory results describe empirical model sensitivity patterns on 15 2D slices per class and should not be interpreted as direct evidence of histopathological infiltration or tumour-boundary biology.




Fig. 9 Core sensitivity by tumour class (Cheng, DeiT-Small, Method A): glioma, meningioma and pituitary adenoma ($N = 15$ each).

- Gliomas. A high margin-to-core sensitivity ratio (67.2%; τ_immed = 0.0268 vs τ_core = 0.0398), indicating substantial model reliance on transitional perilesional parenchyma in this exploratory 15-slice subset.
- Meningiomas. Pronounced core sensitivity (τ_core = 0.3438) with modest immediate margin sensitivity (9.8%; τ_immed = 0.0338), reflecting dominant sensitivity to internal lesion features.
- Pituitary adenomas. Minimal immediate margin sensitivity (τ_immed = 0.0034) with outer annular sensitivity (τ_outer = 0.0392), consistent with circumscribed intrasellar anatomy.

Table 4 Core sensitivity by tumour class (Cheng, DeiT-Small, Method A).

Tumour class	Sample count (N)	Core sensitivity (τcore)	Immediate margin (τimmed)	Outer margin (τouter)	Remote context (τremote)	Margin/Core ratio (%)
Glioma	15	0.0398	0.0268	-0.0066	0.0218	67.2%
Meningioma	15	0.3438	0.0338	-0.0136	0.0165	9.8%
Pituitary adenoma	15	0.0009	0.0034	0.0392	0.0001	360.2%

Note: Exploratory and descriptive; 15 slices per class from the 10 Cheng patient clusters, with no inferential test and no histopathological correlation. No radiological explanation for the class differences is tested here.

5.8 Impact of benchmark duplicate leakage

Comparing performance between the official benchmark ($N = 1{,}000$) and the de-duplicated clean test set ($N = 899$) showed that removing the 101 duplicate test images changed the measured performance in a model-dependent direction (Table 1; Fig. 10). On the four-class test set, DeiT-Small improved from 0.7880 to 0.8265 accuracy (+0.038) and from 0.8030 to 0.8344 macro-F1 (+0.031), whereas ResNet-50 changed little (accuracy 0.7040 to 0.6963; macro-F1 0.7095 to 0.6991). In the two-class neoplasm batch, with macro-F1 computed over the two present labels, DeiT-Small changed from 0.7131 to 0.7431 and ResNet-50 from 0.6919 to 0.6574. The 101 duplicate test images were heavily skewed towards meningiomas (96 of 101), so de-duplication also changed the test-set class balance. The data therefore do not support a uniform inflation of scores by duplicate leakage; the effect depends on the model and on the class composition of the subset.




Fig. 10 Impact of train–test duplicate leakage on benchmark evaluation (BRISC): official test set versus clean de-duplicated test set for DeiT-Small and ResNet-50. (A) Four-class evaluation (N = 1,000 vs 899). (B) Two-class neoplasm batch (N = 480 vs 397). Δ is clean minus official; axes start at zero.

5.9 Evaluation of the study hypotheses

- **Hypothesis 1 (spatial evidence gradient): partially supported.** Lesion-centred edits often produced larger effects than remote edits, but the ordering was not universal across all regions, datasets, architectures and perturbation engines.
- **Hypothesis 2 (margin contribution): partially supported.** The immediate-margin effect for DeiT-Small on BRISC remained significant after Holm adjustment; the corresponding Cheng result did not. Evidence therefore depends on cohort and multiplicity correction.
- **Hypothesis 3 (ViT inductive bias): partially supported; inconclusive for the Cheng-only comparison.** The pooled cluster-level comparison favoured DeiT-Small, but the slice-level Holm-adjusted p-value was 0.0980 and the Cheng-only architectural comparison was not significant in the patient-clustered analysis. DeiT-Small also had lower ECE on the reported baseline partitions, but baseline confidence and calibration differ between the models, which may influence probability-drop estimates.
- **Hypothesis 4 (perturbation method concordance): not uniformly supported.** Cross-method correlations were variable and weak for several regions; the two edit engines should be treated as complementary probes.
- **Hypothesis 5 (leakage and prior distortion): archive-specific finding, partly supported.** Exact duplicate matches were detected in the evaluated BRISC archive and changed the measured metrics in a model-dependent direction, but the result should not be generalised to every BRISC distribution or mirror.

6 Discussion

6.1 Principal findings

Our sham-controlled audit provides evidence that tumour-core perturbations reduce ground-truth-class probabilities in several model/cohort settings and that selected peritumoral edits can also affect predictions. Sham correction helps separate lesion-region effects from the response to an

anatomically separate control edit. The study is a mechanistic audit of spatial reliance, not a test of diagnostic deployability. The results are conditional on the perturbation operator and the evaluated subsets; non-significant remote-region tests do not establish the absence of all shortcut learning or scanner-related confounding.

6.2 Lesion core versus peritumoral margin evidence

The observed immediate-margin effects were modest and varied by cohort and method. For DeiT-Small under Telea inpainting, the mean sham-corrected immediate-margin sensitivity was 0.0213 on Cheng (raw $p = 0.0267$; Holm-adjusted $p = 0.0802$) and 0.0239 on BRISC (raw $p = 0.0094$; Holm-adjusted $p = 0.0188$). These findings are compatible with models using some information in tissue adjacent to the lesion, but do not by themselves identify the underlying radiological substrate or demonstrate histopathological infiltration.

6.3 Architectural inductive bias: attention versus convolution

The results are compatible with, but do not test, the following hypotheses about how architectural inductive biases may shape diagnostic representations:

1. **Possible receptive-field effect.** ResNet-50 showed lower sensitivity to core perturbations ($\tau \approx 0.031$ on Cheng, Method A) and greater sensitivity to edge artefacts under texture filling. Receptive-field constraints are one possible explanation, but this is untested: Δ is a drop in probability and depends on baseline confidence and calibration, which differ substantially between the models (ECE 0.04–0.09 for DeiT-Small vs 0.18–0.39 for ResNet-50).
2. **Possible global-context effect.** DeiT-Small showed higher core sensitivity ($\tau \approx 0.052$–0.128) and lower ECE (0.037 vs 0.184 on Cheng), with no detectable remote-context sensitivity under Method A ($\tau_{\text{remote}} \approx 0.013, p > 0.05$). Self-attention integration of spatial context is a hypothesis that these single-run baselines cannot confirm.

6.4 Benchmark leakage, provenance and data governance

The identical-pixel duplicate matches spanning the official train–test boundary in the evaluated BRISC archive highlight the need for data-governance checks. De-duplication changed the measured metrics in a model-dependent direction and altered the class composition of the test set (96 of 101 duplicated test images were meningiomas). We recommend transparent decoded-pixel and perceptual-hash audits, together with patient- or case-grouped splitting when reliable identifiers are available.

6.5 Limitations

Several limitations must be acknowledged:

1. **2D slices versus volumetric 3D MRI.** Our analysis evaluated 2D contrast-enhanced slices. Neuroradiologists assess multi-sequence 3D volumes (T1, T1+C, T2, FLAIR) in clinical practice.
2. **Synthetic perturbations.** The intervention code preserves pixels outside the target mask, and the repository reports zero measured non-target difference in its QC export. Nevertheless, Telea inpainting and Gaussian texture replacement create synthetic counterfactuals that may be out of distribution and do not simulate true healthy tissue, tumour resection or biological disease progression.
3. **Patient/case cluster sample size.** The Cheng intervention subset (45 slices) represents only 10 patients, and the BRISC audit subset represents 45 slice-level cases. Cluster-level inference is more conservative than treating all slices as independent, but the number of independent clusters remains limited, and case-level independence in BRISC follows the available sample identifiers rather than confirmed patient identities.

4. **Exploratory class inferences.** Class-specific analyses use 15 slices per tumour class in the Cheng intervention subset and 2D image masks/labels; they are descriptive and are not linked to independently reviewed histopathology or volumetric tumour boundaries.
5. **Single-run baselines.** Both models were fine-tuned for at most five epochs with one seed, and the Cheng analyses use Fold 1 only. The baselines (e.g. ResNet-50 accuracy 0.70 on BRISC four-class) are weaker than published benchmark results, so the audited models are not state-of-the-art classifiers and the architectural comparisons may not carry over to converged models.
6. **Region area and perturbation realism.** Only the sham is area-matched to the core, and Method B matches local intensity statistics but not spatial texture; per-pixel sensitivities are reported for context, not as a corrected estimate.

7 Conclusion

This study presents a sham-controlled, mechanistic counterfactual audit of ResNet-50 and DeiT-Small on two contrast-enhanced brain MRI cohorts. Lesion-core perturbations produced measurable reductions in predicted probability in several settings; immediate and outer-margin effects were more variable, and remote-region effects were generally smaller in the highlighted comparisons. DeiT-Small achieved higher baseline performance and lower ECE on the reported partitions, while its pooled core-sensitivity advantage requires qualified interpretation because slice-level Holm adjustment and clustered analyses answer different inferential questions. The archive-specific duplicate audit illustrates why benchmark provenance, patient/case grouping, intervention quality control and transparent sample accounting should accompany performance metrics. The study makes no claim of diagnostic deployability; external multi-centre and volumetric validation, and converged multi-seed baselines, remain necessary before any clinical use.

Statements and Declarations

Funding. This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

Competing interests. The authors declare that they have no competing financial or non-financial interests that could have appeared to influence the work reported in this paper.

Ethics approval and consent to participate. This study analysed publicly available, fully de-identified retrospective magnetic resonance imaging datasets (Cheng et al. [2] and BRISC [5]). The study protocol was reviewed and determined to be exempt from full institutional committee review by the Institutional Ethics Committee of Manipal University Jaipur under institutional research exemption reference MUJ/IEC/2026/04-EX (Category: Secondary Analysis of Anonymized Public Data; National Ethical Guidelines for Biomedical and Health Research, ICMR). In accordance with these regulations, individual patient informed consent was waived.

Consent for publication. Not applicable.

Data availability. The Cheng dataset is publicly accessible at Figshare (doi:10.6084/m9.figshare.1512427). The BRISC dataset is available through the dataset distribution described by Fateh et al. (doi:10.1038/s41597-026-06753-y). Audited manifests, split files, de-duplication mappings and prediction files are provided in the project repository; raw datasets are not redistributed, and SHA-256 hashes of the model checkpoints are recorded in environment_manifest.json.

Code and checkpoint availability. Source code, preprocessing scripts, model architectures, counterfactual intervention engines, evaluation scripts and validation test suites are open-source and hosted in the project repository: https://github.com/CSrajput-ux/brain-tumor-prediction-using-deep-leaning. Checkpoint SHA-256 hashes and hardware environment manifests are documented in revision/reports/environment_manifest.json and configs/checkpoint_manifest.json.

Generative AI and AI-assisted technologies. During manuscript preparation, the authors utilized large language model assistance (Google DeepMind Antigravity) for code development, data analysis verification, statistical integrity checking, figure rendering, and scientific language editing. The authors reviewed and edited all resulting content, performed all data checks, and take full responsibility for the content of the published article.

Dataset Licences: The Cheng dataset is distributed under the Creative Commons Attribution 4.0 International license (CC BY 4.0; Figshare doi:10.6084/m9.figshare.1512427). The BRISC dataset is distributed under the Creative Commons Attribution 4.0 International license (CC BY 4.0; Scientific Data doi:10.1038/s41597-026-06753-y).

Author contributions. Chhotu Rajput: conceptualisation, methodology, counterfactual intervention engine design, data auditing, experimental execution, statistical analysis, figure generation, writing – original draft. Rishi Gupta: supervision, project administration, methodology review, scientific validation, writing – review and editing. Both authors approved the final version.

References

[1] Wallis, D., Buvat, I.: Clever Hans effect found in a widely used brain tumour MRI dataset. Med. Image Anal. 77, 102368 (2022) https://doi.org/10.1016/j.media.2022.102368

[2] Cheng, J., Huang, W., Cao, S., et al.: Enhanced performance of brain tumor classification via tumor region augmentation and partition. PLOS ONE 10(10), e0140381 (2015) https://doi.org/10.1371/journal.pone.0140381

[3] Rossi, E.B., Lopez, E., Comminiello, D.: TACE: Tumor-aware counterfactual explanations for medical image analysis. arXiv preprint arXiv:2409.13045 (2024) https://doi.org/10.48550/arXiv.2409.13045

[4] Sourget, T., Hestbek-Møller, M., Jiménez-Sánchez, A., Xu, J.J., Cheplygina, V.: Mask of truth: model sensitivity to unexpected regions of medical images. J. Imaging Inform. Med. (2025) https://doi.org/10.1007/s10278-025-01531-5

[5] Fateh, A., et al.: BRISC: annotated dataset for brain tumor segmentation and classification. Sci. Data 13, 361 (2026) https://doi.org/10.1038/s41597-026-06753-y

[6] Yagis, E., et al.: Effect of data leakage in brain MRI classification using 2D convolutional neural networks. Sci. Rep. 11(1), 22544 (2021) https://doi.org/10.1038/s41598-021-01681-w

[7] He, K., Zhang, X., Ren, S., Sun, J.: Deep residual learning for image recognition. In: Proc. IEEE Conf. Computer Vision and Pattern Recognition (CVPR), pp. 770–778 (2016) https://doi.org/10.1109/CVPR.2016.90

[8] Touvron, H., Cord, M., Douze, M., et al.: Training data-efficient image transformers & distillation through attention. In: Proc. Int. Conf. Machine Learning (ICML), pp. 10347–10357 (2021)

[9] Guo, C., Pleiss, G., Sun, Y., Weinberger, K.Q.: On calibration of modern neural networks. In: Proc. Int. Conf. Machine Learning (ICML), pp. 1321–1330 (2017)

[10] Telea, A.: An image inpainting technique based on the fast marching method. J. Graph. Tools 9(1), 23–34 (2004) https://doi.org/10.1080/10867651.2004.10487596

[11] Geirhos, R., et al.: Shortcut learning in deep neural networks. Nat. Mach. Intell. 2(11), 665–673 (2020) https://doi.org/10.1038/s42256-020-00257-z

[12] Goyal, Y., Wu, Z., Ernst, J., Batra, D., Parikh, D., Lee, S.: Counterfactual visual explanations. In: Proc. 36th Int. Conf. Machine Learning (ICML), PMLR 97, pp. 2376–2384 (2019)

[13] Rudin, C.: Stop explaining black box machine learning models for high stakes decisions and use interpretable models instead. Nat. Mach. Intell. 1(5), 206–215 (2019) https://doi.org/10.1038/s42256-019-0048-x

[14] DeGrave, A.J., Janizek, J.D., Lee, S.-I.: AI for radiographic COVID-19 detection frequently relies on spurious shortcuts. Nat. Mach. Intell. 3(7), 610–619 (2021) https://doi.org/10.1038/s42256-021-00338-7

Supplementary Table S1 Complete 32-test inferential family across all dataset, architecture, intervention method, and anatomical region combinations.

Dataset & Model	Method	Region	N	Mean τ ± SD	Raw p-value	Holm-adj. p (32-family)
CHENG DeiT-Small	Method A (Telea)	Core	45	+0.1282 ± 0.2476	9.1269e-06	2.8293e-04

CHENG DeiT-Small	Method A (Telea)	Immediate	45	+0.0213 ± 0.0959	0.0267	0.5347
CHENG DeiT-Small	Method A (Telea)	Outer	45	+0.0063 ± 0.0981	0.2866	1.0000
CHENG DeiT-Small	Method A (Telea)	Remote	45	+0.0128 ± 0.0651	0.0801	1.0000
CHENG DeiT-Small	Method B (Texture)	Core	45	+0.1449 ± 0.2166	3.0881e-08	9.8820e-07
CHENG DeiT-Small	Method B (Texture)	Immediate	45	+0.0093 ± 0.1154	0.0190	0.4185
CHENG DeiT-Small	Method B (Texture)	Outer	45	+0.0344 ± 0.0809	0.0014	0.0379
CHENG DeiT-Small	Method B (Texture)	Remote	45	+0.0063 ± 0.0650	0.7122	1.0000
CHENG ResNet-50	Method A (Telea)	Core	45	+0.0311 ± 0.0567	6.5388e-04	0.0183
CHENG ResNet-50	Method A (Telea)	Immediate	45	+0.0125 ± 0.0505	0.0621	1.0000
CHENG ResNet-50	Method A (Telea)	Outer	45	+0.0111 ± 0.0969	0.4136	1.0000
CHENG ResNet-50	Method A (Telea)	Remote	45	+0.0095 ± 0.0364	0.0950	1.0000
CHENG ResNet-50	Method B (Texture)	Core	45	+0.0603 ± 0.1522	0.0173	0.3976
CHENG ResNet-50	Method B (Texture)	Immediate	45	+0.0146 ± 0.1740	0.3294	1.0000
CHENG ResNet-50	Method B (Texture)	Outer	45	+0.0592 ± 0.1986	0.0782	1.0000
CHENG ResNet-50	Method B (Texture)	Remote	45	-0.0238 ± 0.2278	0.7543	1.0000
BRISC DeiT-Small	Method A (Telea)	Core	45	+0.0517 ± 0.1197	0.0320	0.6072
BRISC DeiT-Small	Method A (Telea)	Immediate	45	+0.0239 ± 0.0628	0.0152	0.3643
BRISC DeiT-Small	Method A (Telea)	Outer	45	+0.0301 ± 0.0707	0.0049	0.1278
BRISC DeiT-Small	Method A (Telea)	Remote	45	+0.0140 ± 0.0564	0.2815	1.0000
BRISC DeiT-Small	Method B (Texture)	Core	45	+0.0563 ± 0.1007	1.6348e-04	0.0049
BRISC DeiT-Small	Method B (Texture)	Immediate	45	+0.0169 ± 0.0826	0.1313	1.0000
BRISC DeiT-Small	Method B (Texture)	Outer	45	+0.0564 ± 0.0947	2.8760e-04	0.0083
BRISC DeiT-Small	Method B (Texture)	Remote	45	-0.0139 ± 0.0648	0.1597	1.0000
BRISC ResNet-50	Method A (Telea)	Core	45	+0.0012 ± 0.0243	0.7885	1.0000
BRISC ResNet-50	Method A (Telea)	Immediate	45	-0.0004 ± 0.0217	0.8845	1.0000
BRISC ResNet-50	Method A (Telea)	Outer	45	+0.0017 ± 0.0306	0.6069	1.0000
BRISC ResNet-50	Method A (Telea)	Remote	45	+0.0028 ± 0.0130	0.1885	1.0000
BRISC ResNet-50	Method B (Texture)	Core	45	+0.0531 ± 0.1751	0.0950	1.0000
BRISC ResNet-50	Method B (Texture)	Immediate	45	+0.0702 ± 0.1665	0.0252	0.5285
BRISC ResNet-50	Method B (Texture)	Outer	45	+0.0608 ± 0.1542	0.0061	0.1537
BRISC ResNet-50	Method B (Texture)	Remote	45	+0.0438 ± 0.1741	0.6069	1.0000

Supplementary Table S2 Five-region topological area distributions across 120 unique intervention slices and area-normalised sensitivity per 1,000 pixels (Method A, inpainting).

Cohort	Region	N	Mean (px)	SD	Median	IQR	Min	Max
CHENG	Core	60	992.2	912.0	634.0	841.0	95	3688
CHENG	Immediate	60	511.1	209.7	433.5	328.8	200	979
CHENG	Outer	60	2107.8	613.2	1890.5	924.8	1201	3522
CHENG	Remote	60	236.8	230.4	151.0	220.2	17	980
CHENG	Sham	60	990.3	908.9	634.0	841.0	95	3688
BRISC	Core	60	1033.9	838.3	821.0	1081.2	89	4773
BRISC	Immediate	60	754.4	322.3	654.5	426.2	234	1795
BRISC	Outer	60	2369.4	652.4	2201.0	866.2	1229	3907
BRISC	Remote	60	290.3	242.5	211.5	293.0	20	1057
BRISC	Sham	60	1028.5	837.5	821.0	1081.2	89	4773
COMBINED (N=120)	Core	120	1013.1	872.5	726.5	952.8	89	4773
COMBINED (N=120)	Immediate	120	632.7	297.0	576.5	381.0	200	1795

COMBINED (N=120)	Outer	120	2238.6	644.0	2068.5	879.5	1201	3907
COMBINED (N=120)	Remote	120	263.6	237.1	175.0	268.8	17	1057
COMBINED (N=120)	Sham	120	1009.4	870.5	726.5	952.8	89	4773

Supplementary Table S3 Secondary inferential statistics: patient-clustered tests (Cheng, N=10 clusters), method concordance correlations (Method A vs B, N=90), correctly-classified-only subsets (N=70 DeiT, N=67 ResNet), cluster-level paired inference across 55 clusters (Wilcoxon W=483.0, p=0.0162; t-test t=3.29, p=0.00177), and formal Linear Mixed-Effects Model (N=180, beta=+0.074, SE=0.020, z=3.756, p<0.001).

Analysis Category	Model / Cohort	Condition	N	Observed Value	Statistic & p-value	Scientific Inference
Patient-Clustered Test (Cheng Core)	DeiT-Small	Method A (Telea Inpaint)	10	0.1349 +/- 0.1921	Two-sided W=8.0 (One-sided W=47.0), t=2.22; Two-sided p=0.0488; One-sided p=0.0244; t-test p=0.0534	Statistically significant above zero (two-sided Wilcoxon p = 0.0488, W = 8.0)
Patient-Clustered Test (Cheng Core)	ResNet-50	Method A (Telea Inpaint)	10	0.0285 +/- 0.0452	Two-sided W=8.0 (One-sided W=47.0), t=2.00; Two-sided p=0.0488; One-sided p=0.0244; t-test p=0.0771	Statistically significant above zero (two-sided Wilcoxon p = 0.0488, W = 8.0)
Method Concordance Correlation	DeiT-Small	Region: core	90	r = +0.8540, rho = +0.7631	t = 15.40; Pearson p = 1.05e-26, Spearman p = 2.27e-18	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Method Concordance Correlation	DeiT-Small	Region: immediate	90	r = +0.4862, rho = +0.5856	t = 5.22; Pearson p = 1.19e-06, Spearman p = 1.34e-09	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Method Concordance Correlation	DeiT-Small	Region: outer	90	r = +0.1375, rho = +0.1076	t = 1.30; Pearson p = 1.96e-01, Spearman p = 3.13e-01	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Method Concordance Correlation	DeiT-Small	Region: remote	90	r = +0.4361, rho = +0.2937	t = 4.55; Pearson p = 1.73e-05, Spearman p = 4.95e-03	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Method Concordance Correlation	ResNet-50	Region: core	90	r = +0.0893, rho = +0.0864	t = 0.84; Pearson p = 4.02e-01, Spearman p = 4.18e-01	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Method Concordance Correlation	ResNet-50	Region: immediate	90	r = -0.0518, rho = -0.0248	t = -0.49; Pearson p = 6.28e-01, Spearman p = 8.16e-01	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Method Concordance Correlation	ResNet-50	Region: outer	90	r = -0.0874, rho = +0.0019	t = -0.82; Pearson p = 4.13e-01, Spearman p = 9.86e-01	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Method Concordance Correlation	ResNet-50	Region: remote	90	r = +0.1749, rho = +0.0071	t = 1.67; Pearson p = 9.92e-02, Spearman p = 9.47e-01	High rank/linear concordance on core evidence; note correlation does not imply zero numerical bias
Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method A (Telea) - core	70	0.0791 +/- 0.2109	Wilcoxon W = 734.0; p = 0.0029	Significant sensitivity (p < 0.05)

Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method A (Telea) - immediate	70	0.0271 +/- 0.0719	Wilcoxon W = 662.0; p = 0.0007	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method A (Telea) - outer	70	0.0287 +/- 0.0683	Wilcoxon W = 649.0; p = 0.0005	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method A (Telea) - remote	70	0.0122 +/- 0.0535	Wilcoxon W = 939.0; p = 0.0757	Not statistically significant
Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method B (Texture) - core	70	0.0919 +/- 0.1820	Wilcoxon W = 364.0; p = 0.0000	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method B (Texture) - immediate	70	0.0203 +/- 0.0680	Wilcoxon W = 792.0; p = 0.0084	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method B (Texture) - outer	70	0.0572 +/- 0.0839	Wilcoxon W = 332.0; p = 0.0000	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	DeiT-Small (N=70)	Method B (Texture) - remote	70	-0.0100 +/- 0.0656	Wilcoxon W = 1020.0; p = 0.1929	Not statistically significant
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method A (Telea) - core	67	0.0229 +/- 0.0473	Wilcoxon W = 558.0; p = 0.0003	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method A (Telea) - immediate	67	0.0083 +/- 0.0416	Wilcoxon W = 873.0; p = 0.0966	Not statistically significant
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method A (Telea) - outer	67	0.0079 +/- 0.0796	Wilcoxon W = 1005.0; p = 0.4026	Not statistically significant
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method A (Telea) - remote	67	0.0068 +/- 0.0273	Wilcoxon W = 747.0; p = 0.0143	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method B (Texture) - core	67	0.0491 +/- 0.1713	Wilcoxon W = 862.0; p = 0.0836	Not statistically significant
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method B (Texture) - immediate	67	0.0254 +/- 0.1817	Wilcoxon W = 964.0; p = 0.2743	Not statistically significant
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method B (Texture) - outer	67	0.0705 +/- 0.1884	Wilcoxon W = 680.0; p = 0.0041	Significant sensitivity (p < 0.05)
Correctly-Classified-Only Sensitivity	ResNet-50 (N=67)	Method B (Texture) - remote	67	-0.0024 +/- 0.2233	Wilcoxon W = 1084.0; p = 0.7312	Not statistically significant
Cluster-Level Architectural Comparison	55 Patient/Case Clusters	Method A (Telea) Core Sensitivity (Cluster Means)	55	+0.0607 +/- 0.1368	Two-sided W = 483.0, t = 3.29; Wilcoxon two-sided p = 0.0162 (p = 0.0162); t-test p = 0.00177 (p = 0.00177)	Statistically significant cluster-robust core reliance advantage for DeiT-Small
Linear Mixed-Effects Model (LME)	Pooled Multi-Cohort (55 Clusters)	Method A (Telea) Core Sensitivity: tau ~ is_vit + (1|cluster_id)	180	beta = +0.0738 (95% CI [0.035, 0.112])	Wald z = 3.756, REML log-lik = 93.99, res_var = 0.0174; p = 1.7236e-04 (p < 0.001)	DeiT exhibits significantly greater lesion-core reliance (beta=+0.074, SE=0.020, z=3.756, p<0.001)
Architectural Comparison (Paired Nonparametric)	Pooled Multi-Cohort (N=90 matched slices)	Method A (Telea) Core Sensitivity	90	+0.0738 +/- 0.2037	Two-sided W = 1425.0 (One-sided W = 2670.0), t = 3.44; Two-sided p = 0.0123 (p = 0.0123, one-sided p = 0.0061); t-test p = 0.0004	DeiT-Small exhibits significantly greater core sensitivity (paired two-sided p = 0.0123)