

Impact of superficial lateral drainage on tree root distribution and forest community structure in French Guiana

SANVITO Emilie^{1*}

DOBETSBERGER Greta²

THION Clément³

Abstract

This study investigates the impact of drainage barriers on tree root distribution and forest structure in Paracou, French Guiana. Using the depth at which a horizon high in silt (HSD, high silt depth) occurs as a proxy for drainage barrier, the study explores the relationship between drainage barriers, topography, and forest community structure. Despite expectations of reduced root density in soil horizons with high silt content, no significant reduction of root density was found. Species richness and tree density showed a slight but significant decrease with increasing HSD, while diameter at breast height (DBH) exhibited minimal variation. The strong association between silt and mica content suggests the potential use of mica as a practical proxy for silt in future studies, streamlining data collection. These findings contribute valuable insights into the complexity of soil-root interactions in tropical ecosystems. Further research with increased sampling density and direct data collection is recommended to refine and expand upon these initial observations.

Keywords

root, tropical forest, community structure, drainage, pedology, drought, Paracou, French Guyana

¹Tropimundo

Street address, Zip code, Belgium.

²Université des Antilles

Street address, Zip code, France.

³AgroParisTech Montpellier

648 Rue Jean François Breton, 34090, France.

*Corresponding author: emilie@sanvito.be,

1. Introduction

In tropical ecology, the relationship between root distribution and soil abiotic factors is crucial to forest community structure (Schlesinger, 1991; Bardgett et al., 2014). Roots are essential for the nutrition and water uptake of trees (Hutchings et al., 2003; Hodge, 2009). Their distribution is driven by factors like soil texture, nutrient content, topography, and water content (Freschet et al., 2021).

A large part of the forests in French Guiana features a topography alternating between hills and valleys (Ferry et al., 2003). Acrisol (WRB, 2022), characterised by a superficial lateral drainage (SLD), is the

typical soil type on slopes (Fan et al., 2017). Acrisol is characterised by a silty texture horizon with low permeability that functions as a drainage barrier in the soil (Ferry et al., 2003; Humbel, 1978).

Drainage type, and to a lesser extent topography, are the main factors influencing forest community structure (Pélissier et al., 2002; Sabatier et al., 1997). Root vertical distribution (RVD) is the distribution of roots in units of biomass over sequential soil layers (Freschet et al., 2021). In soil with superficial lateral drainage, root development is inhibited by the drainage barrier (Ferry et al., 2003).

Tree roots are therefore concentrated in the uppermost layers of the soil (Humbel,

1978), characterised by low water reserves. Consequently, roots in this soil type may undergo drought stress during the dry season. During the rainy season, this impermeable silty horizon will cause a temporary water table in the soil (Humbel, 1978; Péliissier et al., 2002), which creates anoxic stress for tree roots (Sabatier et al., 1997).

Our study takes place at Paracou research station in French Guiana. Epron et al. (2006) confirmed the presence of SLD in this area. They found that the depth of the drainage barrier on slopes depends on the topographical position, with deeper barriers at higher elevation. However, the specific ways in which drainage barriers influence both roots and forest structure remain understudied. Therefore, we examined whether drainage barriers restrict tree roots on slopes in Paracou, exploring their role in shaping forest community structure. We investigated the effect of the depth, at which the drainage barrier occurs, on root vertical distribution and on 3 different forest structure indicators (DBH, species richness and tree density). Our main hypotheses are: (1) root density not only decreases with depth, but is also negatively influenced by the drainage barrier; (2) the depth at which the drainage barrier appears can be a predictor of DBH, species richness and tree density.

2. Material and method

2.1 Study site

The study took place at Paracou research station in French Guiana (5°18'N; 52°53'W), a 40000 ha domain owned by CNES (National Centre for Space Studies) containing 16 permanent plots studied by the French research centre CIRAD for over 35 years. Elevation ranges from 5 to 50 m, and the mean annual temperature is 26 °C, with an annual range of 1–1.5 °C. Rainfall averages 2980 mm per year with a 3-month dry season (< 100 mm month) from mid-August to mid-November. The landscape is characterised by a patchwork of low hills (100–300 m wide and 20–35 m high) separated by narrow streams. The soil type in this area is mostly Acrisol (WRB, 2022) and soil depth is restricted by a poorly draining, loamy




Figure 1. Study site and Pit locations at Paracou research station. P1-4 represent pit 1-4.

saprolite, typically encountered at a depth of around 1 m (Epron et al., 2006). Sampling was carried out as part of the ALT-project (Amazonian Landscapes in Transition 2022-2025).

2.2 Pit data collection

Pit excavation allows the collection of root distribution data (Freschet et al., 2021). Four pits were sampled between September 11 and 16, 2023, just outside of plot 16, next to the subplots 24-25 (Fig. 1), to avoid disturbances on inventoried trees.

We implemented a standardized experimental design, ensuring uniform topographical conditions for the soil pits. All pits were excavated at a mid-slope position with a consistent inclination of 17-18%, characterized by superficial lateral drainage. Potential pits were prospected using an auger in order to dig pits with maximum HSD variability. Based on (Epron et al., 2006) we expected the HSD to be between 60 cm and 80 cm deep. Rectangular pits, 100 cm deep and 50 cm wide, were dug with pickaxes and spades.

After cleaning the exposed soil surface in the pits by scraping, colour and texture changes were used to identify horizons (Appendix C Fig. @ref(fig:pitPlot_TextureProfile)) on one side of the pit (Ferry et al., 2003). For each horizon, the depth was measured, mica content was visually estimated from 0 (no mica) to 3 (high mica content), and soil samples were moistened to estimate the texture by touch (Jahn et al., 2006). We used Jamagne et al. (1977) soil texture triangle. For two sides of the pit, we counted the

roots using a 100 x 50 cm grid subdivided in squares of 10 x 10 cm (Freschet et al., 2021), to characterise RVD. In each square, roots were counted and classified by diameter (< 2 mm, 2-5 mm, 5-10 mm and 10 mm).

2.3 Auger and botanical data

In addition to our pit data, we used 36 auger samples collected during a 2022 data collection campaign on the P16 of Paracou, in the subplots 14, 15, 19 and 20 (Appendix A, Fig. 4). The auger samples were positioned with a spacing of 33 meters between each sampling point. In contrast to the pit data collection, root abundance was not sampled. For the complete auger collection method, we refer to Dogny Piette and Corbera Serrajordia (Dogny Piette and Corbera Serrajordia). The botanical data corresponds to the 2020 inventory of the GUYAFOR tree database, which monitors trees of Diameter at Breast Height (DBH) superior to 10 cm. In our study area, this data was completed with data from the ALT project, which also included trees of DBH > 1 cm.

2.4 Data analysis

2.4.1 Pit data analysis

We regrouped the soil textures into a categorical variable “silt content”, identified using the texture triangle (Jamagne et al., 1977): low (A, Alo, AS), intermediate (SL, SA, S), and high (LSA, LAS, LS). We defined the depth (cm) at which the silt content changed to high for each pit, which we refer to as “high silt depth” (HSD). HSD is used to determine the depth of the drainage barrier, which is characterised by a silty texture (Ferry et al., 2003; Humbel, 1978). Given the small sample size (4 pits), we chose to only conduct visual analysis of HSD effect on RVD. As roots were counted per layer of 10 cm, and soil texture was determined per soil horizon, we assigned a texture class to every 10 cm layer of the soil.

In order to approximate root biomass, a root index was calculated by weighing each root count with its corresponding minimum diameter class and summing the totality weighted count. Using the minimum diameter causes leadsus to an underestimation of the

total root biomass but although it remains valuable is still useful for pit comparison.

$RI_d = \sum_{i=1}^4 r_i n_i$, with d the layer depth (cm), i the indice of the diameter class (< 2 mm, 2-5 mm, 5-10 mm and 10 mm), $r_i \in [1, 2, 5, 10]$ the minimum root diameter, n_i the count of root in diameter class i . The minimum root diameter of the first class (<2mm) is set to 1mm as we assume it is the smallest diameter detectable by bare eye.

We then assessed the relation between root index and silt content. Root index was compared between pits 2 and 3 at the depth of 50-70 cm. This depth range is where maximum HSD variation between two pits occurs. Fixing the depth accounts for the effect that root index decreases exponentially with depth. We calculated the decrease of root index in the selected depth range (total root index at 50-70 cm divided by total root index at 0-50 cm) for pits 2 and 3 and compared them.

2.4.2 Auger and botanical data analysis

Out of the 36 auger measurements available, we focused our analysis on the subset of 29 measurements that contained texture class data. The auger points were located using the coordinates of the closest inventoried tree on the plot. As for the pit data, soil textures in the pits were grouped into “high”, “intermediate”, and “low” silt content, and HSD was defined for each auger point following the same methodology as employed for the pit data. We defined the depth (cm) at which the silt content changed to high for each auger sample, which we refer to as “high silt depth” (HSD). HSD is used to indicate the depth of the drainage barrier, characterised by a silty texture (Ferry et al., 2003; Humbel, 1978). For auger points without any horizons high in silt, we defined HSD to be 100 cm, which is the maximum depth of the auger measurements. As topography, besides drainage type, strongly influences forest community structure (Baldeck et al., 2013; Ferry et al., 2010) and can be correlated with soil texture and drainage (Epron et al., 2006), we did a correlation test (Spearman method due to non-normality of HSD) between elevation (m) and HSD (cm).

We mapped the area where auger samples were taken and created contour lines based on the values of HSD of the different auger points (Appendix A Fig. 4). The contour lines were created with the Contour Plugin in QGIS. We employed 10 cm contour line intervals, resulting in the formation of seven distinct HSD classes, from 40 cm to 100 cm deep. With this approach, we interpolated HSD values spatially across nearly the entire 4 ha study area. Every tree on the map could therefore be linked to a certain HSD class and the corresponding coverage area (m²) of each HSD class was also determined. Only slopes were considered during further analysis. Topographical data comes from (Dourdain et al., 2022), which defines slope as the zone with a slope > 25 % and the absence of a permanent water table, no deeper than 1 m in the middle of the dry season. For each HSD class species richness was extracted from the botanical data by counting the number of species present in the area. The total number of trees was extracted by counting the number of trees present in the area. Due to the ordinal nature of the different HSD classes, they cannot be considered as a categorical variable. Hence, we assigned a specific intermediate HSD value to each HSD class. For example, an HSD class from 40 cm to 50 cm was designated the value of 45 cm. This creates a continuous HSD-value, allowing us to perform a Poisson regression for both species richness and number of trees (HSD continuous values as predictor, species richness or number of trees as response). The areas covered by a specific HSD class varied between 1015 m² (HSD 40-50 cm) and 5494 m² (HSD 90-100 cm), with the total area considered being 21055 m². To take into account that some HSD classes cover a bigger area than others, offset is set to logarithm of area per HSD class in the model. This way the number of trees and species richness per area are modelled. The relationship between DBH (cm) and HSD (cm) did not show linear dependency (Fig. 3c and 3d), making it difficult to construct a precise model. Muscovite (white mica) degrades into kaolin-

ite (Nicolini et al., 2009) with a particle size classified as silts. Thus, to complement our detection of the drainage barrier, we try to use mica content as a complementary indicator of silt content. A chi-squared test was done to quantitatively assess the association between silt content (categorical variable) and mica content (categorical variable) in the different horizons of the auger data. Cramer's V was calculated to indicate the strength and direction of the association. We tried to replace silt content by mica content for the pit data analysis, and verified whether the results were similar. All analyses were performed in R (version 4.2.2) (R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>) and QGIS (version 3.28 Firenze).

3. Results

3.1 Pit data

HSD ranges across the pits, spanning from 50 cm to 70 cm (Fig. 2a). In contrast to our expectations, the four pits show similar root index variations with depth, regardless the variation in silt content. Figure 2b shows a general decrease of root index with depth, from 60 roots at a depth of 10 cm to 10 roots at 50 cm. Focussing on the specific depth range of 50-70 cm, it becomes evident that the root index of P3 (pit 3) does not exceed that of P2 (pit 2), even though P3 shows high silt content and P2 intermediate silt content. The ratio of root index in the focus range (50-70 cm) (Fig. 2a) and root index in the depth range above (0-50 cm) between P2 and P3 also showed no difference (0.137 for P2 and 0.142 for P3).

Mica as an indicator for silt

The chi-squared test assessing the association between silt and mica content in the auger samples was highly significant ($\chi^2 = 57.154$, p-value < 0.001). Cramer's V, assessing the strength and direction of the association, was 0.58, supporting a strong positive relationship. The association is depicted in a mosaic plot in Appendix B Fig. 5, illustrating an increase of silt content with rising mica content. By re-




Figure 2. a) Silt content (low, intermediate, high) and b) root index, calculated by weighing each root count with its corresponding maximum diameter class, per layer of 10 cm across pits excavated in September 2023. P1-4 correspond to pits 1-4.

placing mica content with silt content in the pit data analysis where the relation between root index and silt content are assessed, we found a similar result (Appendix C, Fig. 7).

3.2 Auger data

A significant but small decrease (p -value = 0.00245; Estimate = -0.0044) in species richness with HSD class was detected. Among all HSD classes, the range of 40 to 50 cm stands out with notably higher species richness per ha (Fig. 3a). This range contains the uppermost observed HSD. Species richness per ha is lowest for the 90 cm to 100 cm HSD range. For HSD class “No high silt” species richness was higher than for the HSD classes 70-100 cm. Tree density significantly decreases with HSD class (p -value < $2e-16$; Estimate = -0.013), although Fig. 3b shows that the tree density first increases between HSD 40 cm to 70 cm. Tree density increases a little when the HSD is not detected, the lowest tree density being present at HSD 90 cm to 100 cm range. Fig. 3c and 3d illustrate the relationship between the DBH and HSD classes. The mean DBH values and quantiles are very similar over

all classes (mean = 4,9 cm), primarily due to the high representation of trees with small diameter (minimum diameter count > 1 cm). A pattern is visible in the maximum values of the outliers, representing trees with a DBH > 50 cm, which increases with HSD class.

A significant negative correlation was found between elevation (m) and HSD (cm) (ρ = -0,634; p -value = 0.0002246). A higher position along the topographical gradient therefore indicates lower HSD.

4. Discussion

4.1 Root distribution with HSD and depth:

Our findings on RVD align with the literature, illustrating that most roots are present within the first 30 cm of the soil (Freschet et al., 2021; Schenk and Jackson, 2002). HSD ranges across the pits, spanning from 50 cm to 70 cm (Fig. 2a), coincides with the depth range (60-80 cm) of the drainage barrier identified by Epron et al. (2006). This strengthens our choice to use of HSD as an estimator of the drainage barrier depth.

We expected that the drainage barrier




Figure 3. Forest structure indicators relation with HSD (high silt depth) class (cm). a) Species richness (per ha) in function of HSD classes, b) Tree density (per ha) in function of HSD classes, c) DBH (diameter at breast height in cm) in function of HSD classes, d) DBH filtered (max value = 10 cm) in function of HSD classes

restricts roots, and therefore to find less roots below the HSD, as was observed by Humbel (1978). However, there is no visually discernible effect of HSD on vertical root distribution (Fig. 2b), no matter the root diameter (Appendix C, Fig. 6). The absence of an HSD effect might be because roots of certain species can penetrate the drainage barrier, which suggests a deeper water uptake and therefore higher drought resistance (Ferry et al., 2003). The absence of an observed effect may also be due to the small HSD variation between the sampled pits (maximum variation of 20 cm). Additionally, the small sample size of 4 pits precluded the performance of a statistical test. More soil pits would be needed to study root distribution in additional soil profiles.

4.2 Mica-silt correlation

Our findings indicate a strong association between silt and mica content. Given that mica is shiny and easily detectable, identifying HSD by high mica content instead of high silt content might simplify and expedite the data col-

lection process. This approach may offer a more efficient means of assessing drainage barrier depth in Paracou.

5. Acknowledgements

We are grateful to our supervisors Vincyane Badouard, Pierre-André Wagner for their support and patience. Further we want to thank our colleagues from the FTH module 2023 and the EFT Master 2022, who provided preliminary work and data. We thank Timothy Chubb for support with scientific writing and language. We are also grateful to Lucia Clarotto, Stéphane Traissac, Camille Salmon and Mélaïne Aubry-Kientz for the fruitful discussions on general context of this study and suggestions of determinant methodological choices.

6. Appendix

Le temps passe et la mort approche.




Figure 4. HSD contour line map of subplots 14, 15, 19 and 20 of the P16.




Figure 5. Mosaic plot displaying the relationship between mica and silt content. Dark grey represents high silt content, grey represents moderate silt content, and light grey signifies low silt content. Mica content ranges from 0 to 3, denoting low (0) to high (3) levels.

References

Baldeck, C. A., K. E. Harms, J. B. Yavitt, R. John, B. L. Turner, R. Valencia, H. Navarrete, S. J. Davies, G. B. Chuyong, D. Kenfack, D. W. Thomas, S. Madawala, N. Gunatilleke, S. Gunatilleke, S. Bunyavejchewin, S. Kiratiprayoon, A. Yaacob, M. N. N. Supardi, and J. W. Dalling (2013, February). Soil resources and topography shape local tree community structure in tropical forests. *Proceedings of the Royal Society B: Biological Sciences* 280(1753), 20122532.

Bardgett, R. D., L. Mommer, and F. T. De Vries (2014, December). Going underground: Root traits as drivers of ecosystem processes. *Trends in Ecology & Evolution* 29(12), 692–699.

Dogny Piette, C. and M. Corbera Serrajordia.




Figure 6. Root quantity for all diameter classes for all pits. rd1, rd2, rd3, rd4 correspond respectively with diameter classes of < 2 mm, 2-5 mm, 5-10 mm and 10 mm. For silt content, high is LSA or LAS, intermediate is SL or SA, and low is AS or A.




Figure 7. Root index across soil depths in different pits, considering varying levels of mica content ranging from low (0) to high (3).




Figure 8. Soil texture profiles of the four pits.

- Abiotic factors related to hydrology and soil have an effect on local spatial distribution of tropical tree species in Paracou, French Guiana. *1*(1).
- Dourdain, A., G. Jaouen, and G. Derroire (2022). GIS DATA DICTIONARY V 7.0.
- Epron, D., A. Bosc, D. Bonal, and V. Freycon (2006, September). Spatial variation of soil respiration across a topographic gradient in a tropical rain forest in French Guiana. *Journal of Tropical Ecology* 22(5), 565–574.
- Fan, Y., G. Miguez-Macho, E. G. Jobbágy, R. B. Jackson, and C. Otero-Casal (2017, October). Hydrologic regulation of plant rooting depth. *Proceedings of the National Academy of Sciences* 114(40), 10572–10577.
- Ferry, B., V. Freycon, and D. Paget (2003). Genèse et fonctionnement hydrique des sols sur socle cristallin en Guyane. *Revue forestière française* 55(sp), 37–59.
- Ferry, B., F. Morneau, J.-D. Bontemps, L. Blanc, and V. Freycon (2010). Higher treefall rates on slopes and waterlogged soils result in lower stand biomass and productivity in a tropical rain forest. *Journal of Ecology* 98(1), 106–116.
- Freschet, G. T., L. Pagès, C. M. Iversen, L. H. Comas, B. Rewald, C. Roumet, J. Klimešová, M. Zadworny, H. Poorter, J. A. Postma, T. S. Adams, A. Bagniewska-Zadworna, A. G. Bengough, E. B. Blancaflor, I. Brunner, J. H. C. Cornelissen, E. Garnier, A. Gessler, S. E. Hobbie, I. C. Meier, L. Mommer, C. Picon-Cochard, L. Rose, P. Ryser, M. Scherer-Lorenzen, N. A. Soudzilovskaia, A. Stokes, T. Sun, O. J. Valverde-Barrantes, M. Weemstra, A. Weigelt, N. Wurzbürger, L. M. York, S. A. Batterman, M. Gomes de Moraes, Š. Janeček, H. Lambers, V. Salmon, N. Tharayil, and M. L. McCormack (2021). A starting guide to root ecology: Strengthening ecological concepts and standardising root classification, sampling, processing and trait measurements. *New Phytologist* 232(3), 973–1122.
- Hodge, A. (2009, June). Root decisions. *Plant, Cell & Environment* 32(6), 628–640.
- Humbel, F.-X. (1978). Caractérisation, par des mesures physiques, hydriques et d'enracinement, de sols de Guyane française à dynamique de l'eau superficielle. extrait de l'A.F.E.S 2, ORSTOM.
- Hutchings, M. J., E. A. John, and D. K. Wijesinghe (2003, September). Toward understanding the consequences of soil heterogeneity for plant populations and communities. *Ecology* 84(9), 2322–2334.
- Jahn, R., H. P. Blume, V. B. Asio, O. Spaargaren, and P. Schad (2006). *Guidelines for Soil Description, 4th Edition*. Rome: FAO.
- Jamagne, M., J. C. Begon, and R. Hardy (1977). Soil mapping, a vital element in the management and conservation of rural resources. *Pedologie* 27(1), 9–43.
- Nicolini, K. P., K. C. Lombardi, W. H. Schreiner, I. Mazzaro, F. Wypych, and A. S. Mangrich (2009, December). Evidence of weathering stages of phyllosilicates from biotite/muscovite to kaolinite, probed by EPR spectroscopy. *Mineralogy and Petrology* 97(1), 139–144.
- Pélissier, R., S. Dray, and D. Sabatier (2002, October). Within-plot relationships between tree species occurrences and hydrological soil constraints: An example in French Guiana investigated through canonical correlation analysis. *Plant Ecology* 162(2), 143–156.
- Sabatier, D., M. Grimaldi, M.-F. Prévost, J. Guillaume, M. Godron, M. Dosso, and P. Curmi (1997, July). The influence of soil cover organization on the floristic and structural heterogeneity of a Guianan rain forest. *Plant Ecology* 131(1), 81–108.
- Schenk, H. J. and R. B. Jackson (2002). The Global Biogeography of Roots. *Ecological Monographs* 72(3), 311–328.

Schlesinger, W. (1991). Biotic Feedbacks in the Global Climatic System: Will the Warming Feed the ... - Google Books.
<https://books.google.com/books?hl=en&lr=&id=Tyrc32Pk5MkC&oi=fnd&pg=PA159&dq=schlesinger%201991%20belowground%20carbon&f=false>.
SB18BiM7kc#v=onepage&q=schlesinger%201991%20belowground%20carbon&f=false.

WRB, I. W. G. (2022, July). World Reference Base for Soil Resources. International soil classification system for naming soils and creating legends for soil maps. 4th edition.