

# Continuous Space MDPs

# Last Time

- Neural Network Function Approximation

$$\hat{y} = f_{\theta}(x)$$


$$\{(x_i, y_i)\}_{i=1}^n$$

# Guiding Questions

# Guiding Questions

- What tools do we have to solve MDPs with continuous  $S$  and  $A$ ?

# Current Tool-Belt



# Today: Four Tools

1. LQR
2. Fitted Value Iteration
3. Sparse UCT/Progressive Widening
4. MPC

# Notation: Continuous Random Variables

Term	Definition	Coinflip Example	Uniform Example						
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{\text{h}, \text{t}\}$ or $\{0, 1\}$	$\text{Bernoulli}(0.5)$						
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	<table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></tbody></table>	x	P(x)	0	0.5	1	0.5
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$							

# Notation: Continuous Random Variables

Term	Definition	Coinflip Example	Uniform Example						
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$\mathcal{U}([0, 1])$						
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	<table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></tbody></table>	x	P(x)	0	0.5	1	0.5
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$							

# Notation: Continuous Random Variables

Term	Definition	Coinflip Example	Uniform Example						
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	<table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></tbody></table>	x	P(x)	0	0.5	1	0.5
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$							

# Notation: Continuous Random Variables

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}([0, 1])$ Uniform Example
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	$p(x) = \mathbf{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$	


# Notation: Continuous Random Variables

Term	Definition	Bernoulli(0.5)	$\mathcal{U}([0, 1])$
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	$p(x) = \mathbf{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ 
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$	


# Notation: Continuous Random Variables

Term	Definition	Bernoulli(0.5)	$\mathcal{U}([0, 1])$
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	$p(x) = \mathbf{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$  $P(X \in [a, b]) = \int_a^b p(x) dx$
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$ $= 0.5$	

# Notation: Continuous Random Variables

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}([0, 1])$ Uniform Example
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	$p(x) = \mathbf{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$  $P(X \in [a, b]) = \int_a^b p(x) dx$ $P(X = 0.5) =$
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$ $= 0.5$	


# Notation: Continuous Random Variables

Term	Definition	Bernoulli(0.5)	$\mathcal{U}([0, 1])$
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	$p(x) = \mathbf{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$  $P(X \in [a, b]) = \int_a^b p(x) dx$ $P(X = 0.5) = 0$
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$ $= 0.5$	

# Notation: Continuous Random Variables

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}([0, 1])$ Uniform Example
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	$p(x) = \mathbf{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$  $P(X \in [a, b]) = \int_a^b p(x) dx$ $P(X = 0.5) = 0$
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$ $= 0.5$	$E[X] = \int_{x \in X} x p(x) dx$

# Notation: Continuous Random Variables

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}([0, 1])$ Uniform Example
$\text{support}(X)$ $x \in X$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table	$p(x) = \mathbf{1}_{[0,1]}(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$  $P(X \in [a, b]) = \int_a^b p(x) dx$ $P(X = 0.5) = 0$
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$ $= 0.5$	$E[X] = \int_{x \in X} x p(x) dx$ $= 0.5$

# Rules for Continuous RVs

## Discrete


- 1) a)  $0 \leq P(X | Y) \leq 1$
- b)  $\sum_{x \in X} P(x | Y) = 1$
- 2)  $P(X) = \sum_{y \in Y} P(X, y)$

## Continuous

- 1)

$$3) P(X | Y) = \frac{P(X, Y)}{P(Y)}$$
$$P(X, Y) = P(X | Y) P(Y)$$

# Rules for Continuous RVs



**Discrete**

- 1) a)  $0 \leq P(X | Y) \leq 1$
- b)  $\sum_{x \in X} P(x | Y) = 1$
- 2)  $P(X) = \sum_{y \in Y} P(X, y)$

**Continuous**

1)  $0 \leq p(X | Y)$

$p(x) = 10 \mathbf{1}(0 \leq x \leq 0)$

3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

# Rules for Continuous RVs

## Discrete

- 1) a)  $0 \leq P(X | Y) \leq 1$
- b)  $\sum_{x \in X} P(x | Y) = 1$
- 2)  $P(X) = \sum_{y \in Y} P(X, y)$

## Continuous

- 1)  $0 \leq p(X | Y)$   
 $\int_X p(x|Y) dx = 1$

$$3) P(X | Y) = \frac{P(X, Y)}{P(Y)}$$
$$P(X, Y) = P(X | Y) P(Y)$$

# Rules for Continuous RVs

## Discrete

- 1) a)  $0 \leq P(X | Y) \leq 1$
- b)  $\sum_{x \in X} P(x | Y) = 1$
- 2)  $P(X) = \sum_{y \in Y} P(X, y)$
- 3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$   
$$P(X, Y) = P(X | Y) P(Y)$$

## Continuous

- 1)  $0 \leq p(X | Y)$   
$$\int_X p(x | Y) dx = 1$$
- 2)  $\underbrace{p(X)}_{\substack{\longrightarrow \\ \text{---}}} = \int_Y \underbrace{p(X, y)}_{\substack{\longleftarrow \\ \text{---}}} dy$

# Rules for Continuous RVs

## Discrete

- 1) a)  $0 \leq P(X | Y) \leq 1$
- b)  $\sum_{x \in X} P(x | Y) = 1$
- 2)  $P(X) = \sum_{y \in Y} P(X, y)$
- 3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$   
$$P(X, Y) = P(X | Y) P(Y)$$

## Continuous

- 1)  $0 \leq p(X | Y)$   
$$\int_X p(x | Y) dx = 1$$
- 2)  $p(X) = \int_Y p(X, y) dy$
- 3)  $p(X | Y) = \frac{p(X, Y)}{p(Y)}$   
$$p(X, Y) = p(X | Y) p(Y)$$

$$\begin{bmatrix} 4 & 2 & 2 \\ 2 & 4 & 2 \\ 2 & 2 & 4 \end{bmatrix}$$

# Multivariate Gaussian Distribution

$$x = [x_1, x_2]$$


$$\mathcal{N}(\mu, \Sigma)$$

## Joint Distribution

$$p(x) = \mathcal{N}(x, \Sigma)$$

$$p(x) = \frac{\exp\left(-\frac{1}{2}(x-\mu)^T \Sigma^{-1} (x-\mu)\right)}{(2\pi)^{n/2} |\Sigma|^{1/2}}$$

$$\mu = [3, 3] \quad \Sigma = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$$




## Conditional Distribution

$$p(x_1 | x_2) = \mathcal{N}(\bar{\mu}_1, \bar{\Sigma}_1)$$


$$\bar{\mu}_1 = \mu_1 + \Sigma_{12} \Sigma_{22}^{-1} (x_2 - \mu_2)$$

$$\bar{\Sigma}_1 = \Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21}$$

$$\mu = [3, 3] \quad \Sigma = \begin{bmatrix} 9 & 2 \\ 2 & 4 \end{bmatrix}$$




$$p(x_1 | x_2 = 4)$$



## Marginal Distribution

$$p(x_1) = \mathcal{N}(\mu_1, \Sigma_{11})$$



# Continuous $S$ and $A$

# Continuous $S$ and $A$

e.g.  $S \subseteq \mathbb{R}^n$ ,  $A \subseteq \mathbb{R}^m$

# Continuous $S$ and $A$

e.g.  $S \subseteq \mathbb{R}^n$ ,  $A \subseteq \mathbb{R}^m$

The old rules still work!

$$V^*(s) = \max_a (R(s, a) + \gamma \mathbb{E}[V^*(s')])$$

$$V^\pi(s) = \dots$$

$$B[V](s) = \max_a (R(s, a) + \gamma \mathbb{E}_{\substack{s' \sim T(s'|s, a)}} [V(s')])$$

hard!!!!

Optimization is harder  
than integration

$$\int_{S' \in S} T(s'|s, a) V(s') ds'$$

hard!

Monte Carlo

# 1. Linear Dynamics, Quadratic Reward

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m$$

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m \quad T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma)$$

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m \quad T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma)$$

$$s' = T_s s + T_a a + w$$

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m$$

$$T(s' | s, a) = \mathcal{N}(\underbrace{T_s s + T_a a}_{\text{red}}, \underline{\Sigma})$$

$$s' = \underbrace{T_s s + T_a a}_{\text{red}} + \underbrace{w}_{\text{red}} \quad w \sim \mathcal{N}(0, \Sigma)$$

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m$$

$$T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma)$$

$$s' = T_s s + T_a a + w \quad w \sim \mathcal{N}(0, \Sigma) \quad (\text{Also works with other zero-mean } w.)$$

# 1. Linear Dynamics, Quadratic Reward


$$S = \mathbb{R}^n, A = \mathbb{R}^m$$

$$T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma)$$

$$R(s, a) = s^\top \underline{R_s} s + a^\top \underline{R_a} a$$

$$s' = T_s s + T_a a + w \quad w \sim \mathcal{N}(0, \Sigma) \quad (\text{Also works with other zero-mean } w.)$$

$$R(s, a) = R(s) + R(a)$$



# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m$$

$$T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma)$$

$$R(s, a) = s^\top R_s s + a^\top R_a a$$

$$s' = T_s s + T_a a + w \quad w \sim \mathcal{N}(0, \Sigma) \quad (\text{Also works with other zero-mean } w.)$$

Finite Horizon:

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m \quad T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma) \quad R(s, a) = s^\top R_s s + a^\top R_a a$$

$$s' = T_s s + T_a a + w \quad w \sim \mathcal{N}(0, \Sigma) \quad (\text{Also works with other zero-mean } w.)$$

Finite Horizon:  $U_h^*(s) = \max_{\pi} \left[ \sum_{t=0}^h R(s_t, a_t) \right]$

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m \quad T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma) \quad R(s, a) = s^\top R_s s + a^\top R_a a$$

$$s' = T_s s + T_a a + w \quad w \sim \mathcal{N}(0, \Sigma) \quad (\text{Also works with other zero-mean } w.)$$

Finite Horizon:  $U_h^*(s) = \max_{\pi} \left[ \sum_{t=0}^h R(s_t, a_t) \right]$   $\pi_h^*$  is "optimal h-step policy"

# 1. Linear Dynamics, Quadratic Reward

$$S = \mathbb{R}^n, A = \mathbb{R}^m \quad T(s' | s, a) = \mathcal{N}(T_s s + T_a a, \Sigma) \quad R(s, a) = s^\top R_s s + a^\top R_a a$$

$$s' = T_s s + \underbrace{T_a a}_w + w \quad w \sim \mathcal{N}(0, \Sigma) \quad (\text{Also works with other zero-mean } w.)$$

Finite Horizon:  $U_h^*(s) = \max_{\pi} \left[ \sum_{t=0}^h R(s_t, a_t) \right]$   $\pi_h^*$  is "optimal h-step policy"

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$

# 1. Linear Dynamics, Quadratic Reward

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

$R(\alpha)$

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

$V_1 = R_s$

$q_1 = 0$



Inductive step: show that if  $U_t^* = s^\top V_t s + q_t$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

Inductive step: show that if  $U_t^* = s^\top V_t s + q_t$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

$$U_{t+1}^*(s) = \max_a (R(s, a) + \gamma E [U_t^*(s)])$$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

Inductive step: show that if  $U_t^* = s^\top V_t s + q_t$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

$$\begin{aligned} U_{t+1}^*(s) &= \max_a (R(s, a) + \gamma E [U_t^*(s)]) \\ &= \max_a \left( s^\top R_s s + a^\top R_a a + \int p(w) U_t(\underbrace{T_s s + T_a a + w}) dw \right) \end{aligned}$$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

Inductive step: show that if  $\underline{U_t^*} = \underline{s^\top V_t s + q_t}$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

$$\begin{aligned} U_{t+1}^*(s) &= \max_a (R(s, a) + \gamma E [U_t^*(s)]) \\ &= \max_a (s^\top R_s s + a^\top R_a a + \int p(w) U_t(T_s s + T_a a + w) dw) \\ &= s^\top R_s s + \max_a \left( a^\top R_a a + \underbrace{\int p(w) (T_s s + T_a a + w)^\top V_t (T_s s + T_a a + w) dw}_{\underbrace{+ q_t}} \right) \end{aligned}$$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

Inductive step: show that if  $U_t^* = s^\top V_t s + q_t$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

$$\begin{aligned} U_{t+1}^*(s) &= \max_a (R(s, a) + \gamma E [U_t^*(s)]) \\ &= \max_a (s^\top R_s s + a^\top R_a a + \int p(w) U_t(T_s s + T_a a + w) dw) \\ &= s^\top R_s s + \max_a (a^\top R_a a + \int p(w) (T_s s + T_a a + w)^\top V_t (T_s s + T_a a + w) + q_t) dw \\ &= \underbrace{s^\top R_s s + s^\top T_s^\top V_t T_s s}_{\text{max}_a (a^\top R_a a + 2s^\top T_s^\top V_t T_a a + a^\top T_a^\top V_t T_a a)} + \underbrace{\int p(w) w^\top V_t w dw + q_t}_{\text{max}_a (a^\top R_a a + 2s^\top T_s^\top V_t T_a a + a^\top T_a^\top V_t T_a a)} \end{aligned}$$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

Inductive step: show that if  $U_t^* = s^\top V_t s + q_t$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

$$\begin{aligned} U_{t+1}^*(s) &= \max_a (R(s, a) + \gamma E [U_t^*(s)]) \\ &= \max_a (s^\top R_s s + a^\top R_a a + \int p(w) U_t(T_s s + T_a a + w) dw) \\ &= s^\top R_s s + \max_a (a^\top R_a a + \int p(w) (T_s s + T_a a + w)^\top V_t (T_s s + T_a a + w) + q_t) dw \\ &= s^\top R_s s + s^\top T_s^\top V_t T_s s + \max_a (a^\top R_a a + 2s^\top T_s^\top V_t T_a a + a^\top T_a^\top V_t T_a a) + \int p(w) w^\top V_t w dw + q_t \end{aligned}$$

$a^*$  is where  $\nabla_a (\text{max term}) = 0$

$$0 = 2R_a a^* + 2T_a^\top V_t T_s s + 2T_a^\top V_t T_a a^*$$

$$a^* = -\underbrace{(R_a + T_a^\top V_t T_a)^{-1} T_a^\top V_t T_s s}_{K_t}$$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

Inductive step: show that if  $U_t^* = s^\top V_t s + q_t$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

$$\begin{aligned} U_{t+1}^*(s) &= \max_a (R(s, a) + \gamma E [U_t^*(s)]) \\ &= \max_a (s^\top R_s s + a^\top R_a a + \int p(w) U_t(T_s s + T_a a + w) dw) \\ &= s^\top R_s s + \max_a (a^\top R_a a + \int p(w) (T_s s + T_a a + w)^\top V_t (T_s s + T_a a + w) + q_t) dw \\ &= s^\top R_s s + s^\top T_s^\top V_t T_s s + \max_a (a^\top R_a a + 2s^\top T_s^\top V_t T_a a + a^\top T_a^\top V_t T_a a) + \int p(w) w^\top V_t w dw + q_t \end{aligned}$$

$a^*$  is where  $\nabla_a (\text{max term}) = 0$

$$0 = 2R_a a^* + 2T_a^\top V_t T_s s + 2T_a^\top V_t T_a a^*$$

$$a^* = -\underbrace{(R_a + T_a^\top V_t T_a)^{-1} T_a^\top V_t T_s s}_{K_t}$$

$$U_{t+1}^*(s) = \underbrace{s^\top (R_s + T_s^\top V_t T_s - (T_a^\top V_t T_s)^\top (R_a + T_a^\top V_t T_a)^{-1} (T_a^\top V_t T_s)) s}_{V_{t+1}} + \underbrace{\int p(w) w^\top V_t w dw + q_t}_{q_{t+1}}$$

# 1. Linear Dynamics, Quadratic Reward

We will show that  $U_h^*(s) = s^\top V_h s + q_h$  and  $\pi_h^*(s) = -K_h s$  by induction.

Base:  $U_1^*(s) = \max_a (s^\top R_s s + a^\top R_a a) = s^\top R_s s$

Inductive step: show that if  $U_t^* = s^\top V_t s + q_t$ , then  $U_{t+1}^* = s^\top V_{t+1} s + q_{t+1}$ .

$$\begin{aligned} U_{t+1}^*(s) &= \max_a (R(s, a) + \gamma E [U_t^*(s)]) \\ &= \max_a (s^\top R_s s + a^\top R_a a + \int p(w) U_t(T_s s + T_a a + w) dw) \\ &= s^\top R_s s + \max_a (a^\top R_a a + \int p(w) (T_s s + T_a a + w)^\top V_t (T_s s + T_a a + w) + q_t) dw \\ &= s^\top R_s s + s^\top T_s^\top V_t T_s s + \max_a (a^\top R_a a + 2s^\top T_s^\top V_t T_a a + a^\top T_a^\top V_t T_a a) + \int p(w) w^\top V_t w dw + q_t \end{aligned}$$

$a^*$  is where  $\nabla_a (\text{max term}) = 0$

$$0 = 2R_a a^* + 2T_a^\top V_t T_s s + 2T_a^\top V_t T_a a^*$$

$$a^* = -\underbrace{(R_a + T_a^\top V_t T_a)^{-1} T_a^\top V_t T_s s}_{K_t}$$

$$U_{t+1}^*(s) = s^\top \underbrace{\left( R_s + T_s^\top V_t T_s - (T_a^\top V_t T_s)^\top (R_a + T_a^\top V_t T_a)^{-1} (T_a^\top V_t T_s) \right)}_{V_{t+1}} s + \underbrace{\int p(w) w^\top V_t w dw + q_t}_{q_{t+1}}$$

$$U_{t+1}^*(s) = s^\top V_{t+1} s + q_{t+1} \quad \square$$

# 1. Linear Dynamics, Quadratic Reward

# 1. Linear Dynamics, Quadratic Reward

As  $h \rightarrow \infty$

# 1. Linear Dynamics, Quadratic Reward

As  $h \rightarrow \infty$

$$V_\infty = T_s^\top \left( V_\infty - V_\infty T_a \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty \right) T_s + R_s$$

# 1. Linear Dynamics, Quadratic Reward

As  $h \rightarrow \infty$

$$V_\infty = T_s^\top \left( V_\infty - V_\infty T_a \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty \right) T_s + R_s$$

$$K_\infty = \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty T_s$$

# 1. Linear Dynamics, Quadratic Reward

As  $h \rightarrow \infty$

$$V_\infty = T_s^\top \left( V_\infty - V_\infty T_a \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty \right) T_s + R_s$$

$$K_\infty = \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty T_s \quad \pi_\infty^*(s) = -K_\infty s$$

# 1. Linear Dynamics, Quadratic Reward

As  $h \rightarrow \infty$

$$V_\infty = T_s^\top \left( V_\infty - V_\infty T_a \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty \right) T_s + R_s$$

$$K_\infty = \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty T_s$$

$$\boxed{\pi_\infty^*(s) = -\underline{K}_\infty s}$$

( $K_\infty$  has no dependence on  $\Sigma$ )

# 1. Linear Dynamics, Quadratic Reward

As  $h \rightarrow \infty$

$$V_\infty = T_s^\top \left( V_\infty - V_\infty T_a \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty \right) T_s + R_s$$

$$K_\infty = \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty T_s \quad \pi_\infty^*(s) = -K_\infty s$$

( $K_\infty$  has no dependence on  $\Sigma$ )

Certainty-Equivalence Principle: For Linear-Quadratic problems, the optimal policy with noise is the same as the optimal policy without noise!

# 1. Linear Dynamics, Quadratic Reward

As  $h \rightarrow \infty$

$$V_\infty = T_s^\top \left( V_\infty - V_\infty T_a \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty \right) T_s + R_s$$

$$K_\infty = \left( T_a^\top V_\infty T_a + R_a \right)^{-1} T_a^\top V_\infty T_s \quad \pi_\infty^*(s) = -K_\infty s$$

( $K_\infty$  has no dependence on  $\Sigma$ )

Certainty-Equivalence Principle: For Linear-Quadratic problems, the optimal policy with noise is the same as the optimal policy without noise!



Practical Implication: If a continuous problem has roughly linear dynamics, a convex cost function, and roughly zero-mean additive noise, you can use *certainty-equivalent control*, i.e. control as if there is no noise.

# 2. Value Function Approximation

# 2. Value Function Approximation

$$V_\theta(s) = f_\theta(s) \quad (\text{e.g. neural network})$$

## 2. Value Function Approximation

$$V_\theta(s) = f_\theta(s) \quad (\text{e.g. neural network})$$

$$V_\theta(s) = \theta^\top \beta(s) \quad (\text{linear feature})$$

# 2. Value Function Approximation

$$V_\theta(s) = f_\theta(s) \quad (\text{e.g. neural network})$$

$$V_\theta(s) = \theta^\top \beta(s) \quad (\text{linear feature})$$

## Fitted Value Iteration

while not converged

$$\theta \leftarrow \theta'$$

$$\hat{V}' \leftarrow B_{\text{approx}}[V_\theta]$$

$$\theta' \leftarrow \text{fit}(\hat{V}')$$

# 2. Value Function Approximation

$$V_\theta(s) = f_\theta(s) \quad (\text{e.g. neural network})$$

$$V_\theta(s) = \theta^\top \beta(s) \quad (\text{linear feature})$$

## Fitted Value Iteration

while not converged

$$\theta \leftarrow \theta'$$

$$\hat{V}' \leftarrow B_{\text{approx}}[V_\theta]$$

$$\theta' \leftarrow \text{fit}(\hat{V}')$$

$$B_{\text{MC}(N)}[V_\theta](s) = \max_a \left( R(s, a) + \gamma \sum_{i=1}^N V_\theta(G(s, a, w_i)) \right)$$

# 2. Value Function Approximation

$$V_\theta(s) = f_\theta(s) \quad (\text{e.g. neural network})$$

$$V_\theta(s) = \theta^\top \beta(s) \quad (\text{linear feature})$$


## Fitted Value Iteration

while not converged

$$\theta \leftarrow \theta'$$

$$\hat{V}' \leftarrow B_{\text{approx}}[V_\theta]$$

$$\theta' \leftarrow \text{fit}(\hat{V}')$$



$$B_{\text{MC}(N)}[V_\theta](s) = \max_a \left( R(s, a) + \gamma \sum_{i=1}^N V_\theta(G(s, a, w_i)) \right)$$

# 2. Value Function Approximation

$$V_\theta(s) = f_\theta(s) \quad (\text{e.g. neural network})$$

$$V_\theta(s) = \theta^\top \beta(s) \quad (\text{linear feature})$$


## Fitted Value Iteration

while not converged

$$\theta \leftarrow \theta'$$



$$\hat{V}' \leftarrow B_{\text{approx}}[V_\theta]$$

$$\theta' \leftarrow \text{fit}(\hat{V}')$$




$$B_{\text{MC}(N)}[V_\theta](s) = \max_a \left( R(s, a) + \gamma \sum_{i=1}^N V_\theta(G(s, a, w_i)) \right)$$

## 2. Value Function Approximation




# Break

What will a Monte Carlo Tree Search tree look like if run on a problem with continuous spaces?




# 3. Sparse Tree Search/Progressive Widening


# 3. Sparse Tree Search/Progressive Widening




# 3. Sparse Tree Search/Progressive Widening





### 3. Sparse Tree Search/Progressive Widening



### 3. Sparse Tree Search/Progressive Widening




### 3. Sparse Tree Search/Progressive Widening




add new branch if  $C < \underline{kN^\alpha}$  ( $\alpha < 1$ )

number of children

### 3. Sparse Tree Search/Progressive Widening



add new branch if  $C < kN^\alpha$  ( $\alpha < 1$ )



# 4. Model Predictive Control

(Use off-the-shelf optimization software, e.g. Ipopt)

# 4. Model Predictive Control

(Use off-the-shelf optimization software, e.g. Ipopt)

Certainty-  
Equivalent

$$\begin{aligned} & \underset{a_{1:d}, s_{1:d}}{\text{maximize}} \quad \sum_{t=1}^d \gamma^t R(s_t, a_t) \\ & \text{subject to} \quad s_{t+1} = \mathbb{E}_{s' \sim T(s'|s, a)}[s'] \quad \forall t \end{aligned}$$

# 4. Model Predictive Control

(Use off-the-shelf optimization software, e.g. Ipopt)

Certainty-  
Equivalent

$$\begin{aligned} & \underset{a_{1:d}, s_{1:d}}{\text{maximize}} \quad \sum_{t=1}^d \gamma^t R(s_t, a_t) \\ & \text{subject to} \quad s_{t+1} = \mathbb{E}_{s' \sim T(s'|s, a)}[s'] \quad \forall t \end{aligned}$$

Open-Loop

$$\begin{aligned} & \underset{a_{1:d}, s_{1:d}^{(1:m)}}{\text{maximize}} \quad \frac{1}{m} \sum_{i=1}^m \sum_{t=1}^d \gamma^t R(s_t^{(i)}, a_t) \\ & \text{subject to} \quad s_{t+1} = G(s_t^{(i)}, a_t, w_t^{(i)}) \quad \forall t, i \end{aligned}$$

# 4. Model Predictive Control

(Use off-the-shelf optimization software, e.g. Ipopt)

Certainty-  
Equivalent

$$\begin{aligned} & \underset{a_{1:d}, s_{1:d}}{\text{maximize}} \quad \sum_{t=1}^d \gamma^t R(s_t, a_t) \\ & \text{subject to} \quad s_{t+1} = \mathbb{E}_{s' \sim T(s'|s,a)}[s'] \quad \forall t \end{aligned}$$

Open-Loop

$$\begin{aligned} & \underset{a_{1:d}, s_{1:d}^{(1:m)}}{\text{maximize}} \quad \frac{1}{m} \sum_{i=1}^m \sum_{t=1}^d \gamma^t R(s_t^{(i)}, a_t) \\ & \text{subject to} \quad s_{t+1} = G(s_t^{(i)}, a_t, w_t^{(i)}) \quad \forall t, i \end{aligned}$$

Hindsight  
Optimization

$$\begin{aligned} & \underset{a_{1:d}^{(1:m)}, s_{1:d}^{(1:m)}}{\text{maximize}} \quad \frac{1}{m} \sum_{i=1}^m \sum_{t=1}^d \gamma^t R(s_t^{(i)}, a_t^{(i)}) \\ & \text{subject to} \quad s_{t+1} = G(s_t^{(i)}, a_t^{(i)}, w_t^{(i)}) \quad \forall t, i \\ & \quad a_1^{(i)} = a_1^{(j)} \quad \forall i, j \end{aligned}$$

# Guiding Questions

- What tools do we have to solve MDPs with continuous  $S$  and  $A$ ?