

Probability and Random Variables

Concepts

1. Utility and Probability
2. Random Variables
3. Relationships between Random Variables

Utility and Probability

Full story: <https://projecteuclid.org/journals/statistical-science/volume-1/issue-3/The-Axioms-of-Subjective-Probability/10.1214/ss/1177013611.full>

Utility and Probability

Consider events A and B :

Full story: <https://projecteuclid.org/journals/statistical-science/volume-1/issue-3/The-Axioms-of-Subjective-Probability/10.1214/ss/1177013611.full>

Utility and Probability

Consider events A and B :

Utility indicates preference

Full story: <https://projecteuclid.org/journals/statistical-science/volume-1/issue-3/The-Axioms-of-Subjective-Probability/10.1214/ss/1177013611.full>

Utility and Probability

Consider events A and B :

Utility indicates preference

Probability indicates plausibility

Full story: <https://projecteuclid.org/journals/statistical-science/volume-1/issue-3/The-Axioms-of-Subjective-Probability/10.1214/ss/1177013611.full>

Utility and Probability

Consider events A and B :

Utility indicates preference

$U(A) > U(B)$ Indicates A is *preferable* to B

$U(A) = U(B)$ Indicates *indifference* between A and B

Probability indicates plausibility

Full story: <https://projecteuclid.org/journals/statistical-science/volume-1/issue-3/The-Axioms-of-Subjective-Probability/10.1214/ss/1177013611.full>

Utility and Probability

Consider events A and B :

Utility indicates preference

$U(A) > U(B)$ Indicates A is *preferable* to B

$U(A) = U(B)$ Indicates *indifference* between A and B

Probability indicates plausibility

$P(A) > P(B)$ Indicates A is *more plausible* (or likely) than B

$P(A) = P(B)$ Indicates A is *equally as plausible* (or equally likely) as B

What is a Random Variable?

R.V. X

Vocabulary/Notation

Vocabulary/Notation

Term

Definition

Coinflip Example

Vocabulary/Notation

Term	Definition	Coinflip Example
$\text{support}(X)$		

Vocabulary/Notation

Term	Definition	Coinflip Example
$\text{support}(X)$	All the values that X can take	

Vocabulary/Notation

Term	Definition	Coinflip Example
$\text{support}(X)$	All the values that X can take	
$x \in X$		

Vocabulary/Notation

Term	Definition	Coinflip Example
$\text{support}(X)$	All the values that X	$\{h, t\}$ or $\{0, 1\}$
$x \in X$	can take	"Binary random variable"

Vocabulary/Notation

Term	Definition	Coinflip Example
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$
$x \in X$		"Binary random variable"
Distribution		

Vocabulary/Notation

Term	Definition	Coinflip Example
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$ "Binary random variable"
$x \in X$		
Distribution	Maps each value in the support to a real number indicating its probability	

Vocabulary/Notation

Term	Definition	Coinflip Example
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$ "Binary random variable"
$x \in X$		
Distribution	Maps each value in the support to a real number indicating its probability	$\text{Bernoulli}(0.6)$ $P(X = 1) = 0.6$ $P(X = 0) = 0.4$

Vocabulary/Notation

Term	Definition	Coinflip Example							
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	"Binary random variable"						
$x \in X$									
Distribution	Maps each value in the support to a real number indicating its probability	$\text{Bernoulli}(0.6)$ $P(X = 1) = 0.6$ $P(X = 0) = 0.4$	$P(X)$ is a table <table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.4</td></tr><tr><td>1</td><td>0.6</td></tr></tbody></table>	x	P(x)	0	0.4	1	0.6
x	P(x)								
0	0.4								
1	0.6								

Vocabulary/Notation

Term	Definition	Coinflip Example							
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	"Binary random variable"						
Distribution	Maps each value in the support to a real number indicating its probability	$\text{Bernoulli}(0.6)$ $P(X = 1) = 0.6$ $P(X = 0) = 0.4$	$P(X)$ is a table <table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.4</td></tr><tr><td>1</td><td>0.6</td></tr></tbody></table>	x	P(x)	0	0.4	1	0.6
x	P(x)								
0	0.4								
1	0.6								
Expectation									

Vocabulary/Notation

Term	Definition	Coinflip Example							
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	"Binary random variable"						
Distribution	Maps each value in the support to a real number indicating its probability	$\text{Bernoulli}(0.6)$ $P(X = 1) = 0.6$ $P(X = 0) = 0.4$	$P(X)$ is a table <table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.4</td></tr><tr><td>1</td><td>0.6</td></tr></tbody></table>	x	P(x)	0	0.4	1	0.6
x	P(x)								
0	0.4								
1	0.6								
Expectation	First moment of the random variable, "mean"								

Vocabulary/Notation

Term	Definition	Coinflip Example							
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	"Binary random variable"						
Distribution	Maps each value in the support to a real number indicating its probability	$\text{Bernoulli}(0.6)$ $P(X = 1) = 0.6$ $P(X = 0) = 0.4$	$P(X)$ is a table <table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.4</td></tr><tr><td>1</td><td>0.6</td></tr></tbody></table>	x	P(x)	0	0.4	1	0.6
x	P(x)								
0	0.4								
1	0.6								
Expectation $E[X]$	First moment of the random variable, "mean"								

Vocabulary/Notation

Term	Definition	Coinflip Example							
$\text{support}(X)$ $x \in X$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$ "Binary random variable"	$P(X)$ is a table						
Distribution	Maps each value in the support to a real number indicating its probability	$\text{Bernoulli}(0.6)$ $P(X = 1) = 0.6$ $P(X = 0) = 0.4$	<table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.4</td></tr><tr><td>1</td><td>0.6</td></tr></tbody></table>	x	P(x)	0	0.4	1	0.6
x	P(x)								
0	0.4								
1	0.6								
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$							

Vocabulary/Notation

Term	Definition	Coinflip Example							
$\text{support}(X)$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	"Binary random variable"						
Distribution	Maps each value in the support to a real number indicating its probability	$\text{Bernoulli}(0.6)$ $P(X = 1) = 0.6$ $P(X = 0) = 0.4$	$P(X)$ is a table <table border="1"><thead><tr><th>x</th><th>P(x)</th></tr></thead><tbody><tr><td>0</td><td>0.4</td></tr><tr><td>1</td><td>0.6</td></tr></tbody></table>	x	P(x)	0	0.4	1	0.6
x	P(x)								
0	0.4								
1	0.6								
Expectation $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$							

Distributions of related R.V.s

Distributions of related R.V.s

Joint Distribution

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X \mid Y, Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	P(X, Y, Z)
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

X	P(X Y=1, Z=1)
0	0.84
1	0.16

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

X	$P(X Y=1, Z=1)$
0	0.84
1	0.16

Marginal Distribution

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

X	$P(X Y=1, Z=1)$
0	0.84
1	0.16

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X \mid Y, Z)$$

(Distribution - valued function)

X	$P(X \mid Y=1, Z=1)$
0	0.84
1	0.16

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

X	$P(X)$	Y	$P(Y)$
0	0.85	0	0.45
1	0.15	1	0.55

Z	$P(Z)$
0	0.20
1	0.80

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

3 Rules

Conditional Distribution

$$P(X \mid Y, Z)$$

(Burrito-level)

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

3 Rules

Conditional Distribution

$$P(X \mid Y, Z)$$

(Burrito-level)

Marginal Distribution

$$P(X) P(Y) P(Z)$$

(Gourmet Level: Axioms of Probability)

AXIOM 1. STRUCTURE OF UNKNOWN REAL NUMBERS AND PLAUSIBLE VALUE. We assume a set T of unknown numbers is a partially ordered commutative algebra over \mathbb{R} with identity, 1.

We assume in addition a given sub-Boolean algebra E of $E(T)$ with $0, 1 \in E$ and denote by E_0 the set of non-zero members of E . We assume that the partial ordering in $E(T)$ as a Boolean algebra coincides with the ordering that $E(T)$ inherits from the algebra T . Finally, we assume a function $PV : T \times E_0 \rightarrow \mathbb{R}$, called **PLAUSIBLE VALUE**, whose value on the pair (x, e) is denoted $PV(x|e)$.

AXIOM 2. STRONG RESCALING FOR PLAUSIBLE VALUE. If a, b belong to \mathbb{R} , if x belongs to T , and if e belongs to E_0 , then

$$PV(ax + b|e) = aPV(x|e) + b. \quad (2)$$

AXIOM 3. ORDER CONSISTENCY FOR PLAUSIBLE VALUE. If $x, y \in T$ and if $e \in E_0$, implies that $x \leq y$, then $PV(x|e) \leq PV(y|e)$.

Notice that if $e \in E(T)$, then $0 \leq e \leq 1$, in T , as it is true in the lattice ordering of $E(T)$.

AXIOM 4. THE COX AXIOM FOR PLAUSIBLE VALUE: If e, c are fixed in E , with $ec \in E_0$, if x_1, x_2 are in T , if $PV(x_1|ec) = PV(x_2|ec)$, then $PV(x_1e|c) = PV(x_2e|c)$. That is, we assume that as a function of x , the plausible value $PV(x|c)$ depends only on $PV(x|ec)$.

AXIOM 5. RESTRICTED ADDITIVITY OF PLAUSIBLE VALUE. For each fixed $y \in T$ and $e \in E_0$, the plausible value $PV(x+y|e)$ as a function of $x \in T$ depends only on $PV(x|e)$, which is to say that if $x_1, x_2 \in T$ and $PV(x_1|e) = PV(x_2|e)$, then $PV(x_1 + y|e) = PV(x_2 + y|e)$.

not on exam

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

3 Rules

Conditional Distribution

$$P(X \mid Y, Z)$$

(Burrito-level)

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

1)

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

(Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$

Distributions of related R.V.s

Joint Distribution	Conditional Distribution	Marginal Distribution
$P(X, Y, Z)$	$P(X Y, Z)$	$P(X) P(Y) P(Z)$
3 Rules	(Burrito-level)	

- 1) a) $0 \leq P(X | Y) \leq 1$
b) $\sum_{x \in X} P(x | Y) = 1$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$
- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

Distributions of related R.V.s

Joint Distribution	Conditional Distribution	Marginal Distribution
$P(X, Y, Z)$	$P(X Y, Z)$	$P(X) P(Y) P(Z)$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X | Y) \leq 1$
b) $\sum_{x \in X} P(x | Y) = 1$
- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

Joint \rightarrow Marginal

Distributions of related R.V.s

Joint Distribution	Conditional Distribution	Marginal Distribution
$P(X, Y, Z)$	$P(X Y, Z)$	$P(X) P(Y) P(Z)$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X | Y) \leq 1$
b) $\sum_{x \in X} P(x | Y) = 1$
- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X | Y) = \frac{P(X, Y)}{P(Y)}$$

Joint \rightarrow Marginal

Distributions of related R.V.s

Joint Distribution	Conditional Distribution	Marginal Distribution
$P(X, Y, Z)$	$P(X Y, Z)$	$P(X) P(Y) P(Z)$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X | Y) \leq 1$
b) $\sum_{x \in X} P(x | Y) = 1$
- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X | Y) = \frac{P(X, Y)}{P(Y)}$$

Joint \rightarrow Marginal

Joint + Marginal \rightarrow Conditional

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

(Burrito-level)

1) a) $0 \leq P(X \mid Y) \leq 1$

b) $\sum_{x \in X} P(x \mid Y) = 1$

2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint → Marginal

Joint + Marginal → Conditional

Marginal + Conditional → Joint

$$P(X, Y) = P(X|Y) P(Y)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Joint \rightarrow Marginal

Joint + Marginal \rightarrow Conditional

Marginal + Conditional \rightarrow Joint

$$P(X, Y) = P(X|Y) P(Y)$$

Naive Inference

(Book introduces unnormalized "factors", but process is the same.)

Naive Inference

Three Random Variables: A, B, C (Works for any number)

Naive Inference

Three Random Variables: A, B, C (Works for any number)

Want to find $P(\underbrace{A = a}_{\text{query}} \mid \underbrace{B = b}_{\text{evidence}})$

Naive Inference

Three Random Variables: A, B, C (Works for any number)

Want to find $P(\underbrace{A = a}_{\text{query}} \mid \underbrace{B = b}_{\text{evidence}})$

C is a "hidden variable"

Naive Inference

Three Random Variables: A, B, C (Works for any number)

Want to find $P(\underbrace{A = a}_{\text{query}} \mid \underbrace{B = b}_{\text{evidence}})$

C is a "hidden variable"

1. Determine the joint distribution $P(A, B, C)$.

Naive Inference

Three Random Variables: A, B, C (Works for any number)

Want to find $P(\underbrace{A = a}_{\text{query}} \mid \underbrace{B = b}_{\text{evidence}})$

C is a "hidden variable"

1. Determine the joint distribution $P(A, B, C)$.
2. Marginalize over hidden and query variables to get

$$P(A = a, B = b) = \sum_c P(A = a, B = b, C = c)$$

and

$$P(B = b) = \sum_{a,c} P(A = a, B = b, C = c)$$

(Book introduces unnormalized "factors", but process is the same.)

Naive Inference

Three Random Variables: A, B, C (Works for any number)

Want to find $P(\underbrace{A = a}_{\text{query}} \mid \underbrace{B = b}_{\text{evidence}})$

C is a "hidden variable"

1. Determine the joint distribution $P(A, B, C)$.
2. Marginalize over hidden and query variables to get

$$P(A = a, B = b) = \sum_c P(A = a, B = b, C = c)$$

and

$$P(B = b) = \sum_{a,c} P(A = a, B = b, C = c)$$

$$3. P(A = a \mid B = b) = \frac{P(A=a, B=b)}{P(B=b)}$$

(Book introduces unnormalized "factors", but process is the same.)

$$1) \text{ a)} 0 \leq P(X | Y) \leq 1$$

$$\text{b)} \sum_{x \in X} P(x | Y) = 1$$

$$2) P(X) = \sum_{y \in Y} P(X, y)$$

$$3) P(X | Y) = \frac{P(X, Y)}{P(Y)}$$

$$P(X, Y) = P(X|Y) P(Y)$$

Break

- $P \in \{0, 1\}$: Powder Day
- $C \in \{0, 1\}$: Pass Clear
- 1 in 5 days is a powder day
- The pass is clear 8 in 10 days
- If it is a powder day, there is a 50% chance the pass is blocked

- Write out the joint probability distribution for P and C.
- Suppose it is a non-powder day, what is the probability that the pass is blocked?

Bayes Rule

- Know: $P(B | A)$, $P(A)$, $P(B)$
- Want: $P(A | B)$

Definitions: Conditional Expectation and Independence

Definitions: Conditional Expectation and Independence

Definition: The conditional expectation of X given Y is

$$E[X | Y] = \sum_x x P(X = x | Y)$$

(function from values of Y to expectations of X)

Definitions: Conditional Expectation and Independence

Definition: The conditional expectation of X given Y is

$$E[X | Y] = \sum_x x P(X = x | Y)$$

(function from values of Y to expectations of X)

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

Definitions: Conditional Expectation and Independence

Definition: The conditional expectation of X given Y is

$$E[X | Y] = \sum_x x P(X = x | Y)$$

(function from values of Y to expectations of X)

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

Definitions: Conditional Expectation and Independence

Definition: The conditional expectation of X given Y is

$$E[X | Y] = \sum_x x P(X = x | Y)$$

(function from values of Y to expectations of X)

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

Definitions: Conditional Expectation and Independence

Definition: The conditional expectation of X given Y is

$$E[X | Y] = \sum_x x P(X = x | Y)$$

(function from values of Y to expectations of X)

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X | Y) = P(X)$$

Definition: X and Y are *conditionally independent* given Z iff

$$P(X, Y | Z) = P(X | Z) P(Y | Z)$$

Definitions: Conditional Expectation and Independence

Definition: The conditional expectation of X given Y is

$$E[X | Y] = \sum_x x P(X = x | Y)$$

(function from values of Y to expectations of X)

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

Definition: X and Y are *conditionally independent* given Z iff

$$P(X, Y | Z) = P(X | Z) P(Y | Z)$$

$$X \perp Y | Z$$

Concepts

1. Utility and Probability
2. Random Variables
3. Relationships between Random Variables