

Operational Definition of the Phoenix Sepsis Score

July 23, 2026

Table of contents

1	Objective	3
2	Notation	3
2.1	Symbols	3
2.2	Function Definitions	5
3	Defining Organ System Dysfunction Scores	6
3.1	Respiratory	6
3.1.1	Data from the EHR	6
3.1.2	Constructed Variables	7
3.1.3	Respiratory Organ Dysfunction Score	8
3.2	Cardiovascular	8
3.2.1	Data from the EHR	9
3.2.2	Constructed Variable	9
3.2.3	Cardiovascular Dysfunction Score	11
3.3	Neurological	11
3.3.1	Data from the EHR	11
3.3.2	Constructed Variable	11
3.3.3	Neurological Organ Dysfunction Score	12
3.4	Coagulation	12
3.4.1	Data from the EHR	12
3.4.2	Constructed Variable	13
3.4.3	Coagulation Organ Dysfunction Score	13
3.5	Endocrine	13
3.5.1	Data from the EHR	13
3.5.2	Constructed Variables	13
3.5.3	Endocrine Organ Dysfunction Score	13
3.6	Immunologic	14
3.6.1	Data from the EHR	14
3.6.2	Constructed Variables	14
3.6.3	Immunologic Organ Dysfunction Score	14
3.7	Hepatic	14
3.7.1	Data from the EHR	14
3.7.2	Constructed Variables	14
3.7.3	Hepatic Organ Dysfunction Score	15
3.8	Renal	15
3.8.1	Data from the EHR	15
3.8.2	Constructed Variables	15
3.8.3	Renal Organ Dysfunction Score	15

3.9 Suspected Infection	16
4 Phoenix Sepsis Score	16
5 Exploratory Aggregation Schemes	17
5.1 Organ-Level Maxima	17
5.2 Organ-Level + Cardiovascular Component Decoupling	18
5.3 Full Component Decoupling	18

List of Tables

1 Notation used in the operational definition of the Phoenix Sepsis Score.	3
2 Logic with NA.	5
3 EHR Inputs for Respiratory Organ Dysfunction Scoring	6
4 EHR Inputs for Cardiovascular Organ Dysfunction Scoring	9
5 EHR Inputs for Neurological Organ Dysfunction Scoring	11
6 EHR Inputs for Coagulation Organ Dysfunction Scoring	13
7 EHR Inputs for Endocrine Organ Dysfunction Scoring	13
8 EHR Inputs for Immunologic Organ Dysfunction Scoring	14
9 EHR Inputs for Hepatic Organ Dysfunction Scoring	14
10 EHR Inputs for Renal Organ Dysfunction Scoring	15
11 Parameter values defining time windows, look-backs, and thresholds for the Phoenix Sepsis Score, Sepsis, and Septic Shock.	17

1 Objective

Provide formal notation for the operational definition of the Phoenix Sepsis Score as published in [Sanchez-Pinto et al. \[2024\]](#) and [Schlapbach et al. \[2024\]](#) along with several exploratory aggregation schemes. The different scoring methods are based on when and how data is aggregated. The differences in the aggregation methods can dramatically change the Phoenix score and the published cut point of two or more points for sepsis may not be applicable to all aggregations.

2 Notation

2.1 Symbols

This section is a quick guide to the symbols used throughout the document. Detailed definitions for the functions will follow in § 2.2

Table 1: Notation used in the operational definition of the Phoenix Sepsis Score.

Symbol	Meaning	Example or Notes
Sets and Membership		
$\{\cdot\}$	Set notation used to define collections of values satisfying a condition.	$\{x : x \leq t\}$ denotes the set of values x less than or equal to t . Also, $A = \{1, 3, 5\}$ is read as "the set A consisting of the elements 1, 3, and 5."
$\emptyset$	Empty set.	Represents a set containing no elements.
$\in$	Set membership operator.	$t \in T$ indicates that time t lies within the time window T .
$A \setminus B$	Remove elements of a set	Example $\{1, 2, 3\} \setminus \{2\} = \{1, 3\}$
Intervals		
(a, b)	Open interval.	$X \in (a, b) \equiv a < X < b$.
$[a, b)$	Left-closed, right-open interval.	$X \in [a, b) \equiv a \leq X < b$.
$(a, b]$	Left-open, right-closed interval.	$X \in (a, b] \equiv a < X \leq b$.
$[a, b]$	Closed interval.	$X \in [a, b] \equiv a \leq X \leq b$.
Definitions, Equality, and Logic		
$:=$	Definition operator indicating that the quantity on the left is defined by the expression on the right.	$f(x) := x^2$ defines the function $f(x)$ as x^2 .
$=$	Equality relation.	$2 + 3 = 5$.
$\equiv$	Logical equivalence or identity.	$A \equiv B$ means that A and B are equivalent statements.
$\vee$	Logical OR operator.	$A \vee B$ is true if either A or B is true.
$\wedge$	Logical AND operator.	$A \wedge B$ is true if both A and B are true.
$\forall$	Universal quantifier ("for all").	$\forall t \in T$ means for every t in the time window T .
$\exists$	Existential quantifier ("there exists").	$\exists t \in T$ means for some time t in T .
Arithmetic and Functional Operators		

Continued on next page

Table 1 – continued from previous page

Symbol	Meaning	Example or Notes
$\sum$	Summation operator.	$\sum_{i=1}^4 A_i = A_1 + A_2 + A_3 + A_4$.
$\max$	Maximum operator.	$\max_{t \in T} f(t)$ evaluates $f(t)$ over all $t \in T$ and returns the maximum value. $\max \mathbf{A}$ returns the largest value of the set $\mathbf{A}$, e.g., $\max \{1, 2, 3\} = 3$.
$\min$	Minimum operator.	$\min_{t \in T} f(t)$ evaluates $f(t)$ over all $t \in T$ and returns the minimum value. $\min \mathbf{A}$ returns the smallest value of the set $\mathbf{A}$, e.g., $\min \{1, 2, 3\} = 1$.
$\times$	Multiplication operator.	
$\mathbb{I}\{A\}$	Indicator function returning 1 if condition A is true and 0 otherwise.	Defined in (4).
Time and Indexing		
t	Time measured in minutes relative to encounter start.	Negative values denote pre-encounter measurements, $t = 0$ marks encounter start, and positive values occur during the encounter.
T	Time window measured in minutes.	$T = [0, 1440)$ represents the first 24 hours following encounter start.
m_{resp}	Look-back window for respiratory related values	$[0, \infty)$
m_{vaso}	Look-back window for vasoactive medications	$[0, \infty)$
m_{bp}	Look-back window for blood pressure measurements	$[0, \infty)$
m_{lac}	Look-back window for lactate	$[0, \infty)$
m_{gcs}	Look-back window for GCS	$[0, \infty)$
m_{pupil}	Look-back window for Pupils	$[0, \infty)$
m_{coag}	Look-back window for coagulation related values	$[0, \infty)$
m_{endo}	Look-back window for endocrine related values	$[0, \infty)$
m_{immu}	Look-back window for immunologic related values	$[0, \infty)$
m_{hepatic}	Look-back window for hepatic related values	$[0, \infty)$
m_{renal}	Look-back window for renal related values	$[0, \infty)$
$\mathbf{m}$	Vector of look-back windows used for carry-forward operations.	
Observation History and Carry-Forward		
$\tau_X(t)$	Most recent observation time for variable X at or before time t .	$\tau_X(t)$ identifies when X was last measured prior to t . Defined in (1).
$\Delta_X(t)$	Staleness of measurement for variable X .	$\Delta_X(t)$ is the elapsed time since $\tau_X(t)$. Defined in (2).

Continued on next page

Table 1 – continued from previous page

Symbol	Meaning	Example or Notes
$\mathcal{C}[X](t; m)$	Carry-forward operator returning the most recent value of X within m minutes of time t .	Returns the last observed value of X if $\Delta_X(t) \leq m$, otherwise NA. Defined in (3).
$\mathcal{D}[x](t; \cdot)$	Scoring operator: the point contribution of scoring entity x at time t .	When the common argument list (m, Age) is used, arguments not relevant to x are ignored.
NA	Missing, unknown, or undefined value.	Returned when a required measurement is unavailable.

Two points are especially important for reading the rest of the document. First, $:=$ is **not** the same as $=$. $:=$ means we are *defining* something. Second, NA means “missing/unknown/undefined” (e.g., a lab value was not measured), which is different from a numeric value like 0. By contrast, $\emptyset$ is the **empty set** (no elements), which we use in set-based definitions like $\max \emptyset = \text{NA}$.

For arithmetic, when NA is involved, the outcome is always NA, e.g., $\text{NA} + X := \text{NA} \forall X$. Table 2 explicitly defines how logical statements are interpreted when working with NA. When a value is missing, we avoid forcing it into “true” or “false.” The rules below show how missingness propagates through logical statements. If either side of an “and/or” statement is unknown, the whole statement may be unknown unless the other side already determines the answer.

Table 2: Logic with NA.

Statement	Value
‘OR’ statements	
$\text{NA} \vee \text{NA}$	NA
$\text{NA} \vee \text{true}$	true
$\text{NA} \vee \text{false}$	NA
‘And’ statements	
$\text{NA} \wedge \text{NA}$	NA
$\text{NA} \wedge \text{true}$	NA
$\text{NA} \wedge \text{false}$	false

2.2 Function Definitions

This section defines how we turn time-stamped clinical data into a single value at any time t . In real EHR data, measurements are taken at irregular times. The definitions below formalize “what is the most recent value?” and “is it too old to trust?” so that scoring is reproducible.

For any time-indexed physiologic variable X , we write $X(t)$ for the value recorded at time t where t is the time, in minutes from the encounter start, i.e., $t = 0$ denotes encounter start, $t = 32$ is 32 minutes into the encounter, and $t = -12$ is 12 minutes before encounter start. The negative times represent measurements recorded before encounter start, such as EMS data.

For any variable X , define the last observation time prior to or at t as:

$$\tau_X(t) := \max \{s \leq t : X(s) \text{ observed or reported}\}, \quad \max \emptyset = \text{NA}. \quad (1)$$

Define the staleness of the measurement:

$$\Delta_X(t) := \begin{cases} t - \tau_X(t) & \tau_X(t) \neq \text{NA} \\ \text{NA} & \text{otherwise.} \end{cases} \quad (2)$$

We define a staleness-limited carry-forward operator. Carry forward when working with retrospective data, a look back limit when working with prospective or real time data. For $m \geq 0$ minutes,

$$\mathcal{C}[X](t; m) := \begin{cases} X(\tau_X(t)) & \Delta_X(t) \leq m \\ \text{NA} & \text{otherwise.} \end{cases} \quad (3)$$

We will use an indicator function that returns 1 or 0:

$$\mathbb{I}\{A\} := \begin{cases} 1 & A \text{ is true} \\ 0 & A \text{ is false or NA.} \end{cases} \quad (4)$$

In words: $\tau_X(t)$ tells you when the last measurement happened, $\Delta_X(t)$ tells you how old it is, and $\mathcal{C}[X](t; m)$ returns that last value if it is recent enough (within m minutes); otherwise it returns “missing.” This is how we prevent using values that are too old to reflect current physiology. The indicator $\mathbb{I}\{A\}$ then lets us convert logical statements (e.g., “lactate is at least 5”) into 0/1 points for scoring.

For potentially incomplete time-indexed function f , extrema are evaluated over the available values of f . Define:

$$\mathcal{A}_f(T) := \{f(t) : t \in T, f(t) \text{ is available}\}. \quad (5)$$

Then

$$\max_{t \in T} f(t) := \begin{cases} \max \mathcal{A}_f(T) & \mathcal{A}_f(T) \neq \emptyset \\ \text{NA} & \mathcal{A}_f(T) = \emptyset, \end{cases} \quad (6)$$

and

$$\min_{t \in T} f(t) := \begin{cases} \min \mathcal{A}_f(T) & \mathcal{A}_f(T) \neq \emptyset \\ \text{NA} & \mathcal{A}_f(T) = \emptyset. \end{cases} \quad (7)$$

3 Defining Organ System Dysfunction Scores

We assess eight organ systems: respiratory, cardiovascular, neurologic, coagulation, endocrine, immunologic, hepatic, and renal. Each subsection below lists the EHR inputs needed to compute that organ score, then shows how those inputs are transformed into constructed variables and point assignments. The goal is to make every step of the scoring reproducible: given the same time-stamped data, you should arrive at the same score.

The listed patient, laboratory, observation, event, and medication fields are as we expect them to appear in the electronic health record. These are inputs, not judgments about clinical severity. We define a domain for the variable. These domains are intentionally broad (wider than clinically plausible values) to keep data validation separate from scoring.

3.1 Respiratory

3.1.1 Data from the EHR

There are several values we expect to get from the Electronic Health Record capturing oxygenation, FiO_2 (fraction of inspired oxygen), SpO_2 (pulse oximetry oxygen saturation), and PaO_2 (arterial oxygen pressure). Additionally, the presence of mechanical or noninvasive respiratory support. Together, they allow us to quantify how well the patient is oxygenating and whether that oxygenation depends on advanced support.

Table 3: EHR Inputs for Respiratory Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
FiO_2	Fraction of Inspired Oxygen	Real	unitless	[0.21, 1.00]

Table 3 – continued from previous page

Variable	Description	Type	Units	Domain
SpO ₂	Pulse oximetry oxygen saturation	Integer	unitless	$\{0, 1, \dots, 100\}$
PaO ₂	Arterial oxygen pressure	Real	mmHg	$[0, \infty)$
P _{aw} ^{VENT}	Mean Airway Pressure, from a ventilator	Real	cm H ₂ O	$[0, \infty)$
P _{aw} ^{HFOV}	Mean Airway Pressure, from a HFOV	Real	cm H ₂ O	$[0, \infty)$
P _{aw} ^{PEEP}	Positive end-expiratory pressure	Real	cm H ₂ O	$[0, \infty)$
VENT	Indicator for invasive mechanical ventilation	Integer		$\{0, 1\}$
O ₂ Support	Any non-invasive oxygen support	Integer		$\{0, 1\}$

3.1.2 Constructed Variables

From the EHR values we construct several variables which are then used for the respiratory score. The constructed variables combine measurements that are not always recorded at the same time (e.g., SpO₂ and FiO₂) and apply clinical logic about timing and support. This keeps us from pairing an outdated FiO₂ with a more recent oxygen measurement or vice versa.

PF Ratio For determining the PF ratio at time t we consider look-back windows for both FiO₂ and PaO₂ and the relative look-back between the two. Specifically, the PFR is only valid when

$$\tau_{\text{FiO}_2}(t) \leq \tau_{\text{PaO}_2}(t) \quad \equiv \quad \Delta_{\text{FiO}_2}(t) \geq \Delta_{\text{PaO}_2}(t). \quad (8)$$

Conceptually, this restriction is in place to protect from using a PaO₂ value which was taken *before* the most current FiO₂ value. That is, PFR is only valid if the PaO₂ value is reported at the same time, or after the FiO₂ is reported.

The PFR at time t is:

$$\text{PFR}(t; m_{\text{resp}}) := \begin{cases} \frac{\mathcal{C}[\text{PaO}_2](t; m_{\text{resp}})}{\mathcal{C}[\text{FiO}_2](t; m_{\text{resp}})} & \tau_{\text{FiO}_2}(t) \leq \tau_{\text{PaO}_2}(t) \\ \text{NA} & \text{otherwise.} \end{cases} \quad (9)$$

With this definition, at time t , the PF ratio will either have a valid numeric value or be NA.

SF Ratio Like the PF ratio, we only consider the SF ratio valid if the SpO₂ value is reported at the same time as the FiO₂ or after.

$$\tau_{\text{FiO}_2}(t) \leq \tau_{\text{SpO}_2}(t) \quad \equiv \quad \Delta_{\text{FiO}_2}(t) \geq \Delta_{\text{SpO}_2}(t). \quad (10)$$

Additionally, the SF ratio is valid if the SpO₂ ≤ 97 . This requirement is based on the PODIUM [Bembea et al., 2022, Squires et al., 2022] organ dysfunction scoring, and is an additional requirement on the pSOFA [Matics and Sanchez-Pinto, 2017] cut points in the scoring.

$$\text{SFR}(t; m_{\text{resp}}) := \begin{cases} \frac{\mathcal{C}[\text{SpO}_2](t; m_{\text{resp}})}{\mathcal{C}[\text{FiO}_2](t; m_{\text{resp}})} & \tau_{\text{FiO}_2}(t) \leq \tau_{\text{SpO}_2}(t) \wedge \mathcal{C}[\text{SpO}_2](t; m_{\text{resp}}) \leq 97 \\ \text{NA} & \text{otherwise.} \end{cases} \quad (11)$$

Invasive Mechanical Ventilation We operationalize the definition for invasive mechanical ventilation based in part of differences in the EHR records we had access to. In some sets, a simple VENT indicator variable was provided. In other EHR sets we had mean airway pressure from either a ventilator or high-frequency oscillatory ventilation (HFOV). Additionally, we had positive end-expiratory pressure (PEEP) to consider as well.

We defined invasive mechanical ventilation (IMV) as an indicator based on four possible conditions.

$$\text{IMV}(t; m_{\text{resp}}) := \mathbb{I} \left\{ \bigvee_{i=1}^4 A_i(t; m_{\text{resp}}) \right\} \quad (12)$$

where

$$\begin{aligned} A_1(t; m_{\text{resp}}) &:= \mathcal{C}[\text{VENT}](t; m_{\text{resp}}) = 1, \\ A_2(t; m_{\text{resp}}) &:= \mathcal{C}[\text{P}_{\text{aw}}^{\text{VENT}}](t; m_{\text{resp}}) > 0, \\ A_3(t; m_{\text{resp}}) &:= \mathcal{C}[\text{P}_{\text{aw}}^{\text{HFOV}}](t; m_{\text{resp}}) > 0, \text{ and} \\ A_4(t; m_{\text{resp}}) &:= \mathcal{C}[\text{P}_{\text{aw}}^{\text{PEEP}}](t; m_{\text{resp}}) > 3. \end{aligned} \quad (13)$$

The PEEP limit of 3 cm H₂O is used to limit false positives while limiting false negatives. Similar limits have been used in other criteria. For example, the lung injury score [Martin and Bernard, 2001] uses a PEEP of 5 cm H₂O for 0 points and 6-8 cm H₂O. Further, Martin and Bernard [2001] states a PEEP of 5 cm H₂O and FiO₂ < 0.5 as part of the criteria for identifying patients ready to accept a trial of spontaneous breathing.

Other Respiratory Support Extend the definitions in (13) to include

$$\begin{aligned} A_5(t; m_{\text{resp}}) &:= \mathcal{C}[\text{FiO}_2](t; m_{\text{resp}}) > 0.21, \\ A_6(t; m_{\text{resp}}) &:= \mathcal{C}[\text{O}_2\text{Support}](t; m_{\text{resp}}) = 1. \end{aligned} \quad (14)$$

Then other respiratory support is defined by (13) and (14) as

$$\text{ORS}(t; m_{\text{resp}}) := \mathbb{I} \left\{ \bigvee_{i=1}^6 A_i(t; m_{\text{resp}}) \right\} \quad (15)$$

3.1.3 Respiratory Organ Dysfunction Score

In the development of Phoenix, we used pSOFA respiratory scoring [Matics and Sanchez-Pinto, 2017]. The pSOFA respiratory scoring does not explicitly require respiratory support to score 1 point, however, for the Phoenix definition the requirement of some respiratory support was made explicit. It is extremely unlikely that a patient with SFR or PFR values sufficiently low enough to score at least one point for respiratory dysfunction would not have at the very least supplemental oxygen.

$$\mathcal{D}[\text{Resp}](t; m_{\text{resp}}) := \text{ORS}(t; m_{\text{resp}}) \mathbb{I}\{B_1(t; m_{\text{resp}})\} + \text{IMV}(t; m_{\text{resp}}) \sum_{i=2}^3 \mathbb{I}\{B_i(t; m_{\text{resp}})\} \quad (16)$$

where

$$\begin{aligned} B_1(t; m_{\text{resp}}) &:= \text{PFR}(t; m_{\text{resp}}) < 400 \vee \text{SFR}(t; m_{\text{resp}}) < 292 \\ B_2(t; m_{\text{resp}}) &:= \text{PFR}(t; m_{\text{resp}}) < 200 \vee \text{SFR}(t; m_{\text{resp}}) < 220 \\ B_3(t; m_{\text{resp}}) &:= \text{PFR}(t; m_{\text{resp}}) < 100 \vee \text{SFR}(t; m_{\text{resp}}) < 148. \end{aligned} \quad (17)$$

The domain for the respiratory organ dysfunction score is

$$\mathcal{D}[\text{Resp}](t; m_{\text{resp}}) \in \{0, 1, 2, 3\} \quad \forall m_{\text{resp}}, t. \quad (18)$$

3.2 Cardiovascular

The cardiovascular score has three components, vasoactive medications, mean arterial pressure, and lactate. A score of 0, 1, or 2 points is possible for each of the components resulting in a total score of 0 to 6 points.

3.2.1 Data from the EHR

The input data from the EHR needed for the cardiovascular dysfunction scores are vasoactive medications (as indicators of pharmacologic cardiovascular support), blood pressure measurements (from cuff or arterial line), lactate, and age (for age-specific mean arterial pressure thresholds).

Table 4: EHR Inputs for Cardiovascular Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
Dobutamine	Indicator of systemic use of vasoactive medication	Integer	–	$\{0, 1\}$
Dopamine	Indicator of systemic use of vasoactive medication	Integer	–	$\{0, 1\}$
Epinephrine	Indicator of systemic use of vasoactive medication	Integer	–	$\{0, 1\}$
Milrinone	Indicator of systemic use of vasoactive medication	Integer	–	$\{0, 1\}$
Norepinephrine	Indicator of systemic use of vasoactive medication	Integer	–	$\{0, 1\}$
Vasopressin	Indicator of systemic use of vasoactive medication	Integer	–	$\{0, 1\}$
Lactate	Lactate	Real	mmol / L	$[0, \infty)$
SBP _c	systolic blood pressure reported from a cuff	Real	mmHg	$[0, \infty)$
DBP _c	diastolic blood pressure reported from a cuff	Real	mmHg	$[1, 200]$
MAP _c	mean arterial pressure reported from a cuff	Real	mmHg	$[0, \infty)$
SBP _a	systolic blood pressure reported from arterial line	Real	mmHg	$[0, \infty)$
DBP _a	diastolic blood pressure reported from arterial line	Real	mmHg	$[1, 200]$
MAP _a	mean arterial pressure reported from arterial line	Real	mmHg	$[0, \infty)$
Age	Patient’s age	Real	months	$[0, 216]$

3.2.2 Constructed Variable

We build three components: vasoactive medication use, mean arterial pressure, and lactate. Each component contributes 0–2 points, and together they sum to the cardiovascular score. The definitions below formalize how we combine multiple data sources (e.g., cuff vs arterial line) and how we handle timing.

Vasoactive Medications Each of the six vasoactive medications

$$V := \{\text{Dobutamine, Dopamine, Epinephrine, Milrinone, Norepinephrine, Vasopressin}\} \quad (19)$$

are to be assessed as 0/1 indicators. Either the patient is receiving systemic vasoactive medication for the purpose of cardiovascular support or they are not. Medications such as epinephrine are used in many different situations for different purposes, it is important that the input data is an indicator of the use of the medication systemically for cardiovascular support.

We define the contribution to the cardiovascular score from vasoactive medications as the count of the number of vasoactive medications received, regardless of dose, with a cap of 2 points.

$$\mathcal{D}[\text{Vasos}](t; m_{\text{vaso}}) := \min \left\{ 2, \sum_{v \in V} \mathbb{I} \{ \mathcal{C}[v](t; m_{\text{vaso}}) = 1 \} \right\}. \quad (20)$$

The domain for the contribution to the cardiovascular score from vasoactive medications is:

$$\mathcal{D}[\text{Vasos}](t; m_{\text{vaso}}) \in \{0, 1, 2\} \quad \forall m_{\text{vaso}}, t \quad (21)$$

This was based on the use of the vasoactive inotrope score (VIS)[[Haque et al., 2015](#)]. The VIS used the same six vasoactive medications and based the score on the dosage. During the development of the Phoenix criteria we found some lower resourced sites did not have all six medications. As such, patients at these

lower resourced sites had a ceiling on the severity of cardiovascular dysfunction based on VIS. By ignoring the dosage and just relying on the fact that the patient was receiving the medication, the Phoenix criteria was determined to have greater utility across high-, medium-, and low- resourced sites.

Mean Arterial Pressure (MAP) Mean arterial pressure (MAP) is non-trivial to define. We have limits based on age (in months) and a hierarchy for the preferable source of information used to define the MAP. In practice, MAP may be recorded directly or inferred from systolic and diastolic pressures. We prioritize arterial line measurements when available because they are more reliable in unstable patients, then fall back to cuff measurements.

The hierarchy for the pressure: use the MAP from an arterial line (MAP_a) first. If that is not available, then approximate using systolic and diastolic values reported from the arterial line. SBP_a and DBP_a . If none of the arterial line values are available, then use values from a cuff preferentially taking a reported mean arterial pressure, MAP_c , and lastly, approximate via SBP_c and DBP_c . That is, let

$$\begin{aligned}\widehat{\text{MAP}}_1(t; m_{\text{bp}}) &:= \mathcal{C}[\text{MAP}_a](t; m_{\text{bp}}), \\ \widehat{\text{MAP}}_2(t; m_{\text{bp}}) &:= \frac{1}{3}\mathcal{C}[\text{SBP}_a](t; m_{\text{bp}}) + \frac{2}{3}\mathcal{C}[\text{DBP}_a](t; m_{\text{bp}}), \\ \widehat{\text{MAP}}_3(t; m_{\text{bp}}) &:= \mathcal{C}[\text{MAP}_c](t; m_{\text{bp}}), \text{ and} \\ \widehat{\text{MAP}}_4(t; m_{\text{bp}}) &:= \frac{1}{3}\mathcal{C}[\text{SBP}_c](t; m_{\text{bp}}) + \frac{2}{3}\mathcal{C}[\text{DBP}_c](t; m_{\text{bp}}).\end{aligned}\tag{22}$$

Then

$$\text{MAP}^*(t; m_{\text{bp}}) := \begin{cases} \widehat{\text{MAP}}_{i^*}(t; m_{\text{bp}}) & \text{if } \exists i^* \in \{1, 2, 3, 4\} \text{ where} \\ & i^* := \min \{i \in \{1, 2, 3, 4\} : \widehat{\text{MAP}}_i(t; m_{\text{bp}}) \neq \text{NA}\} \\ \text{NA} & \text{otherwise.} \end{cases}\tag{23}$$

The limits for the MAP are based on age (in months). Age-based thresholds reflect pediatric physiology: younger patients have lower expected MAPs, so the cut-points increase with age. Let

$$\theta_1(\text{Age}) := \begin{cases} 31 & \text{Age} \in [0, 1) \\ 39 & \text{Age} \in [1, 12) \\ 44 & \text{Age} \in [12, 24) \\ 45 & \text{Age} \in [24, 60) \\ 49 & \text{Age} \in [60, 144) \\ 52 & \text{Age} \in [144, 216), \end{cases}\tag{24}$$

and

$$\theta_2(\text{Age}) := \begin{cases} 17 & \text{Age} \in [0, 1) \\ 25 & \text{Age} \in [1, 12) \\ 31 & \text{Age} \in [12, 24) \\ 32 & \text{Age} \in [24, 60) \\ 36 & \text{Age} \in [60, 144) \\ 38 & \text{Age} \in [144, 216). \end{cases}\tag{25}$$

Then, the part of the cardiovascular score based on MAP is:

$$\mathcal{D}[\text{MAP}](t; m_{\text{bp}}, \text{Age}) := \mathbb{I}\{\text{MAP}^*(t; m_{\text{bp}}) < \theta_2(\text{Age})\} + \mathbb{I}\{\text{MAP}^*(t; m_{\text{bp}}) < \theta_1(\text{Age})\}.\tag{26}$$

The domain for this component of the cardiovascular dysfunction score is

$$\mathcal{D}[\text{MAP}](t; m_{\text{bp}}, \text{Age}) \in \{0, 1, 2\} \quad \forall m_{\text{bp}}, t.\tag{27}$$

Lactate Lactate is treated as a marker of impaired perfusion and severity of shock. Higher lactate thresholds contribute more points. The definition for lactate used in the cardiovascular dysfunction score is:

$$\mathcal{D}[\text{Lactate}](t; m_{\text{lac}}) := \mathbb{I}\{\mathcal{C}[\text{Lactate}](t; m_{\text{lac}}) \geq 5\} + \mathbb{I}\{\mathcal{C}[\text{Lactate}](t; m_{\text{lac}}) \geq 11\}. \quad (28)$$

That is, one point for a lactate value of at least 5 mmol/L; two points for a lactate value of at least 11 mmol/L. The domain for the contribution of lactate to the cardiovascular dysfunction score is:

$$\mathcal{D}[\text{Lactate}](t; m_{\text{lac}}) \in \{0, 1, 2\} \quad \forall m_{\text{lac}}, t. \quad (29)$$

3.2.3 Cardiovascular Dysfunction Score

The cardiovascular dysfunction is based on (20), (26), and (28).

$$\mathcal{D}[\text{Card}](t; m_{\text{vaso}}, m_{\text{bp}}, m_{\text{lac}}, \text{Age}) := \mathcal{D}[\text{Vasos}](t; m_{\text{vaso}}) + \mathcal{D}[\text{MAP}](t; m_{\text{bp}}, \text{Age}) + \mathcal{D}[\text{Lactate}](t; m_{\text{lac}}). \quad (30)$$

The domain for the cardiovascular dysfunction score is

$$\mathcal{D}[\text{Card}](t; m_{\text{vaso}}, m_{\text{bp}}, m_{\text{lac}}, \text{Age}) \in \{0, 1, 2, 3, 4, 5, 6\} \quad \forall m, t; \quad (31)$$

3.3 Neurological

The neurological score is based on the Glasgow Coma score (GCS) and whether or not the patient has bilaterally fixed pupils. These capture level of consciousness and brainstem function, respectively.

3.3.1 Data from the EHR

Table 5: EHR Inputs for Neurological Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
GCS _{motor}	Glasgow Coma Motor Score	Integer	unitless	{1, 2, 3, 4, 5, 6}
GCS _{eye}	Glasgow Coma Eye Score	Integer	unitless	{1, 2, 3, 4}
GCS _{verbal}	Glasgow Coma Verbal Score	Integer	unitless	{1, 2, 3, 4, 5}
GCS _{total}	Total Glasgow Coma Score	Integer	unitless	{3, 4, ..., 15}
Pupils	Bilaterally fixed pupils	Integer		{0, 1}

3.3.2 Constructed Variable

We derive two variables: a time-aligned GCS total (which may need to be reconstructed from components) and a binary indicator for fixed pupils. This helps address common EHR issues where component scores are updated without the total.

GCS In general the GCS total is the sum of the three components, eye, verbal, and motor. In the development of the Phoenix criteria we found that there were cases where one or two of the components would be updated but the other component(s) and the total were not updated. When this occurred we chose to assume it was because there was no change in the other component scores. This rule prevents missing totals from artificially lowering the score when only one component is updated.

$$GCS(t; m_{gcs}) := \begin{cases} \mathcal{C}[GCS_{total}](t; m_{gcs}) & \Delta_{GCS_{total}}(t) < \min_{i \in \mathbf{G}} \Delta_{GCS_i}(t) \\ \sum_{i \in \mathbf{G}} \mathcal{C}[GCS_i](t; \infty) & \left[\Delta_{GCS_{total}}(t) = \text{NA} \vee \min_{i \in \mathbf{G}} \Delta_{GCS_i}(t) \leq \Delta_{GCS_{total}}(t) \right] \wedge \\ & \min_{i \in \mathbf{G}} \Delta_{GCS_i}(t) \leq m_{gcs} \\ \text{NA} & \text{otherwise,} \end{cases} \quad (32)$$

where

$$\mathbf{G} := \{\text{motor, eye, verbal}\}. \quad (33)$$

The use of $\mathcal{C}[\cdot](t; \infty)$ means “carry forward the most recent value with no time limit.” In practice, this is only applied when each of the three component scores has been observed at least once. So if the latest eye, verbal, and motor scores were recorded at different times, we take each component’s most recent value and sum them to form a consistent total, even if those component timestamps are not identical.

The score contribution from GCS is

$$\mathcal{D}[GCS](t; m_{gcs}) := \mathbb{I}\{GCS(t; m_{gcs}) \leq 10\} \quad (34)$$

Pupils This is treated as a severe neurologic finding and can dominate the neurologic score. The bilaterally fixed pupils are defined as:

$$\mathcal{D}[Pupils](t; m_{pupil}) := 2 \times \mathbb{I}\{\mathcal{C}[Pupils](t; m_{pupil}) = 1\}. \quad (35)$$

3.3.3 Neurological Organ Dysfunction Score

The neurological component on the score is:

$$\mathcal{D}[Neuro](t; m_{gcs}, m_{pupil}) := \max\{\mathcal{D}[GCS](t; m_{gcs}), \mathcal{D}[Pupils](t; m_{pupil})\}. \quad (36)$$

3.4 Coagulation

The coagulation score uses common laboratory markers of clotting and fibrinolysis. Each marker contributes at most one point, and the total is capped at 2. This keeps the score sensitive to abnormalities without overweighting multiple abnormal labs that can occur together in the same patient.

For readers without a clinical background: Clotting (coagulation) is the body’s process for stopping bleeding by forming fibrin clots. Fibrinolysis is the counter-process that breaks down those clots once they are no longer needed. INR is a standardized measure of how long it takes blood to clot; higher INR means slower clotting and can reflect coagulopathy, liver dysfunction, or anticoagulation. D-dimer is a breakdown product of cross-linked fibrin, so elevated levels suggest recent or ongoing clot formation and breakdown, though it is not specific to any single disease.

In severe infection, clotting can be activated even without trauma. Inflammatory signals and endothelial injury can trigger coagulation pathways, leading to micro-clot formation in small vessels. At the same time, fibrinolysis can be overwhelmed or dysregulated. In extreme cases this becomes disseminated intravascular coagulation (DIC), where clotting and bleeding risk coexist.

3.4.1 Data from the EHR

Table 6: EHR Inputs for Coagulation Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
Platelets	Platelets	Real	1000 / μL	$[0, \infty)$
INR	INR	Real	unitless	$[0, \infty)$
DDimer	D-Dimer	Real	mg/L (FEU)	$[0, \infty)$
Fibrinogen	Fibrinogen	Real	mg/dL	$[0, \infty)$

3.4.2 Constructed Variable

There are no variables to construct.

3.4.3 Coagulation Organ Dysfunction Score

Each indicator below tests whether a lab value crosses a clinically relevant threshold. The score sums how many thresholds are met, then caps at 2 points.

Define:

$$\begin{aligned}
\mathcal{D}[\text{Platelets}](t; m_{\text{coag}}) &:= \mathbb{I}\{\mathcal{C}[\text{Platelets}](t; m_{\text{coag}}) < 100\} \\
\mathcal{D}[\text{INR}](t; m_{\text{coag}}) &:= \mathbb{I}\{\mathcal{C}[\text{INR}](t; m_{\text{coag}}) > 1.3\} \\
\mathcal{D}[\text{DDimer}](t; m_{\text{coag}}) &:= \mathbb{I}\{\mathcal{C}[\text{DDimer}](t; m_{\text{coag}}) > 2\} \\
\mathcal{D}[\text{Fibrinogen}](t; m_{\text{coag}}) &:= \mathbb{I}\{\mathcal{C}[\text{Fibrinogen}](t; m_{\text{coag}}) < 100\},
\end{aligned} \tag{37}$$

and then

$$\mathcal{D}[\text{Coag}](t; m_{\text{coag}}) := \min \left\{ 2, \sum_{x \in \mathcal{C}_{\text{coag}}} \mathcal{D}[x](t; m_{\text{coag}}) \right\}, \tag{38}$$

where

$$\mathcal{C}_{\text{coag}} := \{\text{Platelets}, \text{INR}, \text{DDimer}, \text{Fibrinogen}\}. \tag{39}$$

3.5 Endocrine

The endocrine score is part of the Phoenix-8 scoring system.

3.5.1 Data from the EHR

There is only one element from the EHR used in the endocrine dysfunction assessment: glucose, Table 7.

Table 7: EHR Inputs for Endocrine Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
Glucose	Blood Glucose	Real	mg/dL	$[0, \infty)$

3.5.2 Constructed Variables

There are no constructed variables to define.

3.5.3 Endocrine Organ Dysfunction Score

$$\mathcal{D}[\text{Endo}](t; m_{\text{endo}}) := \mathbb{I}\{\mathcal{C}[\text{Glucose}](t; m_{\text{endo}}) < 50 \vee \mathcal{C}[\text{Glucose}](t; m_{\text{endo}}) > 150\} \tag{40}$$

The domain for the endocrine score is

$$\mathcal{D}[\text{Endo}](t; m_{\text{endo}}) \in \{0, 1\}, \quad \forall m_{\text{endo}}, t. \tag{41}$$

3.6 Immunologic

The immunologic score is part of the Phoenix-8 scoring system.

3.6.1 Data from the EHR

Two values are required from the EHR: Absolute neutrophil count (ANC) and absolute lymphocyte count (ALC). Both are expected in units of 1000 cells per cubic millimeter, Table 8.

Table 8: EHR Inputs for Immunologic Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
ANC	Absolute neutrophil count	Real	1000 cells / mm ³	[0, ∞)
ALC	Absolute lymphocyte count	Real	1000 cells / mm ³	[0, ∞)

3.6.2 Constructed Variables

There are no constructed variables for the immunologic score.

3.6.3 Immunologic Organ Dysfunction Score

Define

$$\mathcal{D}[\text{ALC}](t; m_{\text{immu}}) := \mathbb{I}\{\mathcal{C}[\text{ALC}](t; m_{\text{immu}}) < 1.00\}, \quad (42)$$

and

$$\mathcal{D}[\text{ANC}](t; m_{\text{immu}}) := \mathbb{I}\{\mathcal{C}[\text{ANC}](t; m_{\text{immu}}) < 0.50\}. \quad (43)$$

Then

$$\mathcal{D}[\text{Immu}](t; m_{\text{immu}}) := \max\{\mathcal{D}[\text{ALC}](t; m_{\text{immu}}), \mathcal{D}[\text{ANC}](t; m_{\text{immu}})\}. \quad (44)$$

The domain for the immunologic score is

$$\mathcal{D}[\text{Immu}](t; m_{\text{immu}}) \in \{0, 1\}, \quad \forall m_{\text{immu}}, t. \quad (45)$$

3.7 Hepatic

3.7.1 Data from the EHR

Two values are required from the EHR: total bilirubin reported in units of mg/dL and alanine aminotransferase (ALT) reported in units of IU/L, Table 9.

Table 9: EHR Inputs for Hepatic Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
Bilirubin	Total bilirubin	Real	mg/dL	[0, ∞)
ALT	Alanine aminotransferase	Real	IU / L	[0, ∞)

3.7.2 Constructed Variables

There are no constructed variables for the hepatic organ dysfunction assessment.

3.7.3 Hepatic Organ Dysfunction Score

As with endocrine and immunologic scores, the hepatic score can be a 0 or 1 from either the bilirubin or ALT values.

Define:

$$\mathcal{D}[\text{Bilirubin}](t; m_{\text{hepatic}}) := \mathbb{I}\{\mathcal{C}[\text{Bilirubin}](t; m_{\text{hepatic}}) \geq 4\}, \quad (46)$$

and

$$\mathcal{D}[\text{ALT}](t; m_{\text{hepatic}}) := \mathbb{I}\{\mathcal{C}[\text{ALT}](t; m_{\text{hepatic}}) > 102\}. \quad (47)$$

Then

$$\mathcal{D}[\text{Hepatic}](t; m_{\text{hepatic}}) := \max\{\mathcal{D}[\text{Bilirubin}](t; m_{\text{hepatic}}), \mathcal{D}[\text{ALT}](t; m_{\text{hepatic}})\}. \quad (48)$$

The domain for the hepatic score is

$$\mathcal{D}[\text{Hepatic}](t; m_{\text{hepatic}}) \in \{0, 1\}, \quad \forall m_{\text{hepatic}}, t. \quad (49)$$

3.8 Renal

The renal score is part of the Phoenix-8 scoring system.

3.8.1 Data from the EHR

The patient's age (at admission, in months) and their creatinine levels (in mg/dL) are required to assess renal dysfunction. Age is also required for assessment of cardiovascular dysfunction as part of the Phoenix scoring and thus the only additional variable to collect to extend to Phoenix-8 from Phoenix for the renal dysfunction score is the creatinine.

Table 10: EHR Inputs for Renal Organ Dysfunction Scoring

Variable	Description	Type	Units	Domain
Creatinine	Creatinine	Real	mg / dL	$[0, \infty)$
Age	Patient's age	Real	months	$[0, 216)$

3.8.2 Constructed Variables

There are no constructed variables required for the assessment of renal dysfunction.

3.8.3 Renal Organ Dysfunction Score

The renal dysfunction score is based on an age adjusted creatinine level (50). The cut points for the age ranges are the same as used in the cardiovascular scoring, (24) and (25).

$$\mathcal{D}[\text{Renal}](t; m_{\text{renal}}, \text{Age}) := \begin{cases} 1 & \text{Age} \in [0, 1) \wedge \mathcal{C}[\text{Creatinine}](t; m_{\text{renal}}) \geq 0.8 \\ 1 & \text{Age} \in [1, 12) \wedge \mathcal{C}[\text{Creatinine}](t; m_{\text{renal}}) \geq 0.3 \\ 1 & \text{Age} \in [12, 24) \wedge \mathcal{C}[\text{Creatinine}](t; m_{\text{renal}}) \geq 0.4 \\ 1 & \text{Age} \in [24, 60) \wedge \mathcal{C}[\text{Creatinine}](t; m_{\text{renal}}) \geq 0.6 \\ 1 & \text{Age} \in [60, 144) \wedge \mathcal{C}[\text{Creatinine}](t; m_{\text{renal}}) \geq 0.7 \\ 1 & \text{Age} \in [144, 216) \wedge \mathcal{C}[\text{Creatinine}](t; m_{\text{renal}}) \geq 1.0 \\ 0 & \text{otherwise.} \end{cases} \quad (50)$$

3.9 Suspected Infection

A patient is defined to have a suspected infection if they have at least one infectious test *ordered* and at least one dose of a systemic anti-infective agent. Results of the tests are not needed, just that the tests have been ordered. Anti-infective agents include anti-bacterials, anti-fungals, anti-virals, or anti-mycobacterial (anti-buberculosis). This definition is operational: it identifies when clinicians treated the case as infectious, not whether infection was ultimately confirmed.

Let t_1 denote the time when an infectious test was ordered. Let t_2 denote the time when the first dose of a systemic anti-infectious agent was administered.

Suspected infection status is defined as

$$\text{SI}(t) := \mathbb{I}\{t_1 \leq t\} \times \mathbb{I}\{t_2 \leq t\}. \quad (51)$$

4 Phoenix Sepsis Score

Define the look-back windows as the vector

$$\mathbf{m} := (m_{\text{resp}}, m_{\text{vaso}}, m_{\text{bp}}, m_{\text{lac}}, m_{\text{gcs}}, m_{\text{pupil}}, m_{\text{coag}}, m_{\text{endo}}, m_{\text{immu}}, m_{\text{hepatic}}, m_{\text{renal}}). \quad (52)$$

For notational compactness, $\mathcal{D}[x](t; \mathbf{m}, \text{Age})$ uses a common argument list for all scoring entities x . Each scoring function depends only on the elements of $\mathbf{m}$, and on Age, that are relevant to its definition; all other arguments are ignored.

In this definition, the time interval of interest is the same for both SI and Phoenix Sepsis Score (PSS), but the maxima are taken separately. That is, a patient can have $\text{SI}(t) = 1$ at one time and the maximum organ dysfunction score at another. The Phoenix Sepsis Score combines organ dysfunction with evidence of suspected infection (via SI), so it reflects organ dysfunction *in the presence of* suspected infection.

$$\text{PSS}_d(T; \mathbf{m}, \text{Age}) := \left[\max_{t \in T} \text{SI}(t) \right] \left[\max_{t \in T} \sum_{x \in \Omega_d} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) \right], \quad d \in \{4, 8\}, \quad (53)$$

where

$$\Omega_4 := \{\text{Resp}, \text{Card}, \text{Neuro}, \text{Coag}\} \quad (54)$$

and

$$\Omega_8 := \Omega_4 \cup \{\text{Endo}, \text{Immu}, \text{Hepatic}, \text{Renal}\}. \quad (55)$$

Sepsis is defined as organ-dysfunction in the presence of a suspected infection. Specifically, a score of at least $\sigma > 0$ points:

$$\text{Sepsis}(T; \mathbf{m}, \text{Age}) := \mathbb{I}\{\text{PSS}_4(T; \mathbf{m}, \text{Age}) \geq \sigma\}. \quad (56)$$

Septic Shock is sepsis with at least κ point(s) from cardiovascular dysfunction.

$$\text{SepticShock}(T; \mathbf{m}, \text{Age}) := \mathbb{I}\left\{ \max_{t \in T} \text{SI}(t) \times \max_{t \in T} \left(\sum_{x \in \Omega_4} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) \mathbb{I}\{\mathcal{D}[\text{Card}](t; \mathbf{m}, \text{Age}) \geq \kappa\} \right) \geq \sigma \right\}, \quad (57)$$

where $\sigma \geq \kappa > 0$. Note that the specific shock definition does not simply consider any cardiovascular dysfunction in the window T . It must occur with the organ dysfunction associated with reaching the sepsis score threshold.

For the published Phoenix Sepsis Criteria [Sanchez-Pinto et al., 2024, Schlappbach et al., 2024] the following parameter values in Table 11 were used:

Table 11: Parameter values defining time windows, look-backs, and thresholds for the Phoenix Sepsis Score, Sepsis, and Septic Shock.

Parameter	Value	Notes
Time Window		
T	$[0, 1440)$	The first 24-hours of the encounter
Thresholds		
σ	2	
κ	1	
Look-backs		
m_{resp}	360 minutes	all respiratory EHR variables
m_{vaso}	720 minutes	vasoactive medications
m_{bp}	360 minutes	blood pressure / MAP
m_{lac}	360 minutes	lactate
m_{gcs}	720 minutes	GCS components or total
m_{pupil}	720 minutes	pupils
m_{coag}	1440 minutes	coagulation labs
m_{endo}	720 minutes	glucose
m_{immu}	1440 minutes	ANC and ALC
m_{hepatic}	1440 minutes	ALT and Total bilirubin
m_{renal}	1440 minutes	creatinine

Importantly, restricting the maximization to $t \in T$ does **not** truncate the underlying data. For any $t \in T$, the carry-forward operator $\mathcal{C}[\cdot](t; m)$ can still use measurements recorded before T (even at negative times) as long as they fall within the look-back window m . Conceptually, this JAMA definition is time-aligned: at each time t we sum the organ dysfunction components evaluated at that same time, then take the maximum of that sum over T . This preserves temporal alignment across organ systems, so a high score reflects concurrent multi-organ dysfunction at a single time point.

5 Exploratory Aggregation Schemes

There are several exploratory aggregation schemes. Daily and block-based “worst value” aggregation is common in pediatric organ dysfunction scoring (e.g., PELOD/qPELOD and pSOFA), which motivates the time-decoupled exploratory aggregation schemes below. See, for example, daily PELOD scoring using the most abnormal value each day and pSOFA evaluations that use maximum/mean daily scores or the highest score in fixed time blocks. [Leteurtre et al., 2003, Bestati et al., 2010, Matics and Sanchez-Pinto, 2017, Badke et al., 2025]

5.1 Organ-Level Maxima

The primary definition for PSS in (53) looks for the max of the sum of the organ dysfunction scores. This exploratory scheme looks at the sum of the max of each of the organ system dysfunction scores.

$$\text{PSS}_d^{\text{OLM}}(T; \mathbf{m}, \text{Age}) := \left[\max_{t \in T} \text{SI}(t) \right] \left[\sum_{x \in \Omega_d} \max_{t \in T} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) \right], \quad d \in \{4, 8\}. \quad (58)$$

$\text{PSS}_d^{\text{OLM}}(T; \mathbf{m}, \text{Age})$ decouples time across organ systems. It takes the maximum score for each organ system independently and then sums those maxima. This measures the overall burden of dysfunction over the interval, not necessarily dysfunction occurring at the same time. As a result, $\text{PSS}_d^{\text{OLM}}(T; \mathbf{m}, \text{Age})$ can

be larger than the $PSS_d(T; \mathbf{m}, \text{Age})$, because peaks from different time points can accumulate. That is, by construction,

$$PSS_d(T; \mathbf{m}, \text{Age}) \leq PSS_d^{\text{OLM}}(T; \mathbf{m}, \text{Age}). \quad (59)$$

This follows from the max-sum inequality:

$$\max_t \sum_x f_x(t) \leq \sum_x \max_t f_x(t). \quad (60)$$

Using (58) to define sepsis is not structurally different from the primary definition (56).

$$\text{Sepsis}^{\text{OLM}}(T; \mathbf{m}, \text{Age}) := \mathbb{I} \{ PSS_4^{\text{OLM}}(T; \mathbf{m}, \text{Age}) \geq \sigma \}. \quad (61)$$

Septic Shock, for organ-level maxima, is an easier calculation than the primary definition (57).

$$\text{SepticShock}^{\text{OLM}}(T; \mathbf{m}, \text{Age}) := [\text{Sepsis}^{\text{OLM}}(T; \mathbf{m}, \text{Age})] \mathbb{I} \left\{ \max_{t \in T} \mathcal{D}[\text{Card}](t; \mathbf{m}, \text{Age}) \geq \kappa \right\}. \quad (62)$$

for $\sigma \geq \kappa > 0$.

5.2 Organ-Level + Cardiovascular Component Decoupling

In this exploratory aggregation scheme we decouple all organ systems **and** the three cardiovascular components: vasoactive medications, MAP, and lactate.

$$PSS_d^{\text{CCD}}(T; \mathbf{m}, \text{Age}) := \left[\max_{t \in T} \text{SI}(t) \right] \left[\sum_{x \in \Omega_d \setminus \{\text{Card}\}} \max_{t \in T} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) + \sum_{x \in \mathbf{C}_{\text{card}}} \max_{t \in T} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) \right], \quad d \in \{4, 8\}, \quad (63)$$

where

$$\mathbf{C}_{\text{card}} := \{\text{Vasos}, \text{MAP}, \text{Lactate}\}. \quad (64)$$

The construction here creates the relationship:

$$PSS_d(T; \mathbf{m}, \text{Age}) \leq PSS_d^{\text{OLM}}(T; \mathbf{m}, \text{Age}) \leq PSS_d^{\text{CCD}}(T; \mathbf{m}, \text{Age}). \quad (65)$$

Sepsis and Septic Shock are then defined as:

$$\text{Sepsis}^{\text{CCD}}(T; \mathbf{m}, \text{Age}) := \mathbb{I} \{ PSS_4^{\text{CCD}}(T; \mathbf{m}, \text{Age}) \geq \sigma \}, \quad (66)$$

and

$$\text{SepticShock}^{\text{CCD}}(T; \mathbf{m}, \text{Age}) := [\text{Sepsis}^{\text{CCD}}(T; \mathbf{m}, \text{Age})] \mathbb{I} \left\{ \sum_{x \in \mathbf{C}_{\text{card}}} \max_{t \in T} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) \geq \kappa \right\} \quad (67)$$

for $\sigma \geq \kappa > 0$.

5.3 Full Component Decoupling

Under full component decoupling, each point-producing component is aggregated independently over T and then combined using the original organ-specific capping or maximum rules. Component scores are maximized over T ; for respiratory oxygenation, lower PFR and SFR values represent greater dysfunction and are therefore minimized.

$$\begin{aligned} \text{Resp}^{\text{FCD}}(T; m_{\text{resp}}) &:= \max_{t \in T} (\text{IMV}(t; m_{\text{resp}})) \times (\mathbb{I}\{B_3(T; m_{\text{resp}})\} + \mathbb{I}\{B_2(T; m_{\text{resp}})\}) \\ &\quad + \max_{t \in T} (\text{ORS}(t; m_{\text{resp}})) \times \mathbb{I}\{B_1(T; m_{\text{resp}})\}, \end{aligned} \quad (68)$$

where

$$\begin{aligned} B_1(T; m_{\text{resp}}) &:= \min_{t \in T} \text{PFR}(t; m_{\text{resp}}) < 400 \vee \min_{t \in T} \text{SFR}(t; m_{\text{resp}}) < 292, \\ B_2(T; m_{\text{resp}}) &:= \min_{t \in T} \text{PFR}(t; m_{\text{resp}}) < 200 \vee \min_{t \in T} \text{SFR}(t; m_{\text{resp}}) < 220, \\ B_3(T; m_{\text{resp}}) &:= \min_{t \in T} \text{PFR}(t; m_{\text{resp}}) < 100 \vee \min_{t \in T} \text{SFR}(t; m_{\text{resp}}) < 148. \end{aligned} \quad (69)$$

This scheme decouples respiratory support from oxygenation severity; the required support and oxygenation threshold need not occur at the same time within T .

The overall scores are then defined as:

$$\begin{aligned} \text{PSS}_d^{\text{FCD}}(T; \mathbf{m}, \text{Age}) &:= \\ &\left[\max_{t \in T} \text{SI}(t) \right] \left[\text{Resp}^{\text{FCD}}(T; m_{\text{resp}}) + \min \left\{ 2, \sum_{x \in \mathcal{C}_{\text{coag}}} \max_{t \in T} \mathcal{D}[x](t; \mathbf{m}) \right\} \right. \\ &\quad \left. + \sum_{x \in \omega_d} \max_{t \in T} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) \right], \quad d \in \{4, 8\}, \end{aligned} \quad (70)$$

where

$$\omega_4 := \{\text{Vasos}, \text{MAP}, \text{Lactate}, \text{Neuro}\}, \quad (71)$$

and

$$\omega_8 := \omega_4 \cup \{\text{Endo}, \text{Immu}, \text{Hepatic}, \text{Renal}\}. \quad (72)$$

Note: the contributions from neurologic, immunologic, and hepatic scores do consider more than one input from the EHR. However, because

$$\max_{t \in T} \max \{ \mathcal{D}[\text{GCS}](t; m_{\text{gcs}}), \mathcal{D}[\text{Pupils}](t; m_{\text{pupil}}) \} = \max \left\{ \max_{t \in T} \mathcal{D}[\text{GCS}](t; m_{\text{gcs}}), \max_{t \in T} \mathcal{D}[\text{Pupils}](t; m_{\text{pupil}}) \right\} \quad (73)$$

we need not explicitly decouple these component scores.

The constuction here creates the relationship:

$$\text{PSS}_d(T; \mathbf{m}, \text{Age}) \leq \text{PSS}_d^{\text{OLM}}(T; \mathbf{m}, \text{Age}) \leq \text{PSS}_d^{\text{CCD}}(T; \mathbf{m}, \text{Age}) \leq \text{PSS}_d^{\text{FCD}}(T; \mathbf{m}, \text{Age}). \quad (74)$$

Sepsis is defined as:

$$\text{Sepsis}^{\text{FCD}}(T; \mathbf{m}, \text{Age}) := \mathbb{I} \{ \text{PSS}_4^{\text{FCD}}(T; \mathbf{m}, \text{Age}) \geq \sigma \}, \quad (75)$$

for $\sigma > 0$.

Septic shock is defined as follows:

$$\text{SepticShock}^{\text{FCD}}(T; \mathbf{m}, \text{Age}) := [\text{Sepsis}^{\text{FCD}}(T; \mathbf{m}, \text{Age})] \mathbb{I} \left\{ \sum_{x \in \mathcal{C}_{\text{card}}} \max_{t \in T} \mathcal{D}[x](t; \mathbf{m}, \text{Age}) \geq \kappa \right\} \quad (76)$$

for $\sigma \geq \kappa > 0$.

References

- Colleen M. Badke, Austin Wang, Latasha A. Daniels, and L. Nelson Sanchez-Pinto. Validation of pediatric sequential organ failure assessment (psoba) scores to predict critical events in the pediatric intensive care unit. *Journal of Intensive Care Medicine*, 40(5):565–570, 2025. doi: 10.1177/08850666241307630.
- Melania M Bembea, Michael Agus, Ayse Akcan-Arikan, Peta Alexander, Rajit Basu, Tellen D Bennett, Desmond Bohn, Leonardo R Brandão, Ann-Marie Brown, Joseph A Carcillo, et al. Pediatric organ dysfunction information update mandate (podium) contemporary organ dysfunction criteria: executive summary. *Pediatrics*, 149(Supplement_1):S1–S12, 2022.
- Nadine Bestati, Stéphane Leteurtre, Alain Duhamel, François Proulx, Bruno Grandbastien, Jacques Lacroix, and Francis Leclerc. Differences in organ dysfunctions between neonates and older children: A prospective, observational, multicenter study. *Critical Care*, 14(5):R202, 2010. doi: 10.1186/cc9323.
- Anwarul Haque, NR Siddiqui, O Munir, S Saleem, and Asad Mian. Association between vasoactive-inotropic score and mortality in pediatric septic shock. *Indian pediatrics*, 52(4):311–313, 2015.
- Stéphane Leteurtre, Arnaud Martinot, Alain Duhamel, France Gauvin, Bruno Grandbastien, Tuyet V. Nam, François Proulx, Jacques Lacroix, and Francis Leclerc. Validation of the paediatric logistic organ dysfunction (pelod) score: Prospective, observational, multicentre study. *The Lancet*, 362(9379):192–197, 2003. doi: 10.1016/S0140-6736(03)13908-6.
- Greg S Martin and Gordon R Bernard. Airway and lung in sepsis. *Intensive care medicine*, 27:S63, 2001.
- Travis J. Matics and L. Nelson Sanchez-Pinto. Adaptation and validation of a pediatric sequential organ failure assessment score and evaluation of the sepsis-3 definitions in critically ill children. *JAMA Pediatrics*, 171(10):e172352, 2017. doi: 10.1001/jamapediatrics.2017.2352.
- L. Nelson Sanchez-Pinto, Tellen D. Bennett, Peter E. DeWitt, Seth Russell, Margaret N. Rebull, Blake Martin, Samuel Akech, David J. Albers, Elizabeth R. Alpern, Fran Balamuth, Melania Bembea, Mohammad Jobayer Chisti, Idris Evans, Christopher M. Horvat, Juan Camilo Jaramillo-Bustamante, Niranjana Kissoon, Kusum Menon, Halden F. Scott, Scott L. Weiss, Matthew O. Wiens, Jerry J. Zimmerman, Andrew C. Argent, Lauren R. Sorce, Luregn J. Schlapbach, R. Scott Watson, and Society of Critical Care Medicine Pediatric Sepsis Definition Task Force. Development and Validation of the Phoenix Criteria for Pediatric Sepsis and Septic Shock. *JAMA*, 01 2024. ISSN 0098-7484. doi: 10.1001/jama.2024.0196. URL <https://doi.org/10.1001/jama.2024.0196>. Drs Schlapbach, Watson, Sorce, and Argent contributed equally. Drs Sanchez-Pinto and Bennett contributed equally.
- Luregn J. Schlapbach, R. Scott Watson, Lauren R. Sorce, Andrew C. Argent, Kusum Menon, Mark W. Hall, Samuel Akech, David J. Albers, Elizabeth R. Alpern, Fran Balamuth, Melania Bembea, Paolo Biban, Enitan D. Carrol, Kathleen Chiotos, Mohammad Jobayer Chisti, Peter E. DeWitt, Idris Evans, Cláudio Flauzino de Oliveira, Christopher M. Horvat, David Inwald, Paul Ishimine, Juan Camilo Jaramillo-Bustamante, Michael Levin, Rakesh Lodha, Blake Martin, Simon Nadel, Satoshi Nakagawa, Mark J. Peters, Adrienne G. Randolph, Suchitra Ranjit, Margaret N. Rebull, Seth Russell, Halden F. Scott, Daniela Carla de Souza, Pierre Tissieres, Scott L. Weiss, Matthew O. Wiens, James L. Wynn, Niranjana Kissoon, Jerry J. Zimmerman, L. Nelson Sanchez-Pinto, Tellen D. Bennett, and Society of Critical Care Medicine Pediatric Sepsis Definition Task Force. International consensus criteria for pediatric sepsis and septic shock. *JAMA*, 01 2024. ISSN 0098-7484. doi: 10.1001/jama.2024.0179. URL <https://doi.org/10.1001/jama.2024.0179>. Drs Sanchez-Pinto and Bennett contributed equally. Drs DeWitt and Mr Russell contributed equally. Drs Argent, Sorce, Schlapbach, and Watson contributed equally.

James E. Squires, Patrick J. McKiernan, and on behalf of the Pediatric Organ Dysfunction Information Update Mandate (PODIUM) Collaborative Squires, Robert H. Acute Liver Dysfunction Criteria in Critically Ill Children: The PODIUM Consensus Conference. *Pediatrics*, 149(Supplement_1):S59–S65, 01 2022. ISSN 0031-4005. doi: 10.1542/peds.2021-052888I. URL <https://doi.org/10.1542/peds.2021-052888I>.

DRAFT