The Invertible Matrix Theorem

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit https://github.com/CU-Denver-MathStats-DER

Let A be a square $n \times n$ matrix.

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix I_n .
- (c) A has n pivots when forming the reduced row echelon form of A.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.

Let A be a square $n \times n$ matrix.

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix I_n .
- (c) A has n pivots when forming the reduced row echelon form of A.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- (d) \Rightarrow (b) If $A\mathbf{x} = \mathbf{0}$ has only the trivial solution, that means the augmented matrix $[A\mathbf{0}]$ has no free variables. Thus A has n pivots.
- (c) \Leftrightarrow (b) A has n pivots if and only if A is row equivalent to the $n \times n$ identity matrix.

Let A be a square $n \times n$ matrix.

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix I_n .
- (c) A has n pivots when forming the reduced row echelon form of A.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- (d) \Rightarrow (b) If $A\mathbf{x} = \mathbf{0}$ has only the trivial solution, that means the augmented matrix $[A\mathbf{0}]$ has no free variables. Thus A has n pivots.
- (c) \Leftrightarrow (b) A has n pivots if and only if A is row equivalent to the $n \times n$ identity matrix.
- (a) \Rightarrow (d) If A is invertible, then $A\mathbf{x} = \mathbf{b}$ has a unique solution for any b in \mathbb{R}^n .
- (b) \Rightarrow (a) If A is row equivalent to I_n , then there are elementary matrices E_i such that $E_p E_{p-1} \cdots E_1 A = I_n$. Thus, $A^{-1} = E_p E_{p-1} \cdots E_1$, so A is invertible.

Let A be a square $n \times n$ matrix.

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix I_n .
- (c) A has n pivots when forming the reduced row echelon form of A.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- (d) \Rightarrow (b) If $A\mathbf{x} = \mathbf{0}$ has only the trivial solution, that means the augmented matrix $[A\mathbf{0}]$ has no free variables. Thus A has n pivots.
- (c) \Leftrightarrow (b) A has n pivots if and only if A is row equivalent to the $n \times n$ identity matrix.
- (a) \Rightarrow (d) If A is invertible, then $A\mathbf{x} = \mathbf{b}$ has a unique solution for any b in \mathbb{R}^n .
- (b) \Rightarrow (a) If A is row equivalent to I_n , then there are elementary matrices E_i such that $E_p E_{p-1} \cdots E_1 A = I_n$. Thus, $A^{-1} = E_p E_{p-1} \cdots E_1$, so A is invertible.

Statements (a), (b), (c), (d) are all equivalent!

Equivalence of Statements

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix, I_n .
- (c) A had n pivots.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- (j) There is an $n \times n$ matrix C such that $CA = I_n$.

Equivalence of Statements

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix, I_n .
- (c) A had n pivots.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- (j) There is an $n \times n$ matrix C such that $CA = I_n$.
- (a) \Rightarrow (j) If A is invertible, then $C = A^{-1}$ is a matrix such that $CA = I_n$.
- (j) \Rightarrow (d) If there is an $n \times n$ matrix C such that $CA = I_n$, then $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.

Equivalence of Statements

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix, I_n .
- (c) A had n pivots.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- (j) There is an $n \times n$ matrix C such that $CA = I_n$.
- (a) \Rightarrow (j) If A is invertible, then $C = A^{-1}$ is a matrix such that $CA = I_n$.
- (j) \Rightarrow (d) If there is an $n \times n$ matrix C such that $CA = I_n$, then $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.

Statements (a), (b), (c), (d), and (j) are all equivalent!

Adding to the Chain of Equivalence

- (a) A is an invertible matrix.
- (g) The equation $A\mathbf{x} = \mathbf{b}$ has at least one solution for each \mathbf{b} in \mathbb{R}^n .
- (k) There is an $n \times n$ matrix D such that $AD = I_n$.

Adding to the Chain of Equivalence

- (a) A is an invertible matrix.
- (g) The equation $A\mathbf{x} = \mathbf{b}$ has at least one solution for each \mathbf{b} in \mathbb{R}^n .
- (k) There is an $n \times n$ matrix D such that $AD = I_n$.
- (a) \Rightarrow (k) If A is invertible, then there is an $n \times n$ matrix $D = A^{-1}$ such that $AD = I_n$.
- (k) \Rightarrow (g) If there is an $n \times n$ matrix D such that $AD = I_n$, then the equation $AD\mathbf{b} = I_n\mathbf{b} = \mathbf{b}$. Thus, for any \mathbf{b} in \mathbb{R}^n , the equation $A\mathbf{x} = \mathbf{b}$ has the solution $\mathbf{x} = D\mathbf{b}$.
- (g) \Rightarrow (a) If the equation $A\mathbf{x} = \mathbf{b}$ has at least one solution for each \mathbf{b} in \mathbb{R}^n , each row in A must have a pivot. Since A is $n \times n$, A has n pivots, and A therefore is invertible.

Adding to the Chain of Equivalence

- (a) A is an invertible matrix.
- (g) The equation $A\mathbf{x} = \mathbf{b}$ has at least one solution for each \mathbf{b} in \mathbb{R}^n .
- (k) There is an $n \times n$ matrix D such that $AD = I_n$.
- (a) \Rightarrow (k) If A is invertible, then there is an $n \times n$ matrix $D = A^{-1}$ such that $AD = I_n$.
- (k) \Rightarrow (g) If there is an $n \times n$ matrix D such that $AD = I_n$, then the equation $AD\mathbf{b} = I_n\mathbf{b} = \mathbf{b}$. Thus, for any \mathbf{b} in \mathbb{R}^n , the equation $A\mathbf{x} = \mathbf{b}$ has the solution $\mathbf{x} = D\mathbf{b}$.
- (g) \Rightarrow (a) If the equation $A\mathbf{x} = \mathbf{b}$ has at least one solution for each \mathbf{b} in \mathbb{R}^n , each row in A must have a pivot. Since A is $n \times n$, A has n pivots, and A therefore is invertible.

Statements (a), (b), (c), (d), (g), (j) and (k) are all equivalent statements!

The Invertible Matrix Theorem (so far)

Let A be a square $n \times n$ matrix. Then all of the following statements are equivalent.

- (a) A is an invertible matrix.
- (b) A is row equivalent to the $n \times n$ identity matrix I_n .
- (c) A has n pivots.
- (d) The equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- (g) The equation $A\mathbf{x} = \mathbf{b}$ has at least one solution for each \mathbf{b} in \mathbb{R}^n .
- (h) The columns of A span \mathbb{R}^n .
- (j) There is an $n \times n$ matrix C such that $CA = I_n$.
- (k) There is an $n \times n$ matrix D such that $AD = I_n$.
- (I) A^T is an invertible matrix.

Example

Consider the matrix

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 2 & 3 & 4 \\ 2 & 3 & 4 \end{bmatrix}.$$

Does A have an inverse? Explain why or why not.

Example

Consider the matrix

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 2 & 3 & 4 \\ 2 & 3 & 4 \end{bmatrix}.$$

Does A have an inverse? Explain why or why not.

- ightharpoonup A is not row equivalent to I_3 .
- ▶ The columns of A do not span \mathbb{R}^3 .
- \triangleright A^T is not invertible.
- ► And so on . . .