

The Characteristic Equation

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Find the eigenvalues of $A = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix}$.

How do we find eigenvalues of matrix A ? For which values of λ :

Find the eigenvalues of $A = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix}$.

How do we find eigenvalues of matrix A ? For which values of λ :

- ▶ Does $A\mathbf{x} = \lambda\mathbf{x}$ have a **nontrivial** solution?
- ▶ Does $(A - \lambda I)\mathbf{x} = \mathbf{0}$ have a nontrivial solution?

Find the eigenvalues of $A = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix}$.

How do we find eigenvalues of matrix A ? For which values of λ :

- ▶ Does $A\mathbf{x} = \lambda\mathbf{x}$ have a **nontrivial** solution?
- ▶ Does $(A - \lambda I)\mathbf{x} = \mathbf{0}$ have a nontrivial solution?
- ▶ Is the matrix $(A - \lambda I)$ **NOT invertible**?

Introduction

Find the eigenvalues of $A = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix}$.

$$\begin{aligned}\det(A - \lambda I) &= \det\left(\begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix}\right) \\ &= \det\begin{bmatrix} 0 - \lambda & 1 \\ -6 & 5 - \lambda \end{bmatrix} \\ &= (0 - \lambda)(5 - \lambda) - (1)(-6) \\ &= \lambda^2 - 5\lambda + 6\end{aligned}$$

How do we find eigenvalues of matrix A ? For which values of λ :

- ▶ Does $A\mathbf{x} = \lambda\mathbf{x}$ have a **nontrivial** solution?
- ▶ Does $(A - \lambda I)\mathbf{x} = \mathbf{0}$ have a nontrivial solution?
- ▶ Is the matrix $(A - \lambda I)$ **NOT invertible**?
- ▶ Is $\det(A - \lambda I) = 0$?

The Characteristic Equation

Find the eigenvalues of $A = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix}$.

The Characteristic Equation

Find the eigenvalues of $A = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix}$.

To find all eigenvalues, we solve the following equation:

$$\det(A - \lambda I) = \lambda^2 - 5\lambda + 6 = 0.$$

- ▶ The scalar equation $\det(A - \lambda I) = 0$ is called the **characteristic equation** of A .
- ▶ The polynomial $\det(A - \lambda I)$ is called the **characteristic polynomial** of A .
- ▶ A scalar λ is an **eigenvalue** of A if and only if λ satisfies $\det(A - \lambda I) = 0$.

Example

Find and solve the characteristic equation

$$\text{of } A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -5 & 0 \\ 1 & 8 & 1 \end{bmatrix}.$$

Example

Find and solve the characteristic equation

$$\text{of } A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -5 & 0 \\ 1 & 8 & 1 \end{bmatrix}.$$

$$\begin{aligned} \det(A - \lambda I) &= \det \begin{bmatrix} 1 - \lambda & 2 & 1 \\ 0 & -5 - \lambda & 0 \\ 1 & 8 & 1 - \lambda \end{bmatrix} \\ &= (-5 - \lambda) \det \begin{bmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{bmatrix} \\ &= -(5 + \lambda) [(1 - \lambda)^2 - 1] \\ &= -(5 + \lambda)(\lambda^2 - 2\lambda) \\ &= -\lambda(5 + \lambda)(\lambda - 2) \end{aligned}$$

The **characteristic equation** $-\lambda(5 + \lambda)(\lambda - 2) = 0$ has solutions $\lambda = 0$, -5 , and 2 .

So the eigenvalues of A are $\lambda = 0$, -5 , and 2 .

Invertible Matrix Theorem Continued, Even More!

When is 0 an eigenvalue of an $n \times n$ matrix A ?

Invertible Matrix Theorem Continued, Even More!

When is 0 an eigenvalue of an $n \times n$ matrix A ?

- ▶ **Zero is an eigenvalue** of A if and only if there is a nonzero vector \mathbf{x} such that $A\mathbf{x} = 0\mathbf{x} = \mathbf{0}$, which happens if and only if $0 = \det(A - 0I) = \det A$.

Invertible Matrix Theorem Continued, Even More!

When is 0 an eigenvalue of an $n \times n$ matrix A ?

- ▶ **Zero is an eigenvalue** of A if and only if there is a nonzero vector \mathbf{x} such that $A\mathbf{x} = 0\mathbf{x} = \mathbf{0}$, which happens if and only if $0 = \det(A - 0I) = \det A$.
- ▶ $A\mathbf{x} = \mathbf{0}$ has a nontrivial solution if and only if A is **NOT invertible**.
- ▶ Hence A is invertible if and only if 0 is NOT an eigenvalue.

Invertible Matrix Theorem Continued, Even More!

When is 0 an eigenvalue of an $n \times n$ matrix A ?

- ▶ **Zero is an eigenvalue** of A if and only if there is a nonzero vector \mathbf{x} such that $A\mathbf{x} = 0\mathbf{x} = \mathbf{0}$, which happens if and only if $0 = \det(A - 0I) = \det A$.
- ▶ $A\mathbf{x} = \mathbf{0}$ has a nontrivial solution if and only if A is **NOT invertible**.
- ▶ Hence A is invertible if and only if 0 is NOT an eigenvalue.

Theorem (The Invertible Matrix Theorem (continued))

Let A be an $n \times n$ matrix. Then the following are equivalent statements:

- (a) A is an invertible matrix.
- (r) The number **0 is not an eigenvalue** of A .

Eigenvalues of Triangular Matrices

Find the eigenvalues of

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 2 & 3 & 0 & 0 & 0 & 0 \\ 3 & 4 & 5 & 0 & 0 & 0 \\ 6 & 7 & 8 & 9 & 0 & 0 \\ -1 & -2 & -3 & -4 & -5 & 0 \\ -6 & -7 & -8 & -9 & -10 & 1 \end{bmatrix}.$$

- ▶ If A is a **triangular matrix**, then the **entries on the main diagonal** are the eigenvalues of A .
- ▶ The **algebraic multiplicity of an eigenvalue** is the multiplicity of the corresponding root of the characteristic equation of A .

Definition

If A and B are two $n \times n$ matrices, we say A is similar to B if there is an invertible matrix P such that $P^{-1}AP = B$, or equivalently $A = PBP^{-1}$.

- ▶ If A is similar to B , then B is similar to A .

Similar Matrices

Definition

If A and B are two $n \times n$ matrices, we say A is similar to B if there is an invertible matrix P such that $P^{-1}AP = B$, or equivalently $A = PBP^{-1}$.

► If A is similar to B , then B is similar to A .

$$A = \begin{bmatrix} -13 & -8 & -4 \\ 12 & 7 & 4 \\ 24 & 16 & 7 \end{bmatrix} \text{ is similar to } B = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{bmatrix} \text{ since}$$

$$P^{-1}AP = \begin{bmatrix} -6 & -4 & -1 \\ -3 & -2 & -1 \\ 5 & 3 & 1 \end{bmatrix} \begin{bmatrix} -13 & -8 & -4 \\ 12 & 7 & 4 \\ 24 & 16 & 7 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ -2 & -1 & -3 \\ 1 & -2 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{bmatrix} = B.$$

Eigenvalues of Similar Matrices

Theorem

If $n \times n$ matrices A and B are **similar**, then they have the **same characteristic polynomial** and hence the **same eigenvalues** (with the same algebraic multiplicities).

Eigenvalues of Similar Matrices

Theorem

If $n \times n$ matrices A and B are **similar**, then they have the **same characteristic polynomial** and hence the **same eigenvalues** (with the same algebraic multiplicities).

Proof.

Let A and B be similar matrices.

So $B = P^{-1}AP$ for some inv matrix P .

Eigenvalues of Similar Matrices

Theorem

If $n \times n$ matrices A and B are **similar**, then they have the **same characteristic polynomial** and hence the **same eigenvalues** (with the same algebraic multiplicities).

Proof.

Let A and B be similar matrices.

So $B = P^{-1}AP$ for some inv matrix P .

$$\begin{aligned} B - \lambda I &= P^{-1}AP - \lambda P^{-1}P \\ &= P^{-1}AP - P^{-1}(\lambda I)P \\ &= P^{-1}(A - \lambda I)P. \end{aligned}$$

Eigenvalues of Similar Matrices

Theorem

If $n \times n$ matrices A and B are **similar**, then they have the **same characteristic polynomial** and hence the **same eigenvalues** (with the same algebraic multiplicities).

Proof.

Let A and B be similar matrices.

So $B = P^{-1}AP$ for some inv matrix P .

$$\begin{aligned} B - \lambda I &= P^{-1}AP - \lambda P^{-1}P \\ &= P^{-1}AP - P^{-1}(\lambda I)P \\ &= P^{-1}(A - \lambda I)P. \end{aligned}$$

This gives

$$\begin{aligned} \det(B - \lambda I) &= \det(P^{-1}(A - \lambda I)P) \\ &= (\det P^{-1})(\det(A - \lambda I))(\det P) \\ &= (\det P^{-1})(\det P)(\det(A - \lambda I)) \\ &= \det(A - \lambda I). \end{aligned}$$

Eigenvalues of Similar Matrices

Theorem

If $n \times n$ matrices A and B are **similar**, then they have the **same characteristic polynomial** and hence the **same eigenvalues** (with the same algebraic multiplicities).

Proof.

Let A and B be similar matrices.

So $B = P^{-1}AP$ for some inv matrix P .

$$\begin{aligned} B - \lambda I &= P^{-1}AP - \lambda P^{-1}P \\ &= P^{-1}AP - P^{-1}(\lambda I)P \\ &= P^{-1}(A - \lambda I)P. \end{aligned}$$

This gives

$$\begin{aligned} \det(B - \lambda I) &= \det(P^{-1}(A - \lambda I)P) \\ &= (\det P^{-1})(\det(A - \lambda I))(\det P) \\ &= (\det P^{-1})(\det P)(\det(A - \lambda I)) \\ &= \det(A - \lambda I). \end{aligned}$$

Since $\det(B - \lambda I) = \det(A - \lambda I)$, the two similar matrices have the same characteristic polynomial, and thus the same eigenvalues. □

Example

Find the eigenvalues of $A = \begin{bmatrix} -13 & -8 & -4 \\ 12 & 7 & 4 \\ 24 & 16 & 7 \end{bmatrix}$.

Example

Find the eigenvalues of $A = \begin{bmatrix} -13 & -8 & -4 \\ 12 & 7 & 4 \\ 24 & 16 & 7 \end{bmatrix}$.

Recall A is similar to the diagonal matrix B below.

$$\begin{bmatrix} -6 & -4 & -1 \\ -3 & -2 & -1 \\ 5 & 3 & 1 \end{bmatrix} \begin{bmatrix} -13 & -8 & -4 \\ 12 & 7 & 4 \\ 24 & 16 & 7 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ -2 & -1 & -3 \\ 1 & -2 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

Since B is a triangular (diagonal) matrix, the eigenvalues of B are $\lambda = -1, 3$, where $\lambda = -1$ has multiplicity 2. Thus A has eigenvalues $\lambda = -1, 3$, where $\lambda = -1$ has multiplicity 2.

Caution

- ▶ If two matrices are similar, then they have the same eigenvalues. **TRUE**

Caution

- ▶ If two matrices are similar, then they have the same eigenvalues. **TRUE**
- ▶ If two matrices have the same eigenvalues, then the two matrices are similar.

Caution

- ▶ If two matrices are similar, then they have the same eigenvalues. **TRUE**
- ▶ If two matrices have the same eigenvalues, then the two matrices are similar. **FALSE!**

Example

$$\begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 1 \\ 0 & 5 \end{bmatrix}$$

Both matrices have eigenvalues $\lambda = 1, 5$, however they are **not similar** matrices.

Similarity is **not** the same as row equivalence. The two matrices above are row equivalent, but they are not similar since there is no matrix P such that $P^{-1}AP = B$.