


Dot Product, Length, and Orthogonality

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Calculating the Length of a Vector




$$\mathbf{v} = (-4, -5)$$

Use the Pythagorean theorem:

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2} = \sqrt{(-4)^2 + (-5)^2} = \sqrt{41}$$

Calculating the Length of a Vector



$$\mathbf{v} = (-4, -5)$$


Use the Pythagorean theorem:

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2} = \sqrt{(-4)^2 + (-5)^2} = \sqrt{41}$$

Let $\mathbf{w} = (1, 2, 3)$ be a vector in \mathbb{R}^3 .

$$\|\mathbf{w}\| = \sqrt{w_1^2 + w_2^2 + w_3^2} = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

Calculating the Length of a Vector



$$\mathbf{v} = (-4, -5)$$

Use the Pythagorean theorem:

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2} = \sqrt{(-4)^2 + (-5)^2} = \sqrt{41}$$

Let $\mathbf{w} = (1, 2, 3)$ be a vector in \mathbb{R}^3 .

$$\|\mathbf{w}\| = \sqrt{w_1^2 + w_2^2 + w_3^2} = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

The **length** (or **norm**) of \mathbf{v} is the nonnegative scalar $\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$.

The Dot Product

Definition

Let \mathbf{u} and \mathbf{v} denote two $n \times 1$ column vectors in \mathbb{R}^n . The **dot product** of \mathbf{u} and \mathbf{v} is the scalar

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n = \sum_{i=1}^n u_i v_i$$

The Dot Product

Definition

Let \mathbf{u} and \mathbf{v} denote two $n \times 1$ column vectors in \mathbb{R}^n . The **dot product** of \mathbf{u} and \mathbf{v} is the scalar

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n = \sum_{i=1}^n u_i v_i = \mathbf{u}^T \mathbf{v} = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}.$$

The Dot Product

Definition

Let \mathbf{u} and \mathbf{v} denote two $n \times 1$ column vectors in \mathbb{R}^n . The **dot product** of \mathbf{u} and \mathbf{v} is the scalar

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n = \sum_{i=1}^n u_i v_i = \mathbf{u}^T \mathbf{v} = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}.$$

Example.

If $\mathbf{u} = \begin{bmatrix} 2 \\ -5 \\ 1 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} 6 \\ 2 \\ -7 \end{bmatrix}$ in \mathbb{R}^3 , then $\mathbf{u} \cdot \mathbf{v} = (2)(6) + (-5)(2) + (1)(-7) = -5$.

Properties of the Dot Product

Theorem

Let \mathbf{u} , \mathbf{v} , and \mathbf{w} be vectors in \mathbb{R}^n , and let c be a scalar. Then

- (a) $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$.
- (b) $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = (\mathbf{u} \cdot \mathbf{w}) + (\mathbf{v} \cdot \mathbf{w})$.
- (c) $(c\mathbf{u}) \cdot \mathbf{v} = c(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (c\mathbf{v})$.
- (d) $\mathbf{v} \cdot \mathbf{v} \geq 0$, and $\mathbf{v} \cdot \mathbf{v} = 0$ if and only if $\mathbf{v} = \mathbf{0}$.

Properties of the Dot Product

Theorem


Let \mathbf{u} , \mathbf{v} , and \mathbf{w} be vectors in \mathbb{R}^n , and let c be a scalar. Then

- (a) $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$.
- (b) $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = (\mathbf{u} \cdot \mathbf{w}) + (\mathbf{v} \cdot \mathbf{w})$.
- (c) $(c\mathbf{u}) \cdot \mathbf{v} = c(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (c\mathbf{v})$.
- (d) $\mathbf{v} \cdot \mathbf{v} \geq 0$, and $\mathbf{v} \cdot \mathbf{v} = 0$ if and only if $\mathbf{v} = \mathbf{0}$.

Definition


The **length** (or **norm**) $\|\mathbf{v}\|$ of \mathbf{v} is the nonnegative scalar $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}}$.

Scaling Vectors



1. Find a vector \mathbf{w} that has the same direction as $\mathbf{v} = (-4, -5)$ but is twice as long.
2. Find a vector \mathbf{u} that has the same direction as \mathbf{v} that has length 1.


Scaling Vectors



1. Find a vector \mathbf{w} that has the same direction as $\mathbf{v} = (-4, -5)$ but is twice as long.
2. Find a vector \mathbf{u} that has the same direction as \mathbf{v} that has length 1.

Let c be a scalar. Then $\|c\mathbf{v}\| = |c| \|\mathbf{v}\|$.

Scaling Vectors



1. Find a vector \mathbf{w} that has the same direction as $\mathbf{v} = (-4, -5)$ but is twice as long.
2. Find a vector \mathbf{u} that has the same direction as \mathbf{v} that has length 1.

Let c be a scalar. Then $\|c\mathbf{v}\| = |c| \|\mathbf{v}\|$.

1. Let $\mathbf{w} = 2(\mathbf{v}) = \begin{bmatrix} -8 \\ -10 \end{bmatrix}$ has the same direction as \mathbf{v} and is twice as long.
2. Note that $\|\mathbf{v}\| = \sqrt{41}$. If we scale \mathbf{v} by a factor of $1/\|\mathbf{v}\|$, then we will obtain a vector in the same direction as \mathbf{v} with magnitude equal to 1:

$$\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v} = \frac{1}{\sqrt{41}} \mathbf{v} = \begin{bmatrix} -4/\sqrt{41} \\ -5/\sqrt{41} \end{bmatrix}; \quad \|\mathbf{u}\| = 1$$

Normalizing Vectors

Definition

A vector \mathbf{u} whose length is 1 is called a **unit vector**. The process of creating a unit vector \mathbf{u} in the same direction as a vector \mathbf{v} is called **normalizing** \mathbf{v} . We have

$$\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v}.$$

Normalizing Vectors

Definition

A vector \mathbf{u} whose length is 1 is called a **unit vector**. The process of creating a unit vector \mathbf{u} in the same direction as a vector \mathbf{v} is called **normalizing** \mathbf{v} . We have

$$\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v}.$$

Example.

Compute the unit vector in the direction of $\mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$.

Normalizing Vectors

Definition

A vector \mathbf{u} whose length is 1 is called a **unit vector**. The process of creating a unit vector \mathbf{u} in the same direction as a vector \mathbf{v} is called **normalizing** \mathbf{v} . We have

$$\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v}.$$


Example.

Compute the unit vector in the direction of $\mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$. $\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v} = \frac{1}{\sqrt{14}} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{14} \\ 2/\sqrt{14} \\ 3/\sqrt{14} \end{bmatrix}.$

Distance between two vectors

Example


Compute the distance between $\mathbf{u} = \begin{bmatrix} 6 \\ 6 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$.



Distance between two vectors

Example

Compute the distance between $\mathbf{u} = \begin{bmatrix} 6 \\ 6 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$.




We want to find the length of vector \mathbf{w} where $\mathbf{v} + \mathbf{w} = \mathbf{u}$. Thus,

$$\mathbf{w} = \mathbf{u} - \mathbf{v} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}.$$

$$\text{dist}(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\| = \sqrt{2^2 + 4^2} = \sqrt{20}$$

Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .




Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.

► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$



Orthogonality


Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.

► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$

$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$$



Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .


► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.

► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$

$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$$

$$\mathbf{u} \cdot (\mathbf{u} - \mathbf{v}) - \mathbf{v} \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) + \mathbf{v} \cdot (\mathbf{u} + \mathbf{v})$$



Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.


► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$

$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$$

$$\mathbf{u} \cdot (\mathbf{u} - \mathbf{v}) - \mathbf{v} \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) + \mathbf{v} \cdot (\mathbf{u} + \mathbf{v})$$

$$(\mathbf{u} \cdot \mathbf{u}) - (\mathbf{u} \cdot \mathbf{v}) - (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v}) = (\mathbf{u} \cdot \mathbf{u}) + (\mathbf{u} \cdot \mathbf{v}) + (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v})$$



Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.

► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.


$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$

$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$$

$$\mathbf{u} \cdot (\mathbf{u} - \mathbf{v}) - \mathbf{v} \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) + \mathbf{v} \cdot (\mathbf{u} + \mathbf{v})$$

$$(\mathbf{u} \cdot \mathbf{u}) - (\mathbf{u} \cdot \mathbf{v}) - (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v}) = (\mathbf{u} \cdot \mathbf{u}) + (\mathbf{u} \cdot \mathbf{v}) + (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v})$$

$$\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2(\mathbf{u} \cdot \mathbf{v}) = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2(\mathbf{u} \cdot \mathbf{v})$$



Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.

► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$


$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$$

$$\mathbf{u} \cdot (\mathbf{u} - \mathbf{v}) - \mathbf{v} \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) + \mathbf{v} \cdot (\mathbf{u} + \mathbf{v})$$

$$(\mathbf{u} \cdot \mathbf{u}) - (\mathbf{u} \cdot \mathbf{v}) - (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v}) = (\mathbf{u} \cdot \mathbf{u}) + (\mathbf{u} \cdot \mathbf{v}) + (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v})$$

$$\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2(\mathbf{u} \cdot \mathbf{v}) = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2(\mathbf{u} \cdot \mathbf{v})$$

$$-4(\mathbf{u} \cdot \mathbf{v}) = 0$$



Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.

► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$

$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$$


$$\mathbf{u} \cdot (\mathbf{u} - \mathbf{v}) - \mathbf{v} \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) + \mathbf{v} \cdot (\mathbf{u} + \mathbf{v})$$

$$(\mathbf{u} \cdot \mathbf{u}) - (\mathbf{u} \cdot \mathbf{v}) - (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v}) = (\mathbf{u} \cdot \mathbf{u}) + (\mathbf{u} \cdot \mathbf{v}) + (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v})$$

$$\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2(\mathbf{u} \cdot \mathbf{v}) = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2(\mathbf{u} \cdot \mathbf{v})$$

$$-4(\mathbf{u} \cdot \mathbf{v}) = 0$$

$$\mathbf{u} \cdot \mathbf{v} = 0$$



Orthogonality

Consider two perpendicular vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

► We see that $\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} - (-\mathbf{v})\|$.

► Thus $\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} - (-\mathbf{v})\|^2$.

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u} + \mathbf{v}\|^2$$

$$(\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$$


$$\mathbf{u} \cdot (\mathbf{u} - \mathbf{v}) - \mathbf{v} \cdot (\mathbf{u} - \mathbf{v}) = \mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) + \mathbf{v} \cdot (\mathbf{u} + \mathbf{v})$$

$$(\mathbf{u} \cdot \mathbf{u}) - (\mathbf{u} \cdot \mathbf{v}) - (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v}) = (\mathbf{u} \cdot \mathbf{u}) + (\mathbf{u} \cdot \mathbf{v}) + (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{v} \cdot \mathbf{v})$$

$$\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2(\mathbf{u} \cdot \mathbf{v}) = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + 2(\mathbf{u} \cdot \mathbf{v})$$

$$-4(\mathbf{u} \cdot \mathbf{v}) = 0$$

$$\mathbf{u} \cdot \mathbf{v} = 0$$



Definition

Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if and only if $\mathbf{u} \cdot \mathbf{v} = 0$.

Orthogonality

Definition

Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if and only if $\mathbf{u} \cdot \mathbf{v} = 0$.

Example.

Are $\mathbf{u} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$ orthogonal?

Orthogonality

Definition

Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if and only if $\mathbf{u} \cdot \mathbf{v} = 0$.

Example.

Are $\mathbf{u} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$ orthogonal?

$$\mathbf{u} \cdot \mathbf{v} = 5(-1) + 2(2) = -1 \neq 0.$$

\mathbf{u} and \mathbf{v} are **not** orthogonal.

Orthogonality

Definition

Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if and only if $\mathbf{u} \cdot \mathbf{v} = 0$.

Example.

Are $\mathbf{u} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$ orthogonal?

$$\mathbf{u} \cdot \mathbf{v} = 5(-1) + 2(2) = -1 \neq 0.$$

\mathbf{u} and \mathbf{v} are **not** orthogonal.

Are $\mathbf{u} = \begin{bmatrix} 4 \\ -7 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -5 \\ -2 \\ 3 \end{bmatrix}$ orthogonal?

Orthogonality

Definition

Two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if and only if $\mathbf{u} \cdot \mathbf{v} = 0$.

Example.

Are $\mathbf{u} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$ orthogonal?

$$\mathbf{u} \cdot \mathbf{v} = 5(-1) + 2(2) = -1 \neq 0.$$

\mathbf{u} and \mathbf{v} are **not** orthogonal.

Are $\mathbf{u} = \begin{bmatrix} 4 \\ -7 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -5 \\ -2 \\ 3 \end{bmatrix}$ orthogonal?

$$\begin{aligned} \mathbf{u} \cdot \mathbf{v} &= 4(-5) + (-7)(-2) + 2(3) \\ &= -20 + 14 + 6 = 0. \end{aligned}$$

\mathbf{u} and \mathbf{v} are **orthogonal**.


Orthogonal Complements 1

Definition

If a vector \mathbf{z} is orthogonal to every vector in a subspace W of \mathbb{R}^n , then \mathbf{z} is said to be orthogonal to W . The set of all vectors \mathbf{z} that are orthogonal to W is called the orthogonal complement of W and is denoted W^\perp .

Consider the plane $W = \text{Span}\{\mathbf{v}, \mathbf{w}\}$ in \mathbb{R}^3 plotted in the figure to the right.

- ▶ The vector \mathbf{z} is orthogonal to W .
- ▶ The orthogonal complement $W^\perp = \{c\mathbf{z} : \text{for any scalar } c\}$.



Orthogonal Complements 2

Theorem

Let W be a subspace of \mathbb{R}^n . Then W^\perp is a **subspace** of \mathbb{R}^n .

Orthogonal Complements 2

Theorem

Let W be a subspace of \mathbb{R}^n . Then W^\perp is a **subspace** of \mathbb{R}^n .

Proof. We verify the following three properties.

1. W^\perp is **nonempty**. Note that $\mathbf{0}$ is orthogonal to every vector in \mathbb{R}^n , and so $\mathbf{0}$ is orthogonal to every vector in W . Thus, $\mathbf{0}$ is in W^\perp .

Orthogonal Complements 2

Theorem

Let W be a subspace of \mathbb{R}^n . Then W^\perp is a **subspace** of \mathbb{R}^n .

Proof. We verify the following three properties.

1. W^\perp is **nonempty**. Note that $\mathbf{0}$ is orthogonal to every vector in \mathbb{R}^n , and so $\mathbf{0}$ is orthogonal to every vector in W . Thus, $\mathbf{0}$ is in W^\perp .
2. W^\perp is **closed under addition**. Let \mathbf{u} and \mathbf{v} be vectors in W^\perp . Let \mathbf{w} be any vector in W . Then $\mathbf{w} \cdot (\mathbf{u} + \mathbf{v}) = (\mathbf{w} \cdot \mathbf{u}) + (\mathbf{w} \cdot \mathbf{v}) = 0 + 0 = 0$. Thus, $\mathbf{u} + \mathbf{v}$ is in W^\perp .

Orthogonal Complements 2

Theorem

Let W be a subspace of \mathbb{R}^n . Then W^\perp is a **subspace** of \mathbb{R}^n .

Proof. We verify the following three properties.

1. W^\perp is nonempty. Note that $\mathbf{0}$ is orthogonal to every vector in \mathbb{R}^n , and so $\mathbf{0}$ is orthogonal to every vector in W . Thus, $\mathbf{0}$ is in W^\perp .
2. W^\perp is closed under addition. Let \mathbf{u} and \mathbf{v} be vectors in W^\perp . Let \mathbf{w} be any vector in W . Then $\mathbf{w} \cdot (\mathbf{u} + \mathbf{v}) = (\mathbf{w} \cdot \mathbf{u}) + (\mathbf{w} \cdot \mathbf{v}) = 0 + 0 = 0$. Thus, $\mathbf{u} + \mathbf{v}$ is in W^\perp .
3. W^\perp is closed under scalar multiplication. Let \mathbf{u} be a vector in W^\perp , and let c be a scalar. Let \mathbf{w} be any vector in W . Then $\mathbf{w} \cdot (c\mathbf{u}) = c(\mathbf{w} \cdot \mathbf{u}) = c0 = 0$. Thus, $c\mathbf{u}$ is in W^\perp .

Orthogonal Complements 2

Theorem

Let W be a subspace of \mathbb{R}^n . Then W^\perp is a **subspace** of \mathbb{R}^n .

Proof. We verify the following three properties.

1. W^\perp is **nonempty**. Note that $\mathbf{0}$ is orthogonal to every vector in \mathbb{R}^n , and so $\mathbf{0}$ is orthogonal to every vector in W . Thus, $\mathbf{0}$ is in W^\perp .
2. W^\perp is **closed under addition**. Let \mathbf{u} and \mathbf{v} be vectors in W^\perp . Let \mathbf{w} be any vector in W . Then $\mathbf{w} \cdot (\mathbf{u} + \mathbf{v}) = (\mathbf{w} \cdot \mathbf{u}) + (\mathbf{w} \cdot \mathbf{v}) = 0 + 0 = 0$. Thus, $\mathbf{u} + \mathbf{v}$ is in W^\perp .
3. W^\perp is **closed under scalar multiplication**. Let \mathbf{u} be a vector in W^\perp , and let c be a scalar. Let \mathbf{w} be any vector in W . Then $\mathbf{w} \cdot (c\mathbf{u}) = c(\mathbf{w} \cdot \mathbf{u}) = c0 = 0$. Thus, $c\mathbf{u}$ is in W^\perp .

Thus, W^\perp is a **subspace** of \mathbb{R}^n . □

Orthogonal Complements 3

Theorem

Let W be a subspace of \mathbb{R}^n . A vector \mathbf{z} is in W^\perp if and only if \mathbf{z} is orthogonal to **every** vector in a set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ that **spans** W .

Orthogonal Complements 3

Theorem

Let W be a subspace of \mathbb{R}^n . A vector \mathbf{z} is in W^\perp if and only if \mathbf{z} is orthogonal to **every** vector in a set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ that **spans** W .

Proof. (\Rightarrow) If \mathbf{z} is in W^\perp , then \mathbf{z} is orthogonal to each vector in W , including $\mathbf{v}_1, \dots, \mathbf{v}_p$.

Orthogonal Complements 3

Theorem

Let W be a subspace of \mathbb{R}^n . A vector \mathbf{z} is in W^\perp if and only if \mathbf{z} is orthogonal to **every** vector in a set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ that **spans** W .

Proof. (\Rightarrow) If \mathbf{z} is in W^\perp , then \mathbf{z} is orthogonal to each vector in W , including $\mathbf{v}_1, \dots, \mathbf{v}_p$.

(\Leftarrow) Assume that \mathbf{z} is orthogonal to each \mathbf{v}_i .

Let \mathbf{w} be any vector in W . Since $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ spans W , there exist scalars c_1, \dots, c_p such that $\mathbf{w} = c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p$. We compute

$$\mathbf{z} \cdot \mathbf{w} = \mathbf{z} \cdot (c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p)$$

Orthogonal Complements 3

Theorem

Let W be a subspace of \mathbb{R}^n . A vector \mathbf{z} is in W^\perp if and only if \mathbf{z} is orthogonal to **every** vector in a set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ that **spans** W .

Proof. (\Rightarrow) If \mathbf{z} is in W^\perp , then \mathbf{z} is orthogonal to each vector in W , including $\mathbf{v}_1, \dots, \mathbf{v}_p$.

(\Leftarrow) Assume that \mathbf{z} is orthogonal to each \mathbf{v}_i .

Let \mathbf{w} be any vector in W . Since $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ spans W , there exist scalars c_1, \dots, c_p such that $\mathbf{w} = c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p$. We compute

$$\mathbf{z} \cdot \mathbf{w} = \mathbf{z} \cdot (c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p) = c_1(\mathbf{z} \cdot \mathbf{v}_1) + \dots + c_p(\mathbf{z} \cdot \mathbf{v}_p) = 0.$$

Thus, \mathbf{z} is orthogonal to every vector in W , and hence \mathbf{z} is in W^\perp . □

Matrix Subspaces

Let $A = \begin{bmatrix} 1 & 0 & -1 \\ -2 & 0 & 2 \end{bmatrix}$. Find bases for Null A , Col A , Row A , and Null A^T .

Matrix Subspaces

Let $A = \begin{bmatrix} 1 & 0 & -1 \\ -2 & 0 & 2 \end{bmatrix}$. Find bases for Null A , Col A , Row A , and Null A^T .

We have $\begin{bmatrix} 1 & 0 & -1 \\ -2 & 0 & 2 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 0 & 0 \end{bmatrix}$.

Thus we have bases

$$\text{Null } A = \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

$$\text{Row } A = \{[1 \quad 0 \quad -1]\}$$

$$\text{Col } A = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix} \right\}$$

Matrix Subspaces

Let $A = \begin{bmatrix} 1 & 0 & -1 \\ -2 & 0 & 2 \end{bmatrix}$. Find bases for Null A , Col A , Row A , and Null A^T .

We have $\begin{bmatrix} 1 & 0 & -1 \\ -2 & 0 & 2 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 0 & 0 \end{bmatrix}$.

Thus we have bases

$$\text{Null } A = \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

$$\text{Row } A = \{[1 \quad 0 \quad -1]\}$$

$$\text{Col } A = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix} \right\}$$

We have $A^T = \begin{bmatrix} 1 & -2 \\ 0 & 0 \\ -1 & 2 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & -2 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$.

Thus we have basis

$$\text{Null } A^T = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$$

Orthogonal Complement of Matrix Subspaces

Example


Let $A = \begin{bmatrix} 1 & 0 & -1 \\ -2 & 0 & 2 \end{bmatrix}$.

1. Find a basis for Null A , Col A and Row A
2. Find a basis for Null A^T .


We have bases

$$\text{Null } A = \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\} \quad \text{Row } A = \left\{ \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \right\}$$

$$\text{Col } A = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix} \right\} \quad \text{Null } A^T = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}.$$



Subspaces Nul A and Row A



Subspaces Nul A^T and Col A

Null, Col, and Row Spaces Revisited

Theorem

Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A . The orthogonal complement of the column space of A is the null space of A^T .

$$(\text{Row } A)^\perp = \text{Null } A \quad \text{and} \quad (\text{Col } A)^\perp = \text{Null } A^T$$

Null, Col, and Row Spaces Revisited

Theorem

Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A . The orthogonal complement of the column space of A is the null space of A^T .

$$(\text{Row } A)^\perp = \text{Null } A \quad \text{and} \quad (\text{Col } A)^\perp = \text{Null } A^T$$

Proof.

Let \mathbf{x} be in $(\text{Row } A)^\perp$. $A\mathbf{x} = \begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$

Null, Col, and Row Spaces Revisited

Theorem

Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A . The orthogonal complement of the column space of A is the null space of A^T .

$$(\text{Row } A)^\perp = \text{Null } A \quad \text{and} \quad (\text{Col } A)^\perp = \text{Null } A^T$$

Proof.

$$\text{Let } \mathbf{x} \text{ be in } (\text{Row } A)^\perp. \quad A\mathbf{x} = \begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \mathbf{r}_1 \cdot \mathbf{x} \\ \mathbf{r}_2 \cdot \mathbf{x} \\ \vdots \\ \mathbf{r}_m \cdot \mathbf{x} \end{bmatrix}$$

Null, Col, and Row Spaces Revisited

Theorem

Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A . The orthogonal complement of the column space of A is the null space of A^T .

$$(\text{Row } A)^\perp = \text{Null } A \quad \text{and} \quad (\text{Col } A)^\perp = \text{Null } A^T$$

Proof.

$$\text{Let } \mathbf{x} \text{ be in } (\text{Row } A)^\perp. \quad A\mathbf{x} = \begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \mathbf{r}_1 \cdot \mathbf{x} \\ \mathbf{r}_2 \cdot \mathbf{x} \\ \vdots \\ \mathbf{r}_m \cdot \mathbf{x} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

Null, Col, and Row Spaces Revisited

Theorem

Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A . The orthogonal complement of the column space of A is the null space of A^T .

$$(\text{Row } A)^\perp = \text{Null } A \quad \text{and} \quad (\text{Col } A)^\perp = \text{Null } A^T$$

Proof.

$$\text{Let } \mathbf{x} \text{ be in } (\text{Row } A)^\perp. \quad A\mathbf{x} = \begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \mathbf{r}_1 \cdot \mathbf{x} \\ \mathbf{r}_2 \cdot \mathbf{x} \\ \vdots \\ \mathbf{r}_m \cdot \mathbf{x} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

Thus \mathbf{x} is in $\text{Null } A$.

Similarly, if \mathbf{x} is in $\text{Null } A$, then it satisfies the equation $A\mathbf{x} = \mathbf{0}$, which implies \mathbf{x} is orthogonal to each row of A . Since the rows of A span $\text{Row } A$ and by a previous theorem, \mathbf{x} is orthogonal to each vector in $\text{Row } A$. Thus, \mathbf{x} is in $(\text{Row } A)^\perp$.

Null, Col, and Row Spaces Revisited

Theorem

Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A . The orthogonal complement of the column space of A is the null space of A^T .

$$(\text{Row } A)^\perp = \text{Null } A \quad \text{and} \quad (\text{Col } A)^\perp = \text{Null } A^T$$

Proof.

$$\text{Let } \mathbf{x} \text{ be in } (\text{Row } A)^\perp. \quad A\mathbf{x} = \begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_m \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \mathbf{r}_1 \cdot \mathbf{x} \\ \mathbf{r}_2 \cdot \mathbf{x} \\ \vdots \\ \mathbf{r}_m \cdot \mathbf{x} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

Thus \mathbf{x} is in $\text{Null } A$.

Similarly, if \mathbf{x} is in $\text{Null } A$, then it satisfies the equation $A\mathbf{x} = \mathbf{0}$, which implies \mathbf{x} is orthogonal to each row of A . Since the rows of A span $\text{Row } A$ and by a previous theorem, \mathbf{x} is orthogonal to each vector in $\text{Row } A$. Thus, \mathbf{x} is in $(\text{Row } A)^\perp$. Therefore we have $(\text{Row } A)^\perp = \text{Null } A$.

Null, Col, and Row Spaces Revisited

Theorem

Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A . The orthogonal complement of the column space of A is the null space of A^T .

$$(\text{Row } A)^\perp = \text{Null } A \quad \text{and} \quad (\text{Col } A)^\perp = \text{Null } A^T$$

Proof.

The second statement follows by noting that $\text{Col } A = \text{Row } A^T$:

$$(\text{Col } A)^\perp = (\text{Row } A^T)^\perp = \text{Null } A^T$$

