

# Eigenvalues and Eigenvectors

## Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

# Introduction

The concepts of **eigenvalues** and **eigenvectors** of a matrix are useful throughout pure and applied mathematics.

- ▶ Eigenvalues are used to study difference equations and continuous dynamical systems.
- ▶ They provide critical information in engineering design, and they arise naturally in such fields as physics and chemistry.

# Introduction

Consider the linear transformation  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  where  $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 0 & -2 \\ -1 & 2 \end{bmatrix} \mathbf{x}$ .

How do  $\mathbf{x}$  and  $A\mathbf{x}$  relate?

<https://www.desmos.com/calculator/jovijh4fad>

# Introduction

Consider the linear transformation  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  where  $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 0 & -2 \\ -1 & 2 \end{bmatrix} \mathbf{x}$ .

How do  $\mathbf{x}$  and  $A\mathbf{x}$  relate?

<https://www.desmos.com/calculator/jovijh4fad>

a) Compute  $A\mathbf{u}$ , where  $\mathbf{u} = \begin{bmatrix} 1 + \sqrt{3} \\ 1 \end{bmatrix}$ .

b) Compute  $A\mathbf{v}$ , where  $\mathbf{v} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ .

$$\begin{bmatrix} 0 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 1 + \sqrt{3} \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 - \sqrt{3} \end{bmatrix} = (1 - \sqrt{3})\mathbf{u}.$$

$$\begin{bmatrix} 0 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 3 \end{bmatrix} \neq c\mathbf{v}.$$

# Eigenvalues and Eigenvectors

Consider the matrix  $A = \begin{bmatrix} 0 & -2 \\ -1 & 2 \end{bmatrix}$  and vectors  $\mathbf{u} = \begin{bmatrix} 1 + \sqrt{3} \\ 1 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ .

$$\begin{bmatrix} 0 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 1 + \sqrt{3} \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 - \sqrt{3} \end{bmatrix} = (1 - \sqrt{3})\mathbf{u}.$$

$$\begin{bmatrix} 0 & -2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 3 \end{bmatrix} \neq c\mathbf{v}.$$

## Definition

An **eigenvalue** of an  $n \times n$  matrix  $A$  is a scalar  $\lambda$  such that there exists a **nonzero** vector  $\mathbf{x}$  where  $A\mathbf{x} = \lambda\mathbf{x}$ . The nonzero vector  $\mathbf{x}$  is called an **eigenvector corresponding to  $\lambda$** .

## Example

Show that  $\lambda = 4$  is an eigenvalue of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

- If  $\lambda = 4$  is an eigenvalue of  $A$ , then we know that  $A\mathbf{x} = 4\mathbf{x}$  has a **nonzero** solution.

## Example

Show that  $\lambda = 4$  is an eigenvalue of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

- ▶ If  $\lambda = 4$  is an eigenvalue of  $A$ , then we know that  $A\mathbf{x} = 4\mathbf{x}$  has a **nonzero** solution.
- ▶ This gives the equivalent equation  $A\mathbf{x} - 4\mathbf{x} = \mathbf{0}$ .

## Example

Show that  $\lambda = 4$  is an eigenvalue of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

- ▶ If  $\lambda = 4$  is an eigenvalue of  $A$ , then we know that  $A\mathbf{x} = 4\mathbf{x}$  has a **nonzero** solution.
- ▶ This gives the equivalent equation  $A\mathbf{x} - 4\mathbf{x} = \mathbf{0}$ .
- ▶ Note that we can represent the scalar product  $4\mathbf{x} = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \mathbf{x} = 4I_2\mathbf{x}$ .
- ▶ We solve the equation  $A\mathbf{x} - 4\mathbf{x} = A\mathbf{x} - 4I_2\mathbf{x} = (A - 4I)\mathbf{x} = \mathbf{0}$ .

$$(A - 4I)\mathbf{x} = \left( \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} - \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \right) \mathbf{x} = \begin{bmatrix} -4 & -2 \\ -4 & -2 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$



## Example Continued

Show that  $\lambda = 4$  is an eigenvalue of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

$$(A - 4I)\mathbf{x} = \left( \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} - \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \right) \mathbf{x} = \begin{bmatrix} -4 & -2 \\ -4 & -2 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

## Example Continued

Show that  $\lambda = 4$  is an eigenvalue of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

$$(A - 4I)\mathbf{x} = \left( \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} - \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \right) \mathbf{x} = \begin{bmatrix} -4 & -2 \\ -4 & -2 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

We have  $\begin{bmatrix} -4 & -2 & 0 \\ -4 & -2 & 0 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{bmatrix}$ , which gives the solution set  $\mathbf{x} = x_2 \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$ .

## Example Continued

Show that  $\lambda = 4$  is an eigenvalue of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

$$(A - 4I)\mathbf{x} = \left( \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} - \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \right) \mathbf{x} = \begin{bmatrix} -4 & -2 \\ -4 & -2 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

We have  $\begin{bmatrix} -4 & -2 & 0 \\ -4 & -2 & 0 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{bmatrix}$ , which gives the solution set  $\mathbf{x} = x_2 \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$ .

Since the equation  $A\mathbf{x} = 4\mathbf{x}$  has a **nontrivial solution**,  $\lambda = 4$  is an **eigenvalue**.

We see that  $\mathbf{v} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$  is an **eigenvector**. In fact, any nonzero scalar multiple of  $\mathbf{v}$ , such as  $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$ , is also an eigenvector corresponding to  $\lambda = 4$ .

# Setting Up A Matrix Equation

A scalar  $\lambda$  is called an **eigenvalue** of  $A$  if there is a nontrivial solution  $\mathbf{x}$  of  $A\mathbf{x} = \lambda\mathbf{x}$ .

In general, we can rewrite  $A\mathbf{x} = \lambda\mathbf{x}$  as a homogeneous matrix equation:

$$(A - \lambda I)\mathbf{x} = \mathbf{0}.$$

The set of all solutions to  $(A - \lambda I)\mathbf{x} = \mathbf{0}$  is called the **eigenspace** of  $A$  corresponding to the eigenvalue  $\lambda$ .

The eigenspace of  $\begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$  corresponding to  $\lambda = 4$  is  $\text{Span} \left\{ \begin{bmatrix} -1 \\ 2 \end{bmatrix} \right\}$ .

## Finding a Basis for the Eigenspace

Let  $A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$ . Find a basis for the eigenspace corresponding to the eigenvalue  $\lambda = 2$ .

## Finding a Basis for the Eigenspace

Let  $A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$ . Find a basis for the eigenspace corresponding to the eigenvalue  $\lambda = 2$ .

$$A - 2I = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} - \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

The augmented matrix for the equation  $(A - 2I)\mathbf{x} = \mathbf{0}$  is

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ -1 & 1 & 1 & 0 \\ -1 & 1 & 1 & 0 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

## Finding a Basis for the Eigenspace

Let  $A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$ . Find a basis for the eigenspace corresponding to the eigenvalue  $\lambda = 2$ .

$$A - 2I = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} - \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

The augmented matrix for the equation  $(A - 2I)\mathbf{x} = \mathbf{0}$  is

$$\left[ \begin{array}{cccc} 0 & 0 & 0 & 0 \\ -1 & 1 & 1 & 0 \\ -1 & 1 & 1 & 0 \end{array} \right] \xrightarrow{\text{RREF}} \left[ \begin{array}{cccc} 1 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Thus we have the solution set

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_2 + x_3 \\ x_2 \\ x_3 \end{bmatrix}.$$


A basis for the eigenspace corresponding to  $\lambda = 2$  is

$$\mathcal{B}_{\lambda=2} = \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

# Geometric Picture

Let  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ . Find bases for the eigenspaces corresponding to the eigenvalues  $\lambda = 4$  and  $\lambda = -2$ .

$$\mathcal{B}_{\lambda=4} = \left\{ \begin{bmatrix} -1 \\ 2 \end{bmatrix} \right\} \quad \text{and} \quad \mathcal{B}_{\lambda=-2} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}.$$






# Geometric Picture


Let  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ . Find bases for the eigenspaces corresponding to the eigenvalues  $\lambda = 4$  and  $\lambda = -2$ .

$$\mathcal{B}_{\lambda=4} = \left\{ \begin{bmatrix} -1 \\ 2 \end{bmatrix} \right\} \quad \text{and} \quad \mathcal{B}_{\lambda=-2} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}.$$



Let  $A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$ . Find a basis for the eigenspace corresponding to the eigenvalue  $\lambda = 2$ .

$$\mathcal{B}_{\lambda=2} = \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}.$$



## Finding Eigenvalues of $A^2$ , $A^3$ , $A^k$

Let  $A$  be an  $n \times n$  matrix with eigenvalue  $\lambda$  and corresponding eigenvector  $\mathbf{x}$ .

1. Show that  $\lambda^2$  is an eigenvalue for  $A^2$ .

## Finding Eigenvalues of $A^2$ , $A^3$ , $A^k$

Let  $A$  be an  $n \times n$  matrix with eigenvalue  $\lambda$  and corresponding eigenvector  $\mathbf{x}$ .

1. Show that  $\lambda^2$  is an eigenvalue for  $A^2$ .

We have

$$A^2\mathbf{x} = A(A\mathbf{x}) = A(\lambda\mathbf{x}) = \lambda(A\mathbf{x}) = \lambda^2\mathbf{x}.$$

We see  $\lambda^2$  is an eigenvalue of the matrix  $A^2 = AA$ .

## Finding Eigenvalues of $A^2$ , $A^3$ , $A^k$

Let  $A$  be an  $n \times n$  matrix with eigenvalue  $\lambda$  and corresponding eigenvector  $\mathbf{x}$ .

1. Show that  $\lambda^2$  is an eigenvalue for  $A^2$ .

We have

$$A^2\mathbf{x} = A(A\mathbf{x}) = A(\lambda\mathbf{x}) = \lambda(A\mathbf{x}) = \lambda^2\mathbf{x}.$$

We see  $\lambda^2$  is an eigenvalue of the matrix  $A^2 = AA$ .

2. Show that  $\lambda^3$  is an eigenvalue for  $A^3$ .

We have

$$A^3\mathbf{x} = A(A^2\mathbf{x}) = A(\lambda^2\mathbf{x}) = \lambda^2(A\mathbf{x}) = \lambda^3\mathbf{x}.$$

We see  $\lambda^3$  is an eigenvalue of the matrix  $A^3$ .

## Finding Eigenvalues of $A^2$ , $A^3$ , $A^k$

Let  $A$  be an  $n \times n$  matrix with eigenvalue  $\lambda$  and corresponding eigenvector  $\mathbf{x}$ .

1. Show that  $\lambda^2$  is an eigenvalue for  $A^2$ .

We have

$$A^2\mathbf{x} = A(A\mathbf{x}) = A(\lambda\mathbf{x}) = \lambda(A\mathbf{x}) = \lambda^2\mathbf{x}.$$

We see  $\lambda^2$  is an eigenvalue of the matrix  $A^2 = AA$ .

2. Show that  $\lambda^3$  is an eigenvalue for  $A^3$ .

We have

$$A^3\mathbf{x} = A(A^2\mathbf{x}) = A(\lambda^2\mathbf{x}) = \lambda^2(A\mathbf{x}) = \lambda^3\mathbf{x}.$$

We see  $\lambda^3$  is an eigenvalue of the matrix  $A^3$ .

In general, if  $\lambda$  is an eigenvalue of  $A$ , then  $\lambda^k$  is an eigenvalue of  $A^k$ , where  $k$  is a positive integer.

## Example 2

Is 5 an **eigenvalue** of  $A = \begin{bmatrix} 6 & -3 & 1 \\ 3 & 0 & 5 \\ 2 & 2 & 6 \end{bmatrix}$ ?

## Example 2

Is 5 an **eigenvalue** of  $A = \begin{bmatrix} 6 & -3 & 1 \\ 3 & 0 & 5 \\ 2 & 2 & 6 \end{bmatrix}$ ?

$$(A - 5I)\mathbf{x} = \begin{bmatrix} 1 & -3 & 1 \\ 3 & -5 & 5 \\ 2 & 2 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

We have  $\begin{bmatrix} 1 & -3 & 1 & 0 \\ 3 & -5 & 5 & 0 \\ 2 & 2 & 1 & 0 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix},$

## Example 2

Is 5 an **eigenvalue** of  $A = \begin{bmatrix} 6 & -3 & 1 \\ 3 & 0 & 5 \\ 2 & 2 & 6 \end{bmatrix}$ ?

$$(A - 5I)\mathbf{x} = \begin{bmatrix} 1 & -3 & 1 \\ 3 & -5 & 5 \\ 2 & 2 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

We have  $\begin{bmatrix} 1 & -3 & 1 & 0 \\ 3 & -5 & 5 & 0 \\ 2 & 2 & 1 & 0 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$ , which has only the **trivial solution**  $\mathbf{x} = \mathbf{0}$ .

Since the equation  $A\mathbf{x} = 5\mathbf{x}$  has only the trivial solution,  $\lambda = 5$  is **NOT** an eigenvalue of  $A$ .



# Eigenvalues of a Triangular Matrix

## Theorem

The eigenvalues of a **triangular matrix** are the entries on its main diagonal.

# Eigenvalues of a Triangular Matrix

## Theorem

The eigenvalues of a **triangular matrix** are the entries on its main diagonal.

**Proof.** Let's consider the **3 × 3** case. Let  $A$  be a  $3 \times 3$  upper triangular matrix. Then

$$A - \lambda I = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix} - \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix} = \begin{bmatrix} a_{11} - \lambda & a_{12} & a_{13} \\ 0 & a_{22} - \lambda & a_{23} \\ 0 & 0 & a_{33} - \lambda \end{bmatrix}.$$

The scalar  $\lambda$  is an eigenvalue of  $A$  if and only if there is a **nontrivial solution** to the homogeneous equation  $(A - \lambda I)\mathbf{x} = \mathbf{0}$ . We have a nontrivial solution when  $(A - \lambda I)$  has at least one **free variable**.

# Eigenvalues of a Triangular Matrix

## Theorem

The eigenvalues of a **triangular matrix** are the entries on its main diagonal.

**Proof.** Let's consider the  $3 \times 3$  case. Let  $A$  be a  $3 \times 3$  upper triangular matrix. Then

$$A - \lambda I = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix} - \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix} = \begin{bmatrix} a_{11} - \lambda & a_{12} & a_{13} \\ 0 & a_{22} - \lambda & a_{23} \\ 0 & 0 & a_{33} - \lambda \end{bmatrix}.$$

The scalar  $\lambda$  is an eigenvalue of  $A$  if and only if there is a **nontrivial solution** to the homogeneous equation  $(A - \lambda I)\mathbf{x} = \mathbf{0}$ . We have a nontrivial solution when  $(A - \lambda I)$  has at least one **free variable**.

- ▶ If we set  $\lambda = a_{11}$ , then  $x_1$  is free since there is no pivot in column 1.
- ▶ If we set  $\lambda = a_{22}$ , then  $x_2$  is free since there is no pivot in column 2 (assuming that  $\lambda \neq a_{11}$ ).
- ▶ If we set  $\lambda = a_{33}$ , then  $x_3$  is free since there is no pivot in column 3 (assuming that  $\lambda \neq a_{11}, a_{22}$ ).

Thus the eigenvalues of  $A$  are  $\lambda = a_{11}$ ,  $a_{22}$ , or  $a_{33}$ . The argument follows similarly for larger matrices and for lower triangular matrices as well. □

# Linear Independence of Eigenvectors

## Theorem

If  $\mathbf{v}_1, \dots, \mathbf{v}_p$  are eigenvectors corresponding to **distinct** eigenvalues  $\lambda_1, \dots, \lambda_p$  of an  $n \times n$  matrix  $A$ , then the set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is **linearly independent**.

# Linear Independence of Eigenvectors

## Theorem

If  $\mathbf{v}_1, \dots, \mathbf{v}_p$  are eigenvectors corresponding to **distinct** eigenvalues  $\lambda_1, \dots, \lambda_p$  of an  $n \times n$  matrix  $A$ , then the set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is **linearly independent**.

**Proof.** Suppose  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is linearly dependent. Thus there are scalars  $c_i$ , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such  $c_i$ s that are not all zero, choose the scalars such that the **fewest**  $c_i$ s are nonzero. Since the  $\mathbf{v}_i$ s are eigenvectors and not the zero vector, at least two of the  $c_i$ s must be nonzero. Assume that  $c_j$  and  $c_k$  are nonzero.

# Linear Independence of Eigenvectors

## Theorem

If  $\mathbf{v}_1, \dots, \mathbf{v}_p$  are eigenvectors corresponding to **distinct** eigenvalues  $\lambda_1, \dots, \lambda_p$  of an  $n \times n$  matrix  $A$ , then the set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is **linearly independent**.

**Proof.** Suppose  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is linearly dependent. Thus there are scalars  $c_i$ , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such  $c_i$ s that are not all zero, choose the scalars such that the **fewest**  $c_i$ s are nonzero. Since the  $\mathbf{v}_i$ s are eigenvectors and not the zero vector, at least two of the  $c_i$ s must be nonzero. Assume that  $c_j$  and  $c_k$  are nonzero.

Multiplying both sides of equation (1) by  $A$  gives

$$A(c_1\mathbf{v}_1 + \dots + c_j\mathbf{v}_j + \dots + c_p\mathbf{v}_p) = c_1\lambda_1\mathbf{v}_1 + \dots + c_j\lambda_j\mathbf{v}_j + \dots + c_p\lambda_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

# Linear Independence of Eigenvectors

## Theorem

If  $\mathbf{v}_1, \dots, \mathbf{v}_p$  are eigenvectors corresponding to **distinct** eigenvalues  $\lambda_1, \dots, \lambda_p$  of an  $n \times n$  matrix  $A$ , then the set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is **linearly independent**.

**Proof.** Suppose  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is linearly dependent. Thus there are scalars  $c_i$ , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such  $c_i$ s that are not all zero, choose the scalars such that the **fewest**  $c_i$ s are nonzero. Since the  $\mathbf{v}_i$ s are eigenvectors and not the zero vector, at least two of the  $c_i$ s must be nonzero. Assume that  $c_j$  and  $c_k$  are nonzero.

Multiplying both sides of equation (1) by  $A$  gives

$$A(c_1\mathbf{v}_1 + \dots + c_j\mathbf{v}_j + \dots + c_p\mathbf{v}_p) = c_1\lambda_1\mathbf{v}_1 + \dots + c_j\lambda_j\mathbf{v}_j + \dots + c_p\lambda_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

Multiplying both sides of equation (1) by  $\lambda_j$  and subtracting from equation (2) gives

$$c_1(\lambda_1 - \lambda_j)\mathbf{v}_1 + c_2(\lambda_2 - \lambda_j)\mathbf{v}_2 + \dots + 0\mathbf{v}_j + \dots + c_p(\lambda_p - \lambda_j)\mathbf{v}_p = \mathbf{0}. \quad (3)$$

# Linear Independence of Eigenvectors

## Theorem

If  $\mathbf{v}_1, \dots, \mathbf{v}_p$  are eigenvectors corresponding to **distinct** eigenvalues  $\lambda_1, \dots, \lambda_p$  of an  $n \times n$  matrix  $A$ , then the set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is **linearly independent**.

**Proof.** Suppose  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is linearly dependent. Thus there are scalars  $c_i$ , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such  $c_i$ s that are not all zero, choose the scalars such that the **fewest**  $c_i$ s are nonzero. Since the  $\mathbf{v}_i$ s are eigenvectors and not the zero vector, at least two of the  $c_i$ s must be nonzero. Assume that  $c_j$  and  $c_k$  are nonzero.

Multiplying both sides of equation (1) by  $A$  gives

$$A(c_1\mathbf{v}_1 + \dots + c_j\mathbf{v}_j + \dots + c_p\mathbf{v}_p) = c_1\lambda_1\mathbf{v}_1 + \dots + c_j\lambda_j\mathbf{v}_j + \dots + c_p\lambda_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

Multiplying both sides of equation (1) by  $\lambda_j$  and subtracting from equation (2) gives

$$c_1(\lambda_1 - \lambda_j)\mathbf{v}_1 + c_2(\lambda_2 - \lambda_j)\mathbf{v}_2 + \dots + 0\mathbf{v}_j + \dots + c_p(\lambda_p - \lambda_j)\mathbf{v}_p = \mathbf{0}. \quad (3)$$

Note that  $c_k(\lambda_k - \lambda_j) \neq 0$  since  $\lambda_k \neq \lambda_j$ . Thus, we have expressed  $\mathbf{0}$  as a linear combination of  $\mathbf{v}_1, \dots, \mathbf{v}_p$  with not all scalars 0 that has **fewer** zero scalars than what we assumed was the fewest.

**Contradiction!**

