

# The Matrix of a Linear Transformation

## Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit

<https://github.com/CU-Denver-MathStats-OER>

# The Matrix of a Linear Transformation

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  denote a linear transformation with

$$T \begin{pmatrix} [1] \\ [0] \\ [0] \end{pmatrix} = T(\mathbf{e}_1) = \begin{bmatrix} 3 \\ -1 \\ 1 \\ 0 \end{bmatrix}, \quad T \begin{pmatrix} [0] \\ [1] \\ [0] \end{pmatrix} = T(\mathbf{e}_2) = \begin{bmatrix} 0 \\ 2 \\ 1 \\ 3 \end{bmatrix}, \quad T \begin{pmatrix} [0] \\ [0] \\ [1] \end{pmatrix} = T(\mathbf{e}_3) = \begin{bmatrix} 1 \\ 0 \\ 1 \\ -1 \end{bmatrix}.$$

Find the image of an arbitrary vector  $\mathbf{x}$  in  $\mathbb{R}^3$ .

# The Matrix of a Linear Transformation

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  denote a linear transformation with

$$T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = T(\mathbf{e}_1) = \begin{bmatrix} 3 \\ -1 \\ 1 \\ 0 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) = T(\mathbf{e}_2) = \begin{bmatrix} 0 \\ 2 \\ 1 \\ 3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = T(\mathbf{e}_3) = \begin{bmatrix} 1 \\ 0 \\ 1 \\ -1 \end{bmatrix}.$$

Find the image of an arbitrary vector  $\mathbf{x}$  in  $\mathbb{R}^3$ .

$$\begin{aligned} T(\mathbf{x}) &= T(x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + x_3 \mathbf{e}_3) = x_1 T(\mathbf{e}_1) + x_2 T(\mathbf{e}_2) + x_3 T(\mathbf{e}_3) \\ &= x_1 \begin{bmatrix} 3 \\ -1 \\ 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 2 \\ 1 \\ 3 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 0 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 1 \\ -1 & 2 & 0 \\ 1 & 1 & 1 \\ 0 & 3 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = A\mathbf{x}. \end{aligned}$$

# Every Linear Transformation is a Matrix Transformation

## Theorem

For any linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ , there is a unique matrix  $A$  (called the **associated matrix for the linear transformation**) such that  $T(\mathbf{x}) = A\mathbf{x}$  for all  $\mathbf{x}$  in  $\mathbb{R}^n$ . The matrix  $A$  can be found by

$$A = [T(\mathbf{e}_1) \quad T(\mathbf{e}_2) \quad \dots \quad T(\mathbf{e}_n)].$$

# Every Linear Transformation is a Matrix Transformation

## Theorem

For any linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ , there is a unique matrix  $A$  (called the **associated matrix for the linear transformation**) such that  $T(\mathbf{x}) = A\mathbf{x}$  for all  $\mathbf{x}$  in  $\mathbb{R}^n$ . The matrix  $A$  can be found by

$$A = [T(\mathbf{e}_1) \quad T(\mathbf{e}_2) \quad \dots \quad T(\mathbf{e}_n)].$$

## Proof.



$$\begin{aligned} T(\mathbf{x}) &= T(x_1\mathbf{e}_1 + x_2\mathbf{e}_2 + \dots + x_n\mathbf{e}_n) = x_1 T(\mathbf{e}_1) + x_2 T(\mathbf{e}_2) + \dots + x_n T(\mathbf{e}_n) \\ &= [T(\mathbf{e}_1) \quad T(\mathbf{e}_2) \quad \dots \quad T(\mathbf{e}_n)] \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = A\mathbf{x}. \end{aligned}$$



# Geometric Interpretation in $\mathbb{R}^2$

Consider the linear transformation given by  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2 : \mathbf{x} \mapsto 0.5\mathbf{x}$ . Find the associated matrix  $A$  for this linear transformation.



We have  $T(\mathbf{e}_1) = \begin{bmatrix} 0.5 \\ 0 \end{bmatrix}$  and  $T(\mathbf{e}_2) = \begin{bmatrix} 0 \\ 0.5 \end{bmatrix}$  giving the matrix  $A = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix}$ .



# Geometric Interpretation in $\mathbb{R}^2$

Consider the linear transformation given by  $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x}$ .



We have  $T(\mathbf{e}_1) = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$  and  $T(\mathbf{e}_2) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  giving the geometric interpretation seen below.



# Contractions and Expansions



Consider the linear transformation given by  $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \mathbf{x}$ .

We have  $T(\mathbf{e}_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $T(\mathbf{e}_2) = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$  giving the geometric interpretation seen below.





# Shear Transformations

Consider the linear transformation given by  $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 1 & 0.75 \\ 0 & 1 \end{bmatrix} \mathbf{x}$ .



# Projections

Consider the linear transformation given by  $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \mathbf{x}$ .




## Rotations

Consider the linear transformation  $T$  of  $\mathbb{R}^2$  that rotates by the angle  $\theta$ . What is the associated matrix of  $T$ ?

## Rotations


Consider the linear transformation  $T$  of  $\mathbb{R}^2$  that rotates by the angle  $\theta$ . What is the associated matrix of  $T$ ?



$$T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \mathbf{x}$$

# Rotations

Consider the linear transformation  $T$  of  $\mathbb{R}^2$  that rotates by the angle  $\theta$ . What is the associated matrix of  $T$ ?



$$A\mathbf{e}_2 = (-\sin \theta, \cos \theta)$$

$$(0, 1) = \mathbf{e}_2$$

$$(\cos \theta, \sin \theta) = A\mathbf{e}_1$$

$$\mathbf{x}_1$$

$$(1, 0) = \mathbf{e}_1$$

$$T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \mathbf{x}$$

# Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. The mapping  $T$  is said to be:

- ▶ **One-to-one** if each  $\mathbf{b}$  in  $\mathbb{R}^m$  has **at most one**  $\mathbf{x}$  in  $\mathbb{R}^n$  such that  $T(\mathbf{x}) = \mathbf{b}$ .
- ▶ **Onto** if each  $\mathbf{b}$  in  $\mathbb{R}^m$  is the image of **at least one**  $\mathbf{x}$  in  $\mathbb{R}^n$  such that  $T(\mathbf{x}) = \mathbf{b}$ .

# Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. The mapping  $T$  is said to be:

- ▶ **One-to-one** if each  $\mathbf{b}$  in  $\mathbb{R}^m$  has **at most one**  $\mathbf{x}$  in  $\mathbb{R}^n$  such that  $T(\mathbf{x}) = \mathbf{b}$ .
- ▶ **Onto** if each  $\mathbf{b}$  in  $\mathbb{R}^m$  is the image of **at least one**  $\mathbf{x}$  in  $\mathbb{R}^n$  such that  $T(\mathbf{x}) = \mathbf{b}$ .

## Example

- ▶ The dilation transformation  $T(\mathbf{x}) = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$  is both one-to-one and onto.
- ▶ The projection transformation  $T(\mathbf{x}) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$  is neither one-to-one nor onto.

## Examples

Determine whether each of the transformations is one-to-one and/or onto.

$$T : \mathbb{R}^3 \rightarrow \mathbb{R}^2 : \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \mapsto \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$
$$R : \mathbb{R}^3 \rightarrow \mathbb{R}^3 : \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \mapsto \begin{bmatrix} -x_2 \\ -x_3 \\ -x_1 \end{bmatrix}$$
$$S : \mathbb{R}^2 \rightarrow \mathbb{R}^3 : \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} x_2 \\ 0 \\ x_1 \end{bmatrix}$$

## Theorem

A linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is **one-to-one** if and only if  $T(\mathbf{x}) = \mathbf{0}$  has only the **trivial solution**.

## Proof.

## Theorem

A linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is **one-to-one** if and only if  $T(\mathbf{x}) = \mathbf{0}$  has only the **trivial solution**.

## Proof.

The forward direction we prove using proof by contradiction. Let's assume that  $T$  is a one-to-one mapping, and that  $T(\mathbf{x}) = \mathbf{0}$  has a non-trivial solution  $\mathbf{v}$  in  $\mathbb{R}^n$ . Then  $T$  is not one-to-one since  $T(\mathbf{0}) = T(\mathbf{v}) = \mathbf{0}$ . Thus we have a contradiction.

## Theorem

A linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is **one-to-one** if and only if  $T(\mathbf{x}) = \mathbf{0}$  has only the **trivial solution**.

## Proof.

The forward direction we prove using proof by contradiction. Let's assume that  $T$  is a one-to-one mapping, and that  $T(\mathbf{x}) = \mathbf{0}$  has a non-trivial solution  $\mathbf{v}$  in  $\mathbb{R}^n$ . Then  $T$  is not one-to-one since  $T(\mathbf{0}) = T(\mathbf{v}) = \mathbf{0}$ . Thus we have a contradiction.

The reverse direction we prove using proof by contradiction as well. Suppose  $T(\mathbf{x}) = \mathbf{0}$  has only the trivial solution, and  $T$  is not one-to-one. This means there exist  $\mathbf{u} \neq \mathbf{v}$  both in  $\mathbb{R}^n$  with  $T(\mathbf{u}) = T(\mathbf{v}) = \mathbf{b}$  in  $\mathbb{R}^m$ .

## Theorem

A linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is **one-to-one** if and only if  $T(\mathbf{x}) = \mathbf{0}$  has only the **trivial solution**.

## Proof.

The forward direction we prove using proof by contradiction. Let's assume that  $T$  is a one-to-one mapping, and that  $T(\mathbf{x}) = \mathbf{0}$  has a non-trivial solution  $\mathbf{v}$  in  $\mathbb{R}^n$ . Then  $T$  is not one-to-one since  $T(\mathbf{0}) = T(\mathbf{v}) = \mathbf{0}$ . Thus we have a contradiction.

The reverse direction we prove using proof by contradiction as well. Suppose  $T(\mathbf{x}) = \mathbf{0}$  has only the trivial solution, and  $T$  is not one-to-one. This means there exist  $\mathbf{u} \neq \mathbf{v}$  both in  $\mathbb{R}^n$  with  $T(\mathbf{u}) = T(\mathbf{v}) = \mathbf{b}$  in  $\mathbb{R}^m$ . Therefore, we have

$$T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v}) = \mathbf{b} - \mathbf{b} = \mathbf{0}.$$

Since  $\mathbf{u} \neq \mathbf{v}$ , we have found a non-trivial solution such that  $T(\mathbf{u} - \mathbf{v}) = \mathbf{0}$ . Thus we have a contradiction. □

## Summary

A linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  with associated matrix  $A$  is **one-to-one** if and only if

- ▶  $T(\mathbf{x}) = \mathbf{0}$  has only the trivial solution.
- ▶ The solution set to  $A\mathbf{x} = \mathbf{0}$  has no free variables.
- ▶ The matrix  $A$  has a pivot in every column.

A linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  with associated matrix  $A$  is **onto** if and only if

- ▶ For any  $\mathbf{b}$  in  $\mathbb{R}^m$  there exists at least one  $\mathbf{x}$  in  $\mathbb{R}^n$  with  $A\mathbf{x} = \mathbf{b}$ .
- ▶ All vectors  $\mathbf{b}$  in  $\mathbb{R}^m$  can be written as a linear combination of the columns of  $A$ .
- ▶ The matrix  $A$  has a pivot in every row.
- ▶ **The columns of  $A$  span all of  $\mathbb{R}^m$ .**