

Change of Basis

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Change of Coordinate Mappings

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$. Suppose that $\mathbf{x} \in V$, and we have $[\mathbf{x}]_{\mathcal{B}}$. How can we find $[\mathbf{x}]_{\mathcal{C}}$?

Change of Coordinate Mappings

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$. Suppose that $\mathbf{x} \in V$, and we have $[\mathbf{x}]_{\mathcal{B}}$. How can we find $[\mathbf{x}]_{\mathcal{C}}$?

We have that $\mathbf{x} = d_1\mathbf{b}_1 + d_2\mathbf{b}_2 + \dots + d_n\mathbf{b}_n$. We want to write \mathbf{x} as a linear combin of $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$. Since \mathcal{C} is a basis, each \mathbf{b}_j is a linear combin of the elements of \mathcal{C} : $\mathbf{b}_j = a_{1j}\mathbf{c}_1 + a_{2j}\mathbf{c}_2 + \dots + a_{nj}\mathbf{c}_n$. Substituting and grouping the \mathbf{c}_i , we have

$$\mathbf{x} = \left(\sum_{j=1}^n a_{1j}d_j \right) \mathbf{c}_1 + \left(\sum_{j=1}^n a_{2j}d_j \right) \mathbf{c}_2 + \dots + \left(\sum_{j=1}^n a_{ij}d_j \right) \mathbf{c}_i + \dots + \left(\sum_{j=1}^n a_{nj}d_j \right) \mathbf{c}_n.$$

Change of Coordinate Mappings

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$. Suppose that $\mathbf{x} \in V$, and we have $[\mathbf{x}]_{\mathcal{B}}$. How can we find $[\mathbf{x}]_{\mathcal{C}}$?

We have that $\mathbf{x} = d_1\mathbf{b}_1 + d_2\mathbf{b}_2 + \dots + d_n\mathbf{b}_n$. We want to write \mathbf{x} as a linear combin of $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$. Since \mathcal{C} is a basis, each \mathbf{b}_j is a linear combin of the elements of \mathcal{C} : $\mathbf{b}_j = a_{1j}\mathbf{c}_1 + a_{2j}\mathbf{c}_2 + \dots + a_{nj}\mathbf{c}_n$. Substituting and grouping the \mathbf{c}_i , we have

$$\mathbf{x} = \left(\sum_{j=1}^n a_{1j}d_j \right) \mathbf{c}_1 + \left(\sum_{j=1}^n a_{2j}d_j \right) \mathbf{c}_2 + \dots + \left(\sum_{j=1}^n a_{ij}d_j \right) \mathbf{c}_i + \dots + \left(\sum_{j=1}^n a_{nj}d_j \right) \mathbf{c}_n.$$

$$\text{Thus, } [\mathbf{x}]_{\mathcal{C}} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nj} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} d_1 \\ d_2 \\ \vdots \\ d_n \end{bmatrix}$$

Change of Coordinate Mappings

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$. Suppose that $\mathbf{x} \in V$, and we have $[\mathbf{x}]_{\mathcal{B}}$. How can we find $[\mathbf{x}]_{\mathcal{C}}$?

We have that $\mathbf{x} = d_1\mathbf{b}_1 + d_2\mathbf{b}_2 + \dots + d_n\mathbf{b}_n$. We want to write \mathbf{x} as a linear combin of $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$. Since \mathcal{C} is a basis, each \mathbf{b}_j is a linear combin of the elements of \mathcal{C} : $\mathbf{b}_j = a_{1j}\mathbf{c}_1 + a_{2j}\mathbf{c}_2 + \dots + a_{nj}\mathbf{c}_n$. Substituting and grouping the \mathbf{c}_i , we have

$$\mathbf{x} = \left(\sum_{j=1}^n a_{1j}d_j \right) \mathbf{c}_1 + \left(\sum_{j=1}^n a_{2j}d_j \right) \mathbf{c}_2 + \dots + \left(\sum_{j=1}^n a_{ij}d_j \right) \mathbf{c}_i + \dots + \left(\sum_{j=1}^n a_{nj}d_j \right) \mathbf{c}_n.$$

$$\text{Thus, } [\mathbf{x}]_{\mathcal{C}} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nj} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} d_1 \\ d_2 \\ \vdots \\ d_n \end{bmatrix} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} & \dots & [\mathbf{b}_n]_{\mathcal{C}} \end{bmatrix} [\mathbf{x}]_{\mathcal{B}} = \underset{\mathcal{C} \leftarrow \mathcal{B}}{P} [\mathbf{x}]_{\mathcal{B}}.$$

Change of Basis Matrix

Theorem

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$ be bases for a vector space V . Then there is a unique $n \times n$ matrix $P_{\mathcal{C} \leftarrow \mathcal{B}}$ such that

$$P_{\mathcal{C} \leftarrow \mathcal{B}} [\mathbf{x}]_{\mathcal{B}} = [\mathbf{x}]_{\mathcal{C}}.$$

The map $\mathbf{z} \mapsto P_{\mathcal{C} \leftarrow \mathcal{B}} \mathbf{z}$ is a linear transformation from \mathbb{R}^n (\mathcal{B} -coordinates) to \mathbb{R}^n (\mathcal{C} -coordinates).

The columns of the **change of basis matrix from \mathcal{B} to \mathcal{C}** , denoted $P_{\mathcal{C} \leftarrow \mathcal{B}}$, are the \mathcal{C} -coordinate vectors in the basis \mathcal{B} .

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} & \dots & [\mathbf{b}_n]_{\mathcal{C}} \end{bmatrix}.$$

Example

Let $V = \mathbb{P}^2$ and consider two bases $\mathcal{B} = \{1, t, t^2\}$ and $\mathcal{C} = \{1, 1 + t, 1 + t + t^2\}$.
Write $\mathbf{x} = 3 - 2t + 7t^2$ in terms of the basis \mathcal{C} .

Example

Let $V = \mathbb{P}^2$ and consider two bases $\mathcal{B} = \{1, t, t^2\}$ and $\mathcal{C} = \{1, 1 + t, 1 + t + t^2\}$. Write $\mathbf{x} = 3 - 2t + 7t^2$ in terms of the basis \mathcal{C} .

We compute the change of basis matrix $P_{\mathcal{C} \leftarrow \mathcal{B}}$.

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} & \dots & [\mathbf{b}_n]_{\mathcal{C}} \end{bmatrix} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Example

Let $V = \mathbb{P}^2$ and consider two bases $\mathcal{B} = \{1, t, t^2\}$ and $\mathcal{C} = \{1, 1+t, 1+t+t^2\}$. Write $\mathbf{x} = 3 - 2t + 7t^2$ in terms of the basis \mathcal{C} .

We compute the change of basis matrix ${}_{\mathcal{C} \leftarrow \mathcal{B}} P$.

$${}_{\mathcal{C} \leftarrow \mathcal{B}} P = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} & \dots & [\mathbf{b}_n]_{\mathcal{C}} \end{bmatrix} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}.$$

We can then compute

$$[\mathbf{x}]_{\mathcal{C}} = {}_{\mathcal{C} \leftarrow \mathcal{B}} P [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ -2 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 \\ -9 \\ 7 \end{bmatrix}.$$

So $3 - 2t + 7t^2 = 5(1) - 9(1+t) + 7(1+t+t^2)$.

Example

Let $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ and $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be bases for a vector space V . Suppose we know that

$$\mathbf{a}_1 = 4\mathbf{b}_1 - \mathbf{b}_2, \quad \mathbf{a}_2 = -\mathbf{b}_1 + \mathbf{b}_2 + \mathbf{b}_3, \quad \text{and} \quad \mathbf{a}_3 = -\mathbf{b}_2 + 2\mathbf{b}_3$$

1. Find the change of basis matrix from \mathcal{A} to \mathcal{B} .

Example

Let $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ and $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be bases for a vector space V . Suppose we know that

$$\mathbf{a}_1 = 4\mathbf{b}_1 - \mathbf{b}_2, \quad \mathbf{a}_2 = -\mathbf{b}_1 + \mathbf{b}_2 + \mathbf{b}_3, \quad \text{and} \quad \mathbf{a}_3 = -\mathbf{b}_2 + 2\mathbf{b}_3$$

1. Find the change of basis matrix from \mathcal{A} to \mathcal{B} .

2. Find $[\mathbf{x}]_{\mathcal{B}}$ for $\mathbf{x} = 3\mathbf{a}_1 + 4\mathbf{a}_2 + \mathbf{a}_3$.

We know that

$$[\mathbf{a}_1]_{\mathcal{B}} = \begin{bmatrix} 4 \\ -1 \\ 0 \end{bmatrix}_{\mathcal{B}}, \quad [\mathbf{a}_2]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}_{\mathcal{B}}, \quad [\mathbf{a}_3]_{\mathcal{B}} = \begin{bmatrix} 0 \\ -1 \\ 2 \end{bmatrix}_{\mathcal{B}}$$

Thus we have

$$P_{\mathcal{B} \leftarrow \mathcal{A}} = \begin{bmatrix} [\mathbf{a}_1]_{\mathcal{B}} & [\mathbf{a}_2]_{\mathcal{B}} & [\mathbf{a}_3]_{\mathcal{B}} \end{bmatrix} = \begin{bmatrix} 4 & -1 & 0 \\ -1 & 1 & -1 \\ 0 & 1 & 2 \end{bmatrix}$$

Example

Let $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ and $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be bases for a vector space V . Suppose we know that

$$\mathbf{a}_1 = 4\mathbf{b}_1 - \mathbf{b}_2, \quad \mathbf{a}_2 = -\mathbf{b}_1 + \mathbf{b}_2 + \mathbf{b}_3, \quad \text{and} \quad \mathbf{a}_3 = -\mathbf{b}_2 + 2\mathbf{b}_3$$

1. Find the change of basis matrix from \mathcal{A} to \mathcal{B} .

2. Find $[\mathbf{x}]_{\mathcal{B}}$ for $\mathbf{x} = 3\mathbf{a}_1 + 4\mathbf{a}_2 + \mathbf{a}_3$.

We know that

$$[\mathbf{a}_1]_{\mathcal{B}} = \begin{bmatrix} 4 \\ -1 \\ 0 \end{bmatrix}_{\mathcal{B}}, \quad [\mathbf{a}_2]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}_{\mathcal{B}}, \quad [\mathbf{a}_3]_{\mathcal{B}} = \begin{bmatrix} 0 \\ -1 \\ 2 \end{bmatrix}_{\mathcal{B}}$$

Thus we have

$$P_{\mathcal{B} \leftarrow \mathcal{A}} = \begin{bmatrix} [\mathbf{a}_1]_{\mathcal{B}} & [\mathbf{a}_2]_{\mathcal{B}} & [\mathbf{a}_3]_{\mathcal{B}} \end{bmatrix} = \begin{bmatrix} 4 & -1 & 0 \\ -1 & 1 & -1 \\ 0 & 1 & 2 \end{bmatrix}$$

We have $\mathbf{x} = \begin{bmatrix} 3 \\ 4 \\ 1 \end{bmatrix}_{\mathcal{A}}$. Using our change of basis matrix we have

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 4 & -1 & 0 \\ -1 & 1 & -1 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \\ 1 \end{bmatrix}_{\mathcal{A}} = \begin{bmatrix} 8 \\ 0 \\ 6 \end{bmatrix}_{\mathcal{B}}.$$

Example in \mathbb{R}^n

Let $V = \mathbb{R}^2$ and consider two bases $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -6 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{C} = \left\{ \begin{bmatrix} 5 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \end{bmatrix} \right\}$.

Write $\begin{bmatrix} 1 \\ 6 \end{bmatrix}_{\mathcal{B}}$ in terms of the basis \mathcal{C} .

Example in \mathbb{R}^n

Let $V = \mathbb{R}^2$ and consider two bases $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -6 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{C} = \left\{ \begin{bmatrix} 5 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \end{bmatrix} \right\}$.

Write $\begin{bmatrix} 1 \\ 6 \end{bmatrix}_{\mathcal{B}}$ in terms of the basis \mathcal{C} .

We find $[\mathbf{b}_1]_{\mathcal{C}}$ and $[\mathbf{b}_2]_{\mathcal{C}}$ by solving

$$\begin{bmatrix} 5 & 2 \\ 3 & -2 \end{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

and

$$\begin{bmatrix} 5 & 2 \\ 3 & -2 \end{bmatrix} [\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} -6 \\ 1 \end{bmatrix}.$$

Example in \mathbb{R}^n

Let $V = \mathbb{R}^2$ and consider two bases $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -6 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{C} = \left\{ \begin{bmatrix} 5 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \end{bmatrix} \right\}$.

Write $\begin{bmatrix} 1 \\ 6 \end{bmatrix}_{\mathcal{B}}$ in terms of the basis \mathcal{C} .

We find $[\mathbf{b}_1]_{\mathcal{C}}$ and $[\mathbf{b}_2]_{\mathcal{C}}$ by solving

$$\begin{bmatrix} 5 & 2 \\ 3 & -2 \end{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

and

$$\begin{bmatrix} 5 & 2 \\ 3 & -2 \end{bmatrix} [\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} -6 \\ 1 \end{bmatrix}.$$

More efficient to solve **both** by row reducing

$$\begin{aligned} \left[\begin{array}{cc|cc} \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{b}_1 & \mathbf{b}_2 \end{array} \right] &= \begin{bmatrix} 5 & 2 & | & 2 & -6 \\ 3 & -2 & | & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & \frac{2}{5} & | & \frac{2}{5} & -\frac{6}{5} \\ 0 & -\frac{16}{5} & | & -\frac{1}{5} & \frac{23}{5} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & | & \frac{3}{16} & -\frac{10}{16} \\ 0 & 1 & | & -\frac{1}{16} & \frac{-23}{16} \end{bmatrix} \end{aligned}$$

Example in \mathbb{R}^n

Let $V = \mathbb{R}^2$ and consider two bases $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -6 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{C} = \left\{ \begin{bmatrix} 5 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \end{bmatrix} \right\}$.

Write $\begin{bmatrix} 1 \\ 6 \end{bmatrix}_{\mathcal{B}}$ in terms of the basis \mathcal{C} .

We find $[\mathbf{b}_1]_{\mathcal{C}}$ and $[\mathbf{b}_2]_{\mathcal{C}}$ by solving

$$\begin{bmatrix} 5 & 2 \\ 3 & -2 \end{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

and

$$\begin{bmatrix} 5 & 2 \\ 3 & -2 \end{bmatrix} [\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} -6 \\ 1 \end{bmatrix}.$$

More efficient to solve **both** by row reducing

$$\begin{aligned} [\mathbf{c}_1 \quad \mathbf{c}_2 \mid \mathbf{b}_1 \quad \mathbf{b}_2] &= \begin{bmatrix} 5 & 2 & \mid & 2 & -6 \\ 3 & -2 & \mid & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & \frac{2}{5} & \mid & \frac{2}{5} & -\frac{6}{5} \\ 0 & -\frac{16}{5} & \mid & -\frac{1}{5} & \frac{23}{5} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & \mid & \frac{3}{16} & -\frac{10}{16} \\ 0 & 1 & \mid & -\frac{1}{16} & \frac{23}{16} \end{bmatrix} \end{aligned}$$

$$[\mathbf{x}]_{\mathcal{C}} = {}_{\mathcal{C} \leftarrow \mathcal{B}} P [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} \frac{3}{16} & -\frac{10}{16} \\ -\frac{1}{16} & \frac{23}{16} \end{bmatrix} \begin{bmatrix} 1 \\ 6 \end{bmatrix}_{\mathcal{B}} = \begin{bmatrix} -27/8 \\ -137/16 \end{bmatrix}_{\mathcal{C}}.$$

Finding the Change of Basis Matrix for Two Bases of \mathbb{R}^n

Let $V = \mathbb{R}^n$ with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$.

(Note that \mathbf{b}_i and \mathbf{c}_j are written in terms of the *standard basis* for \mathbb{R}^n .)

Then the *change of basis matrix* $P_{\mathcal{C} \leftarrow \mathcal{B}}$ can be found by row reducing

$$\left[\begin{array}{cccc|cccc} \mathbf{c}_1 & \mathbf{c}_2 & \dots & \mathbf{c}_n & \mathbf{b}_1 & \mathbf{b}_2 & \dots & \mathbf{b}_n \end{array} \right] \rightarrow \left[\begin{array}{cccc|cccc} I_n & & & & & & & P_{\mathcal{C} \leftarrow \mathcal{B}} \end{array} \right].$$

Note this is reminiscent of computing A^{-1} by row reducing $[A \mid I] \rightarrow [I \mid A^{-1}]$.

Inverse of Change of Basis Matrix

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$.

Let $P_{\mathcal{C} \leftarrow \mathcal{B}}$ be the change of basis matrix. Is $P_{\mathcal{C} \leftarrow \mathcal{B}}$ invertible?

Inverse of Change of Basis Matrix

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$.

Let $P_{\mathcal{C} \leftarrow \mathcal{B}}$ be the change of basis matrix. Is $P_{\mathcal{C} \leftarrow \mathcal{B}}$ invertible?

Recall that the \mathcal{C} -coordinate mapping $V \rightarrow \mathbb{R}^n$ where $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{C}}$ is an isomorphism.

Since $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ is a linearly independent set, so is $\{[\mathbf{b}_1]_{\mathcal{C}}, [\mathbf{b}_2]_{\mathcal{C}}, \dots, [\mathbf{b}_n]_{\mathcal{C}}\}$.

Thus the cols of $P_{\mathcal{C} \leftarrow \mathcal{B}}$ are linearly indep, and by the Invertible Matrix Theorem $P_{\mathcal{C} \leftarrow \mathcal{B}}$ is invertible.

Inverse of Change of Basis Matrix

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$.

Let $P_{\mathcal{C} \leftarrow \mathcal{B}}$ be the change of basis matrix. Is $P_{\mathcal{C} \leftarrow \mathcal{B}}$ invertible?

Recall that the \mathcal{C} -coordinate mapping $V \rightarrow \mathbb{R}^n$ where $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{C}}$ is an isomorphism.

Since $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ is a linearly independent set, so is $\{[\mathbf{b}_1]_{\mathcal{C}}, [\mathbf{b}_2]_{\mathcal{C}}, \dots, [\mathbf{b}_n]_{\mathcal{C}}\}$.

Thus the cols of $P_{\mathcal{C} \leftarrow \mathcal{B}}$ are linearly indep, and by the Invertible Matrix Theorem $P_{\mathcal{C} \leftarrow \mathcal{B}}$ is invertible.

What is $P_{\mathcal{C} \leftarrow \mathcal{B}}^{-1}$?

Inverse of Change of Basis Matrix

Let V be a vector space with two bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n\}$.

Let $P_{\mathcal{C} \leftarrow \mathcal{B}}$ be the change of basis matrix. Is $P_{\mathcal{C} \leftarrow \mathcal{B}}$ invertible?

Recall that the \mathcal{C} -coordinate mapping $V \rightarrow \mathbb{R}^n$ where $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{C}}$ is an isomorphism.

Since $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ is a linearly independent set, so is $\{[\mathbf{b}_1]_{\mathcal{C}}, [\mathbf{b}_2]_{\mathcal{C}}, \dots, [\mathbf{b}_n]_{\mathcal{C}}\}$.

Thus the cols of $P_{\mathcal{C} \leftarrow \mathcal{B}}$ are linearly indep, and by the Invertible Matrix Theorem $P_{\mathcal{C} \leftarrow \mathcal{B}}$ is invertible.

What is $P_{\mathcal{C} \leftarrow \mathcal{B}}^{-1}$? Since $P_{\mathcal{C} \leftarrow \mathcal{B}}[\mathbf{x}]_{\mathcal{B}} = [\mathbf{x}]_{\mathcal{C}}$, $[\mathbf{x}]_{\mathcal{B}} = P_{\mathcal{C} \leftarrow \mathcal{B}}^{-1}[\mathbf{x}]_{\mathcal{C}}$. Thus, $P_{\mathcal{C} \leftarrow \mathcal{B}}^{-1} = P_{\mathcal{B} \leftarrow \mathcal{C}}$.

Example

Let $V = \mathbb{P}^2$ and consider two bases $\mathcal{B} = \{1, t, t^2\}$ and $\mathcal{C} = \{1, 1 + t, 1 + t + t^2\}$.

Find the change of basis matrix $\underset{\mathcal{C} \leftarrow \mathcal{B}}{P}$.

Example

Let $V = \mathbb{P}^2$ and consider two bases $\mathcal{B} = \{1, t, t^2\}$ and $\mathcal{C} = \{1, 1 + t, 1 + t + t^2\}$.

Find the change of basis matrix $\overset{\textcolor{red}{P}}{\underset{\textcolor{red}{\mathcal{C} \leftarrow \mathcal{B}}}{}}$.

We compute the change of basis matrix $\overset{\textcolor{blue}{P}}{\underset{\textcolor{blue}{\mathcal{B} \leftarrow \mathcal{C}}}{}}$:

$$\overset{\textcolor{blue}{P}}{\underset{\textcolor{blue}{\mathcal{B} \leftarrow \mathcal{C}}}{}} = \begin{bmatrix} [1]_{\mathcal{B}} & [1+t]_{\mathcal{B}} & [1+t+t^2]_{\mathcal{B}} \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Now we can compute

$$\overset{\textcolor{red}{P}}{\underset{\textcolor{red}{\mathcal{C} \leftarrow \mathcal{B}}}{}} = \overset{\textcolor{blue}{P}}{\underset{\textcolor{blue}{\mathcal{B} \leftarrow \mathcal{C}}}{}}^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Matrix Vector Space

Let V denote the vector space of **symmetric 2×2 matrices**. We have a basis

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}. \text{ Consider the set } \mathcal{C} = \left\{ \begin{bmatrix} 1 & -2 \\ -2 & 2 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 3 & -1 \\ -1 & 1 \end{bmatrix} \right\}$$

1. Show that \mathcal{C} is a basis for V .

Matrix Vector Space

Let V denote the vector space of **symmetric 2×2 matrices**. We have a basis

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}. \text{ Consider the set } \mathcal{C} = \left\{ \begin{bmatrix} 1 & -2 \\ -2 & 2 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 3 & -1 \\ -1 & 1 \end{bmatrix} \right\}$$

1. Show that \mathcal{C} is a basis for V .

We know that

$$[\mathbf{c}_1]_{\mathcal{B}} = \begin{bmatrix} 1 \\ -2 \\ 2 \end{bmatrix}_{\mathcal{B}}, [\mathbf{c}_2]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 2 \\ 4 \end{bmatrix}_{\mathcal{B}}, [\mathbf{c}_3]_{\mathcal{B}} = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}_{\mathcal{B}}$$

The change of basis matrix below is invertible,
so \mathcal{C} is also a basis.

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = \begin{bmatrix} 1 & 0 & 3 \\ -2 & 2 & -1 \\ 2 & 4 & 1 \end{bmatrix}$$

Matrix Vector Space

Let V denote the vector space of **symmetric 2×2 matrices**. We have a basis

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}. \text{ Consider the set } \mathcal{C} = \left\{ \begin{bmatrix} 1 & -2 \\ -2 & 2 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 3 & -1 \\ -1 & 1 \end{bmatrix} \right\}$$

1. Show that \mathcal{C} is a basis for V .

We know that

$$[\mathbf{c}_1]_{\mathcal{B}} = \begin{bmatrix} 1 \\ -2 \\ 2 \end{bmatrix}_{\mathcal{B}}, [\mathbf{c}_2]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 2 \\ 4 \end{bmatrix}_{\mathcal{B}}, [\mathbf{c}_3]_{\mathcal{B}} = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}_{\mathcal{B}}$$

2. Write the matrix corresponding to $\begin{bmatrix} 2 \\ 1 \\ -5 \end{bmatrix}_{\mathcal{C}}$ in \mathcal{B} coordinates.

The change of basis matrix below is invertible, so \mathcal{C} is also a basis.

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = \begin{bmatrix} 1 & 0 & 3 \\ -2 & 2 & -1 \\ 2 & 4 & 1 \end{bmatrix}$$

Matrix Vector Space

Let V denote the vector space of **symmetric 2×2 matrices**. We have a basis

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}. \text{ Consider the set } \mathcal{C} = \left\{ \begin{bmatrix} 1 & -2 \\ -2 & 2 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 3 & -1 \\ -1 & 1 \end{bmatrix} \right\}$$

1. Show that \mathcal{C} is a basis for V .

We know that

$$[\mathbf{c}_1]_{\mathcal{B}} = \begin{bmatrix} 1 \\ -2 \\ 2 \end{bmatrix}_{\mathcal{B}}, [\mathbf{c}_2]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 2 \\ 4 \end{bmatrix}_{\mathcal{B}}, [\mathbf{c}_3]_{\mathcal{B}} = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}_{\mathcal{B}}$$

The change of basis matrix below is invertible,
so \mathcal{C} is also a basis.

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = \begin{bmatrix} 1 & 0 & 3 \\ -2 & 2 & -1 \\ 2 & 4 & 1 \end{bmatrix}$$

2. Write the matrix corresponding to $\begin{bmatrix} 2 \\ 1 \\ -5 \end{bmatrix}_{\mathcal{C}}$
in \mathcal{B} coordinates.

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 1 & 0 & 3 \\ -2 & 2 & -1 \\ 2 & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ -5 \end{bmatrix}_{\mathcal{C}} = \begin{bmatrix} -13 \\ 3 \\ 3 \end{bmatrix}_{\mathcal{B}}$$

which is the matrix $\begin{bmatrix} -13 & 3 \\ 3 & 3 \end{bmatrix}$.