

Linear Independence and Bases

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Finding an Efficient Set of Vectors that Span V

Consider the vector space $V = \mathbb{R}^3$ with the usual operations, and consider the following set of vectors:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}.$$

1. Do the set of vectors above span all of \mathbb{R}^3 ?

Finding an Efficient Set of Vectors that Span V

Consider the vector space $V = \mathbb{R}^3$ with the usual operations, and consider the following set of vectors:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}.$$

1. Do the set of vectors above span all of \mathbb{R}^3 ?

YES. Given any vector $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$ in \mathbb{R}^3 , we have $\mathbf{b} = b_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + b_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + b_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$, so \mathbf{b} is in the span of the set of vectors above.

Finding an Efficient Set of Vectors that Span V

Consider the vector space $V = \mathbb{R}^3$ with the usual operations, and consider the following set of vectors:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}.$$

1. Do the set of vectors above span all of \mathbb{R}^3 ?

YES. Given any vector $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$ in \mathbb{R}^3 , we have $\mathbf{b} = b_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + b_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + b_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$, so \mathbf{b} is in the span of the set of vectors above.

2. Do we need all five vectors in the set to span all of \mathbb{R}^3 ?

Finding an Efficient Set of Vectors that Span V

Consider the vector space $V = \mathbb{R}^3$ with the usual operations, and consider the following set of vectors:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}.$$

1. Do the set of vectors above span all of \mathbb{R}^3 ? **YES**
2. Do we need all five vectors in the set to span all of \mathbb{R}^3 ?

NO. We only need the three standard column vectors to span \mathbb{R}^3 . Notice

$$\begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} = -2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

Finding an Efficient Set of Vectors that Span V

Consider the vector space $V = \mathbb{R}^3$ with the usual operations, and consider the following set of vectors:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}.$$

1. Do the set of vectors above span all of \mathbb{R}^3 ? **YES**
2. Do we need all five vectors in the set to span all of \mathbb{R}^3 ?

NO. We only need the three standard column vectors to span \mathbb{R}^3 . Notice

$$\begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} = -2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

- ▶ Removing these two vectors from the set above does not change the span.
- ▶ Then also removing one of the standard column vectors would affect the span of the set.
- ▶ We need a minimum of three vectors to span all of \mathbb{R}^3 .

Revisiting Linear Independence

Let V denote a vector space, and consider the set of vectors $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$.

- ▶ The set S of vectors is **linearly independent** if the vector equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}$ has **only the trivial solution**.
- ▶ The set S of vectors is **linearly dependent** if the vector equation above has a **non-trivial solution**.

Revisiting Linear Independence

Let V denote a vector space, and consider the set of vectors $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$.

- ▶ The set S of vectors is **linearly independent** if the vector equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}$ has **only the trivial solution**.
- ▶ The set S of vectors is **linearly dependent** if the vector equation above has a **non-trivial solution**.

Theorem

A set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$ of vectors is **linearly dependent** if and only if some \mathbf{v}_j is a linear combination of the other vectors.

Basis of a Vector Space

Let H denote a subspace of a vector space V . A set of vectors \mathcal{B} in V is **basis for H** if:

- (i) \mathcal{B} is a **linearly independent set**, and
- (ii) the subspace spanned by \mathcal{B} equals H . Namely **$\text{Span } \mathcal{B} = H$** .

Basis of a Vector Space

Let H denote a subspace of a vector space V . A set of vectors \mathcal{B} in V is **basis for H** if:

- (i) \mathcal{B} is a **linearly independent set**, and
- (ii) the subspace spanned by \mathcal{B} equals H . Namely **$\text{Span } \mathcal{B} = H$** .

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}$$

NOT A BASIS

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

NOT A BASIS

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

WE HAVE A WINNER!

Basis of a Vector Space

Let H denote a subspace of a vector space V . A set of vectors \mathcal{B} in V is **basis for H** if:

- (i) \mathcal{B} is a **linearly independent set**, and
- (ii) the subspace spanned by \mathcal{B} equals H . Namely **$\text{Span } \mathcal{B} = H$** .

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}$$

NOT A BASIS

Note that there is **not** a unique basis.

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}$$

ALSO A BASIS

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

WE HAVE A WINNER!

Spanning Set Theorem

Theorem

Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$ be a set in a vector space V , and let $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$ (which is a subspace of V).

- ▶ If one of the vectors in S is a linear combination of the remaining vectors in S , then after removing that vector from the set, the remaining vectors will still span H .
- ▶ If $H \neq \{\mathbf{0}\}$, some nonempty subset of S is a basis for H .

Example

$$H = \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ 5 \end{bmatrix} \right\}$$

Standard Basis for \mathbb{R}^n

We saw that $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is a basis for \mathbb{R}^3 , and probably this is the most natural basis. We extend this basis to higher dimensions.

The set of standard column vectors $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ that forms a basis for \mathbb{R}^n is called the **standard basis for \mathbb{R}^n** .

Examples in \mathbb{R}^3

Determine whether the given set of vectors is a basis for \mathbb{R}^3 .

$$\left\{ \begin{bmatrix} 3 \\ 0 \\ -6 \end{bmatrix}, \begin{bmatrix} -4 \\ 1 \\ 7 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 5 \end{bmatrix} \right\}$$

Examples in \mathbb{R}^3

Determine whether the given set of vectors is a basis for \mathbb{R}^3 .

$$\left\{ \begin{bmatrix} 3 \\ 0 \\ -6 \end{bmatrix}, \begin{bmatrix} -4 \\ 1 \\ 7 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 5 \end{bmatrix} \right\}$$

$$\begin{bmatrix} 3 & -4 & -2 \\ 0 & 1 & 1 \\ -6 & 7 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\det \begin{bmatrix} 3 & -4 & -2 \\ 0 & 1 & 1 \\ -6 & 7 & 7 \end{bmatrix} = 6$$

Examples in \mathbb{R}^3

Determine whether the given set of vectors is a basis for \mathbb{R}^3 .

$$\left\{ \begin{bmatrix} 3 \\ 0 \\ -6 \end{bmatrix}, \begin{bmatrix} -4 \\ 1 \\ 7 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 5 \end{bmatrix} \right\}$$

$$\left\{ \begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 6 \\ 16 \\ -5 \end{bmatrix} \right\}$$

$$\begin{bmatrix} 3 & -4 & -2 \\ 0 & 1 & 1 \\ -6 & 7 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\det \begin{bmatrix} 3 & -4 & -2 \\ 0 & 1 & 1 \\ -6 & 7 & 7 \end{bmatrix} = 6$$

Examples in \mathbb{R}^3

Determine whether the given set of vectors is a basis for \mathbb{R}^3 .

$$\left\{ \begin{bmatrix} 3 \\ 0 \\ -6 \end{bmatrix}, \begin{bmatrix} -4 \\ 1 \\ 7 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 5 \end{bmatrix} \right\}$$

$$\begin{bmatrix} 3 & -4 & -2 \\ 0 & 1 & 1 \\ -6 & 7 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\det \begin{bmatrix} 3 & -4 & -2 \\ 0 & 1 & 1 \\ -6 & 7 & 7 \end{bmatrix} = 6$$

$$\left\{ \begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 6 \\ 16 \\ -5 \end{bmatrix} \right\}$$

$$\begin{bmatrix} 0 & 2 & 6 \\ 2 & 2 & 16 \\ -1 & 0 & -5 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -5x_3 \\ -3x_3 \\ x_3 \end{bmatrix} = x_3 \begin{bmatrix} -5 \\ -3 \\ 1 \end{bmatrix}.$$

Setting $x_3 = -1$ and rearranging, we see that

$$\begin{bmatrix} 6 \\ 16 \\ -5 \end{bmatrix} = 5 \begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix} + 3 \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}$$

Bases for More Abstract Vector Spaces

1. Find a basis for $\text{Mat}_{2 \times 2}$ the vector space of all 2×2 matrices with the usual operations.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + d \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

Thus we have $\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$.

Bases for More Abstract Vector Spaces

1. Find a basis for $\text{Mat}_{2 \times 2}$ the vector space of all 2×2 matrices with the usual operations.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + d \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

Thus we have $\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$.

2. Find a basis for \mathbb{P}_4 , the vector space of all polynomials of degree at most 4 with the usual operations.

An arbitrary vector in \mathbb{P}_4 is given by $p(t) = a_4t^4 + a_3t^3 + a_2t^2 + a_1t + a_0$. Thus we have $\mathcal{B} = \{1, t, t^2, t^3, t^4\}$.

Polynomial Vector Space Example 1

Is $\{2t^2 + 1, t^2 - 1, t + 2\}$ a basis for \mathbb{P}_2 ? (\mathbb{P}_2 = polynomials of degree at most 2)

Polynomial Vector Space Example 1

Is $\{2t^2 + 1, t^2 - 1, t + 2\}$ a basis for \mathbb{P}_2 ? (\mathbb{P}_2 = polynomials of degree at most 2)

We solve $c_1(2t^2 + 1) + c_2(t^2 - 1) + c_3(t + 2) = 0$. This gives

$$(2c_1 + c_2)t^2 + c_3t + (c_1 - c_2) = 0$$

From the linear and constant terms respectively, we see that $c_3 = 0$ and $c_1 = c_2$. Then looking at the coefficient in front of the quadratic term, we have $2c_1 + c_2 = 2c_1 + c_1 = 3c_1 = 0$. Thus the only solution is the trivial solution, $c_1 = c_2 = c_3 = 0$. The set is linearly independent.

Note the standard basis for \mathbb{P}_2 is $\{t^2, t, 1\}$, which contains three vectors. We have a linearly independent set of the same number of vectors, thus this set is a basis for \mathbb{P}_2 .

Polynomial Vector Space Example 2

Is $\{2t^2 + t, t + 1, 2t^2 - 1\}$ a basis for \mathbb{P}_2 ?

Polynomial Vector Space Example 2

Is $\{2t^2 + t, t + 1, 2t^2 - 1\}$ a basis for \mathbb{P}_2 ?

Notice that we have $2t^2 - 1 = (2t^2 + t) - (t + 1)$, or $\mathbf{v}_3 = \mathbf{v}_1 - \mathbf{v}_2$. Since the set is linearly dependent, it cannot be a basis for \mathbb{P}_2 .

Matrix Vector Space Example

Is $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 2 & -4 \\ 4 & 0 \end{bmatrix} \right\}$ a basis for $\text{Mat}_{2 \times 2}$?

Matrix Vector Space Example

Is $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 2 & -4 \\ 4 & 0 \end{bmatrix} \right\}$ a basis for $\text{Mat}_{2 \times 2}$?

Notice that we have

$$\begin{bmatrix} 2 & -4 \\ 4 & 0 \end{bmatrix} = 2 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} - 4 \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Since one matrix can be written as a linear combination of other matrices in the set, this is a **linearly dependent set**, and it cannot be a basis for $\text{Mat}_{2 \times 2}$.

Bases for the Null Space of a Matrix

The **null space** of an $m \times n$ matrix A , denoted **Null A** , is the set of all solutions to the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

Let $A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}$.

Find a basis for **Null A** .

Bases for the Null Space of a Matrix

The **null space** of an $m \times n$ matrix A , denoted **Null** A , is the set of all solutions to the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}. \quad \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Find a basis for **Null** A .

$$\mathbf{x} = \begin{bmatrix} -4x_2 - 2x_4 \\ x_2 \\ x_4 \\ x_4 \\ 0 \end{bmatrix} = x_2 \begin{bmatrix} -4 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -2 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} \quad \mathcal{B} = \left\{ \begin{bmatrix} -4 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} \right\}.$$

Basis for Column Space of a Matrix

The **column space** of an $m \times n$ matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$, denoted **Col** A , is the set of all linear combinations of the columns of A .

Find a basis for **Col** A .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Basis for Column Space of a Matrix

The **column space** of an $m \times n$ matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$, denoted **Col** A , is the set of all linear combinations of the columns of A .

Find a basis for **Col** A .

Note that $A\mathbf{y} = \mathbf{0} \iff R\mathbf{y} = \mathbf{0}$.

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Basis for Column Space of a Matrix

The **column space** of an $m \times n$ matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$, denoted **Col** A , is the set of all linear combinations of the columns of A .

Find a basis for **Col** A .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Note that $A\mathbf{y} = \mathbf{0} \iff R\mathbf{y} = \mathbf{0}$.

So every linear dep of the columns of R is a linear dep of the columns of A .

Basis for Column Space of a Matrix

The **column space** of an $m \times n$ matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$, denoted **Col** A , is the set of all linear combinations of the columns of A .

Find a basis for **Col** A .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Note that $A\mathbf{y} = \mathbf{0} \iff R\mathbf{y} = \mathbf{0}$.

So every linear dep of the columns of R is a linear dep of the columns of A .

Pivot cols of R are \mathbf{e}_i s; non-pivot cols are linear combs.

Basis for Column Space of a Matrix

The **column space** of an $m \times n$ matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$, denoted **Col** A , is the set of all linear combinations of the columns of A .

Find a basis for **Col** A .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$\begin{aligned} R = \\ \text{RREF}(A) = \end{aligned} \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Note that $A\mathbf{y} = \mathbf{0} \iff R\mathbf{y} = \mathbf{0}$.

So every linear dep of the columns of R is a linear dep of the columns of A .

Pivot cols of R are \mathbf{e}_i s; non-pivot cols are linear combs.

So **pivot cols of** A are lin indep, form a basis for **Col** A .

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 3 \\ 2 \\ 5 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -1 \\ 5 \\ 2 \\ 8 \end{bmatrix} \right\}.$$

Basis for Column Space of a Matrix

$$A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}$$

Given \mathbf{b} in Col A , how can \mathbf{b} be written as a linear combination of vectors in \mathcal{B} ?

Basis for Column Space of a Matrix

$$A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}$$

Given \mathbf{b} in Col A , how can \mathbf{b} be written as a linear combination of vectors in \mathcal{B} ?

Solve $A\mathbf{x} = \mathbf{b}$, use the solution where the free variables are 0.

Basis for Column Space of a Matrix

$$A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}$$

Given \mathbf{b} in Col A , how can \mathbf{b} be written as a linear combination of vectors in \mathcal{B} ?

Solve $A\mathbf{x} = \mathbf{b}$, use the solution where the free variables are 0.

$$\mathbf{b} = (1, 12, 3, 17).$$

$$\begin{array}{l} \text{RREF of} \\ [A|\mathbf{b}] = \end{array} \begin{bmatrix} 1 & 4 & 0 & 2 & 0 & 3 \\ 0 & 0 & 1 & -1 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 3 - 4x_2 - 2x_4 \\ x_2 \\ -7 + x_4 \\ x_4 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ -7 \\ 0 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} -4 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -2 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}.$$

$$\text{Set } x_2 = x_4 = 0. \text{ Then } \mathbf{b} = \begin{bmatrix} 1 \\ 12 \\ 3 \\ 17 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 3 \\ 2 \\ 5 \end{bmatrix} - 7 \begin{bmatrix} 0 \\ 1 \\ 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} -1 \\ 5 \\ 2 \\ 8 \end{bmatrix}.$$

Basis for Row Space of a Matrix

The **row space** of an $m \times n$ matrix $A = \begin{bmatrix} \mathbf{r}_1 \\ \vdots \\ \mathbf{r}_m \end{bmatrix}$, denoted **Row A** , is the set of all linear combinations of the rows of A . $\text{Row } A = \text{Span} \{ \mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m \} = \text{Col } A^T$

Find a basis for **Row A** .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Basis for Row Space of a Matrix

The **row space** of an $m \times n$ matrix $A = \begin{bmatrix} \mathbf{r}_1 \\ \vdots \\ \mathbf{r}_m \end{bmatrix}$, denoted **Row A** , is the set of all linear combinations of the rows of A . $\text{Row } A = \text{Span}\{\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m\} = \text{Col } A^T$

Find a basis for **Row A** .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The rows of R are linear combinations of rows of A .

Thus, $\text{span rows } R \subseteq \text{span rows } A$.

Basis for Row Space of a Matrix

The **row space** of an $m \times n$ matrix $A = \begin{bmatrix} \mathbf{r}_1 \\ \vdots \\ \mathbf{r}_m \end{bmatrix}$, denoted **Row A** , is the set of all linear combinations of the rows of A . $\text{Row } A = \text{Span}\{\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m\} = \text{Col } A^T$

Find a basis for **Row A** .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The rows of R are linear combinations of rows of A .

Thus, $\text{span rows } R \subseteq \text{span rows } A$.

But since elementary row ops are **reversible**,
the rows of A are linear combinations of rows of R .

Thus, $\text{span rows } A \subseteq \text{span rows } R \Rightarrow \text{Row } A = \text{Row } R$.

Basis for Row Space of a Matrix

The **row space** of an $m \times n$ matrix $A = \begin{bmatrix} \mathbf{r}_1 \\ \vdots \\ \mathbf{r}_m \end{bmatrix}$, denoted **Row A** , is the set of all linear combinations of the rows of A . $\text{Row } A = \text{Span} \{ \mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_m \} = \text{Col } A^T$

Find a basis for **Row A** .

$$\text{Let } A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

$$R = \text{RREF}(A) = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The rows of R are linear combinations of rows of A .

Thus, $\text{span rows } R \subseteq \text{span rows } A$.

But since elementary row ops are **reversible**, the rows of A are linear combinations of rows of R .

Thus, $\text{span rows } A \subseteq \text{span rows } R \Rightarrow \text{Row } A = \text{Row } R$.

So a basis for **Row R** is a basis for **Row A** .

A basis for **Row A** is

$$\{(1, 4, 0, 2, 0), (0, 0, 1, -1, 0), (0, 0, 0, 0, 1)\}.$$

Basis for Row Space of a Matrix

Theorem

If two matrices A and B are row equivalent, then their **row spaces** are the same.

If B is in (reduced) row echelon form, the nonzero rows of B form a basis for the row space of A as well as for B .

Summary

Given a matrix A :

- ▶ Solve the equation $A\mathbf{x} = \mathbf{0}$ to find a basis for $\text{Null } A$.

Summary

Given a matrix A :

- ▶ Solve the equation $A\mathbf{x} = \mathbf{0}$ to find a basis for **Null A** .
- ▶ Find the reduced row echelon form of A .
- ▶ The pivot columns of the **original matrix A** form a basis for **Col A** .
- ▶ The non-zero rows of the **RREF matrix** form a basis for **Row A** .