

Properties of Determinants

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Cofactor Expansion Definition of Determinant

For an $n \times n$ matrix A , the cofactor expansion about row i is $\det A = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det A_{ij}$.

Cofactor Expansion Definition of Determinant

For an $n \times n$ matrix A , the cofactor expansion about row i is $\det A = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det A_{ij}$.

This recursive definition is too time-consuming for large matrices.

We will discover a more efficient method of calculating the determinant by studying how the elementary row operations change the determinant.

Scaling a Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

Scaling a Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -5 & 15 & 35 & 45 \\ 4 & -2 & -3 & 2 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Scaling a Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -5 & 15 & 35 & 45 \\ 4 & -2 & -3 & 2 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Let A and B be $n \times n$ matrices, and let k be a scalar.

If B is obtained by scaling one row of A by k , then $\det B = k(\det A)$.

Scaling a Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -5 & 15 & 35 & 45 \\ 4 & -2 & -3 & 2 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Let A and B be $n \times n$ matrices, and let k be a scalar.

If B is obtained by scaling one row of A by k , then $\det B = k(\det A)$.

Proof. Let A be an $n \times n$ matrix, and let B be the matrix obtained by multiplying the row i of A by the scalar k . We calculate the determinant of B by cofactor expansion across row i .

$$\det B = \sum_{j=1}^n (-1)^{i+j} b_{ij} \det B_{ij} = \sum_{j=1}^n (-1)^{i+j} (k \cdot a_{ij}) \det A_{ij} = k \sum_{j=1}^n (-1)^{i+j} a_{ij} \det A_{ij} = \det A.$$

Adding a Multiple of a Row to another Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

Adding a Multiple of a Row to another Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 15 & 28 & 35 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Adding a Multiple of a Row to another Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 15 & 28 & 35 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Let A and B be $n \times n$ matrices.

If B is obtained from A by adding a multiple of a row to another, then $\det B = \det A$.

Adding a Multiple of a Row to another Row

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 15 & 28 & 35 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Let A and B be $n \times n$ matrices.

If B is obtained from A by adding a multiple of a row to another, then $\det B = \det A$.

Proof. We'll give a proof later.

Swapping Two Rows

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

Swapping Two Rows

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 4 & -2 & -3 & 2 \\ -1 & 3 & 7 & 9 \\ 5 & 6 & 7 & 8 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Swapping Two Rows

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 4 & -2 & -3 & 2 \\ -1 & 3 & 7 & 9 \\ 5 & 6 & 7 & 8 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Let A and B be $n \times n$ matrices.

If B is obtained by **swapping two rows of A** , then **$\det B = -\det A$** .

Swapping Two Rows

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

$$\det I_4 = 1$$

$$\det E =$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ -1 & 3 & 7 & 9 \\ 4 & -2 & -3 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 & 4 \\ 4 & -2 & -3 & 2 \\ -1 & 3 & 7 & 9 \\ 5 & 6 & 7 & 8 \end{bmatrix}$$

$$\det A = -40$$

$$\det B =$$

Let A and B be $n \times n$ matrices.

If B is obtained by **swapping two rows of A** , then $\det B = -\det A$.

Proof. We'll give a proof later.

Elementary Row Operations and Determinants

Let A and B denote $n \times n$ matrices, and let k denote a scalar.

- ▶ If B is obtained from A by scaling one row of A by k , then $\det B = k(\det A)$.
- ▶ If B is obtained from A by adding a mult of a row to another, then $\det B = \det A$.
- ▶ If B is obtained from A by swapping two rows of A , then $\det B = -\det A$.

Elementary Row Operations and Determinants

Let A and B denote $n \times n$ matrices, and let k denote a scalar.

- ▶ If B is obtained from A by scaling one row of A by k , then $\det B = k(\det A)$.
- ▶ If B is obtained from A by adding a mult of a row to another, then $\det B = \det A$.
- ▶ If B is obtained from A by swapping two rows of A , then $\det B = -\det A$.

Proof.

Let A be an $n \times n$ matrix ($n \geq 2$), and let B denote the matrix obtained from A by adding k times row r to row s , OR swapping rows r and s .

Elementary Row Operations and Determinants

Let A and B denote $n \times n$ matrices, and let k denote a scalar.

- ▶ If B is obtained from A by scaling one row of A by k , then $\det B = k(\det A)$.
- ▶ If B is obtained from A by adding a mult of a row to another, then $\det B = \det A$.
- ▶ If B is obtained from A by swapping two rows of A , then $\det B = -\det A$.

Proof.

Let A be an $n \times n$ matrix ($n \geq 2$), and let B denote the matrix obtained from A by adding k times row r to row s , OR swapping rows r and s .

If $n = 2$, then we can directly calculate the determinant.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow \begin{bmatrix} a & b \\ c + ka & d + kb \end{bmatrix}$$

$$\det A = ad - bc \quad \begin{aligned} \det B &= a(d + kb) - b(c + ka) \\ &= ad - bc + kab - kab = ad - bc \end{aligned}$$

Elementary Row Operations and Determinants

Let A and B denote $n \times n$ matrices, and let k denote a scalar.

- ▶ If B is obtained from A by scaling one row of A by k , then $\det B = k(\det A)$.
- ▶ If B is obtained from A by adding a mult of a row to another, then $\det B = \det A$.
- ▶ If B is obtained from A by swapping two rows of A , then $\det B = -\det A$.

Proof.

Let A be an $n \times n$ matrix ($n \geq 2$), and let B denote the matrix obtained from A by adding k times row r to row s , OR swapping rows r and s .

If $n = 2$, then we can directly calculate the determinant.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow \begin{bmatrix} a & b \\ c + ka & d + kb \end{bmatrix}$$

$$\det A = ad - bc$$

$$\begin{aligned} \det B &= a(d + kb) - b(c + ka) \\ &= ad - bc + kab - kab = ad - bc \end{aligned}$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow \begin{bmatrix} c & d \\ a & b \end{bmatrix}$$

$$\det A = ad - bc$$

$$\det B = bc - ad = -(ad - bc)$$

Proof continued

Proof.

Let A be an $n \times n$ matrix ($n \geq 2$), and let B denote the matrix obtained from A by
adding k times row r to row s , OR swapping rows r and s .

Assume that $n > 2$. Let i be a row other than rows r and s .

We calculate the determinant of matrix B by doing a cofactor expansion across row i .

$$\det B = \sum_{j=1}^n (-1)^{i+j} b_{ij} \det B_{ij}$$

Proof continued

Proof.

Let A be an $n \times n$ matrix ($n \geq 2$), and let B denote the matrix obtained from A by adding k times row r to row s , OR swapping rows r and s .

Assume that $n > 2$. Let i be a row other than rows r and s .

We calculate the determinant of matrix B by doing a cofactor expansion across row i .

$$\det B = \sum_{j=1}^n (-1)^{i+j} b_{ij} \det B_{ij}$$

B_{ij} is A_{ij} with either a multiple of a row added to another or two rows swapped.

If a multiple added, then by induction $\det B_{ij} = \det A_{ij}$. So $\det B = \det A$.

Proof continued

Proof.

Let A be an $n \times n$ matrix ($n \geq 2$), and let B denote the matrix obtained from A by adding k times row r to row s , OR swapping rows r and s .

Assume that $n > 2$. Let i be a row other than rows r and s .

We calculate the determinant of matrix B by doing a cofactor expansion across row i .

$$\det B = \sum_{j=1}^n (-1)^{i+j} b_{ij} \det B_{ij}$$

B_{ij} is A_{ij} with either a multiple of a row added to another or two rows swapped.

If a multiple added, then by induction $\det B_{ij} = \det A_{ij}$. So $\det B = \det A$.

If two rows swapped, then by induction $\det B_{ij} = -\det A_{ij}$. So $\det B = -\det A$.



Example of Computing Determinant

Compute the determinant of $A = \begin{bmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{bmatrix}$.

Example of Computing Determinant

Compute the determinant of $A = \begin{bmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{bmatrix}$.

Row-reduce A , tracking how the elementary row ops change the determinant.

$$\begin{aligned} \det \begin{bmatrix} 1 & -4 & 2 \\ -2 & 8 & -9 \\ -1 & 7 & 0 \end{bmatrix} &= \det \begin{bmatrix} 1 & -4 & 2 \\ 0 & 0 & -5 \\ -1 & 7 & 0 \end{bmatrix} = \det \begin{bmatrix} 1 & -4 & 2 \\ 0 & 0 & -5 \\ 0 & 3 & 2 \end{bmatrix} \\ &= -\det \begin{bmatrix} 1 & -4 & 2 \\ 0 & 3 & 2 \\ 0 & 0 & -5 \end{bmatrix} = -3 \det \begin{bmatrix} 1 & -4 & 2 \\ 0 & 1 & \frac{2}{3} \\ 0 & 0 & -5 \end{bmatrix} \\ &= 15 \det \begin{bmatrix} 1 & -4 & 2 \\ 0 & 1 & \frac{2}{3} \\ 0 & 0 & 1 \end{bmatrix} = 15 \cdot 1 = 15 \end{aligned}$$

Example

A square matrix A is invertible if and only if $\det A \neq 0$.

Compute the determinant of $A = \begin{bmatrix} 3 & -1 & 2 & -5 \\ 0 & 5 & -3 & -6 \\ -6 & 7 & -7 & 4 \\ -5 & -8 & 0 & 9 \end{bmatrix}$.

Example

A square matrix A is invertible if and only if $\det A \neq 0$.

Compute the determinant of $A = \begin{bmatrix} 3 & -1 & 2 & -5 \\ 0 & 5 & -3 & -6 \\ -6 & 7 & -7 & 4 \\ -5 & -8 & 0 & 9 \end{bmatrix}$.

$$\det \begin{bmatrix} 3 & -1 & 2 & -5 \\ 0 & 5 & -3 & -6 \\ -6 & 7 & -7 & 4 \\ -5 & -8 & 0 & 9 \end{bmatrix} = \det \begin{bmatrix} 3 & -1 & 2 & -5 \\ 0 & 5 & -3 & -6 \\ 0 & 5 & -3 & -6 \\ -5 & -8 & 0 & 9 \end{bmatrix}$$

Since A is not invertible, we have $\det A = 0$.

Elementary Matrices

Scaling a row:

$$\det \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = 5$$

Add mult of row to another:

$$\det \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = 1$$

Swapping two rows:

$$\det \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} = -1$$

If E is an elementary matrix, then $\det(EA) = (\det E)(\det A)$.

Product of Square Matrices

Let A and B denote $n \times n$ matrices. If A is not invertible, then AB is not invertible.

We prove this statement using proof by contradiction.

Product of Square Matrices

Let A and B denote $n \times n$ matrices. If A is not invertible, then AB is not invertible.

We prove this statement using proof by contradiction.

Proof.

Suppose A is not invertible and AB is invertible. Since AB is invertible, there exists an inverse matrix M such that $(AB)M = I_n$.

Product of Square Matrices

Let A and B denote $n \times n$ matrices. If A is not invertible, then AB is not invertible.

We prove this statement using proof by contradiction.

Proof.

Suppose A is not invertible and AB is invertible. Since AB is invertible, there exists an inverse matrix M such that $(AB)M = I_n$. Since there exists a matrix $D = BM$ such that $A(BM) = I_n$, by the Invertible Matrix Theorem, it follows that A is invertible. Thus, we have a contradiction, so our original assumption must not be correct. Therefore, if A is not invertible, it must follow that AB is not invertible. \square

Product of Square Matrices

Let A and B denote $n \times n$ matrices. If A is not invertible, then AB is not invertible.

We prove this statement using proof by contradiction.

Proof.

Suppose A is not invertible and AB is invertible. Since AB is invertible, there exists an inverse matrix M such that $(AB)M = I_n$. Since there exists a matrix $D = BM$ such that $A(BM) = I_n$, by the Invertible Matrix Theorem, it follows that A is invertible. Thus, we have a contradiction, so our original assumption must not be correct. Therefore, if A is not invertible, it must follow that AB is not invertible. \square

If AB is invertible, then both A and B are invertible.

Determinant of a Product AB

Let A and B denote two $n \times n$ matrices. Then $\det(AB) = (\det A)(\det B)$.

Determinant of a Product AB

Let A and B denote two $n \times n$ matrices. Then $\det(AB) = (\det A)(\det B)$.

Proof.

Case 1: Suppose either A or B is not invertible. Then we just showed that AB is not invertible. In this case, $\det(AB) = 0$ and $(\det A)(\det B) = 0$, so the property holds.

Determinant of a Product AB

Let A and B denote two $n \times n$ matrices. Then $\det(AB) = (\det A)(\det B)$.

Proof.

Case 1: Suppose either A or B is not invertible. Then we just showed that AB is not invertible. In this case, $\det(AB) = 0$ and $(\det A)(\det B) = 0$, so the property holds.

Case 2: Suppose both A and B are invertible. This means A is row equivalent to I_n . Thus, we have $A = E_p \dots E_2 E_1 I_n$ where each E_i denotes an elementary matrix.

Determinant of a Product AB

Let A and B denote two $n \times n$ matrices. Then $\det(AB) = (\det A)(\det B)$.

Proof.

Case 1: Suppose either A or B is not invertible. Then we just showed that AB is not invertible. In this case, $\det(AB) = 0$ and $(\det A)(\det B) = 0$, so the property holds.

Case 2: Suppose both A and B are invertible. This means A is row equivalent to I_n . Thus, we have $A = E_p \dots E_2 E_1 I_n$ where each E_i denotes an elementary matrix. Therefore we have

$$\begin{aligned}\det(AB) &= \det(E_p \dots E_2 E_1 B) = \det(E_p) \det(E_{p-1} \dots E_2 E_1 B) \\ &= \det(E_p) \det(E_{p-1}) \det(E_{p-2} \dots E_2 E_1 B) \\ &= \dots \\ &= \det(E_p) \det(E_{p-1}) \dots (\det E_1)(\det B) \\ &= \det(E_p E_{p-1} \dots E_2 E_1)(\det B) \\ &= (\det A)(\det B)\end{aligned}$$



If A is an $n \times n$ invertible matrix with inverse A^{-1} , then $\det A^{-1} = \frac{1}{\det A}$.

Determinants of A and A^{-1}

If A is an $n \times n$ invertible matrix with inverse A^{-1} , then $\det A^{-1} = \frac{1}{\det A}$.

Proof.

Let A be an invertible matrix with inverse A^{-1} . This means $AA^{-1} = I_n$. Then we have $\det(AA^{-1}) = \det I_n = 1$. Using the previous result about the determinant for a product of matrices, we therefore have

$$1 = \det(AA^{-1}) = (\det A)(\det A^{-1}).$$

Therefore we have $\det A^{-1} = \frac{1}{\det A}$. □